

Shining Light on Black Rock Coatings in Smelter-Impacted Areas

Michael Schindler, Nathalie Mantha, Kurt T. Kyser, Mitsuhiro Murayama and Michael F. Hochella, Jr.

Volume 39, Number 3, 2012

URI: https://id.erudit.org/iderudit/geocan39_3art01

[See table of contents](#)

Publisher(s)

The Geological Association of Canada

ISSN

0315-0941 (print)

1911-4850 (digital)

[Explore this journal](#)


Cite this article

Schindler, M., Mantha, N., Kyser, K. T., Murayama, M. & Hochella, M. F. (2012). Shining Light on Black Rock Coatings in Smelter-Impacted Areas. *Geoscience Canada*, 39(3), 148–157.

Article abstract

Earth scientists have long known of the existence of black coatings on exposed rocks in smelter-impacted areas such as Sudbury, Ontario or Rouyn-Noranda, Québec. Black rock coatings in the Greater Sudbury area are remarkable geological records of atmospheric conditions, including mixing, scavenging, and oxidation processes, deposition rates, and the nature and source of anthropogenic releases to the atmosphere. The coatings are composed of an amorphous silica matrix that has trapped atmosphere-borne nanoparticles and has preserved their chemical and isotopic signature. These coatings are the product of high emissions of SO₂ and subsequent non-stoichiometric dissolution of exposed siliceous rocks. The coatings contain spherical smelter-derived Cu–Ni-oxide particulate matter (micrometre and nanometre-sized) and metal-sulphate-rich layers composed of nanometre aggregates of Fe–Cu sulphates. Lead, As, and Se-bearing nanoparticles emitted from smelters are incorporated in metal-sulphate-rich layers along the atmosphere-coating interface, presumably during coating formation. On a regional scale, ratios between different metal(loid)s in the coatings indicate that small diameter primary Pb, As and Se-bearing sulphate aerosols have been deposited at higher rates compared to larger, Ni-bearing particulate matter. High sulphur isotope values in coatings closer to smelting centres and their decrease with distance from the smelters is attributed to an increase in mixing of primary and secondary sulphates.

ARTICLE



Shining Light on Black Rock Coatings in Smelter-Impacted Areas

Michael Schindler^{1*}, Nathalie Mantha¹, Kurt T. Kyser², Mitsuhiro Murayama³, and Michael F. Hochella, Jr.⁴

¹ Department of Earth Sciences, Laurentian University, Sudbury, ON, Canada, P3E 2C6

² Department of Geological Sciences and Geological Engineering, Queen's University, Kingston, ON, Canada, K7L 3N6

³ Department of Materials Science and Engineering, Virginia Polytechnic Institute and State University, Blacksburg, VA, 24061, USA

⁴ Center for NanoBioEarth, Department of Geosciences, Virginia Polytechnic Institute and State University, Blacksburg, VA, 24061, USA

*Corresponding Author:
mschindler@laurentian.ca

SUMMARY

Earth scientists have long known of the existence of black coatings on exposed rocks in smelter-impacted areas such as Sudbury, Ontario or

Rouyn-Noranda, Québec. Black rock coatings in the Greater Sudbury area are remarkable geological records of atmospheric conditions, including mixing, scavenging, and oxidation processes, deposition rates, and the nature and source of anthropogenic releases to the atmosphere. The coatings are composed of an amorphous silica matrix that has trapped atmosphere-borne nanoparticles and has preserved their chemical and isotopic signature. These coatings are the product of high emissions of SO₂ and subsequent non-stoichiometric dissolution of exposed siliceous rocks. The coatings contain spherical smelter-derived Cu–Ni-oxide particulate matter (micrometre and nanometre-sized) and metal-sulphate-rich layers composed of nanometre aggregates of Fe–Cu sulphates. Lead, As, and Se-bearing nanoparticles emitted from smelters are incorporated in metal-sulphate-rich layers along the atmosphere-coating interface, presumably during coating formation. On a regional scale, ratios between different metal(loid)s in the coatings indicate that small diameter primary Pb, As and Se-bearing sulphate aerosols have been deposited at higher rates compared to larger, Ni-bearing particulate matter. High sulphur isotope values in coatings closer to smelting centres and their decrease with distance from the smelters is attributed to an increase in mixing of primary and secondary sulphates.

SOMMAIRE

Les géoscientifiques connaissent depuis longtemps l'existence d'une couche noire sur les roches exposées aux abords des fonderies comme celles de Sudbury en Ontario ou Rouyn-Noranda au Québec. Les couches noires des roches de la grande région de Sudbury

constituent de remarquables enregistrements géologiques des phénomènes atmosphériques, notamment des processus de mélange, de piégeage, et d'oxydation, ainsi que des taux de sédimentation et de la nature et de l'origine des rejets anthropiques dans l'atmosphère. Ces couches noires sont constituées d'une matrice de silice amorphe qui a piégé des nanoparticules atmosphériques et conservé leur signature chimique et isotopique. Ces couches noires sont le produit de fortes émissions atmosphériques de SO₂ et d'une dissolution non-stœchiométrique subséquente des roches siliceuses exposées. Ces couches noires contiennent des sphérules de particules atmosphériques d'oxydes de Cu–Ni (de taille micrométrique et nanométrique) issues de la fonderie, et des couches riches en sulfate de métaux constituées d'agrégats nanométriques de sulfates de Fe–Cu. Les nanoparticules de plomb, d'As et de Se émises par les fonderies sont incorporées dans les couches riches en sulfate de métal à l'interface de l'atmosphère et de cette couche, probablement lors de la formation de cette couche. À l'échelle régionale, les rapports de concentration des différents métaux ou métalloïdes dans les couches noires indiquent que les aérosols de faible diamètre de sulfate de Pb, d'As et de Se primaires ont été déposés à des taux plus élevés que les particules nickélicifères de plus grande dimension. Les valeurs plus élevées des isotopes du soufre observées dans les couches à proximité des fonderies et leur diminution en fonction de l'éloignement des fonderies sont attribuées à une augmentation du mélange entre sulfates à l'émission et post-émission.




Figure 1. (a) Map of the Greater Sudbury area with the location of the 30 sampling sites. The smelter centres at Copper Cliff, Coniston and Falconbridge are indicated with a sketched chimney; an index in the upper left corner shows the location of Sudbury within Ontario, Canada. Triangles and circles are sampling locations affected primarily by proximity to the Coniston and Falconbridge smelters, respectively. Areas dominantly affected by the Copper Cliff smelter are shown as stars (Copper Cliff), squares (Sudbury Centre) and crosses (Sudbury South). (b) Black coatings of the Sudbury area developed on exposed rocks. (c) A cross-section of black rock coating; amorphous silica matrix, embedded detrital amphiboles, a spherical smelter-derived particle and the metal-sulphate-rich layer are labelled 1, 2, 3 and 4, respectively.

INTRODUCTION

The Greater Sudbury area in Ontario, Canada, is one of the major base-metal smelting regions in North America. Three major Cu–Ni smelters (Copper Cliff, Coniston and Falconbridge; Fig. 1a) operated in Sudbury during the last century and annually emitted over 2 million tons of SO₂ and over 1000 tons of Cu, Ni, Pb and As during peak years of emissions. Logging of area forests prior to, and during, the mining activities, in conjunction with emissions of SO₂ and metal(loid)s, resulted in loss of vegetation in the Greater Sudbury area and development of large areas of barren rock and soil. These conditions yielded a landscape characterized by thin, acidic soils and black rock coatings (Gundermann and Hutchinson 1995). The black rock coatings of the Greater Sudbury area (Fig. 1b), which have been observed since the first half of the 20th century, are the product of atmosphere–rock interactions on exposed surfaces and infill fractures (Mantha et al. 2012a, b).

Inorganic rock coatings occur around the world (Dorn 1998); thick coatings are commonly observed in deserts (Fisk 1971; Smith and Whalley 1988), volcanic weathering environments (Curtiss et al. 1985; Minitti et al. 2007; Schindler et al. 2010), mine tailings (Schindler et al. 2009; Durocher and Schindler 2011), Antarctica (Friedmann and Weed 1987; Weed and Norton 1991; Giorgetti and Baroni 2007) and glaciated environments (Whalley et al. 1990). They contain a variety of environmental products, such as dust particulate matter, *in situ*-formed minerals, complex organic molecules and microfossils (Langworthy et al. 2010).

Surface coatings on rocks and building stones are useful forensic tools for monitoring recent and past air pollutants and tracing their sources. Coatings formed in deserts, in proximity to smelters and in highly polluted urban environments, may capture common pollutants such as metal-bearing particulate matter, sulphate aerosols and radionuclides during their forma-

tion (Nord et al. 1994; Hodge et al. 2005; Wayne et al. 2006; Smith and Prikryl 2007; Hoar et al. 2011; Mantha et al. 2012a, b). For example, rock varnishes collected around the Nevada Test Site and near power plants in Nevada contain anomalously low ²⁴⁰Pu/²³⁹Pu mass ratios in comparison to observed ratios worldwide (Hoar et al. 2011). Gypsum coatings on building stones in Prague record a change in the sulphur isotope composition from the centre to the periphery of the city, which was initially explained by a sulphur-isotope (S-isotope) fractionation process (Buzek and Šrámek 1985), but has been recently interpreted as a mixing process of primary and secondary sulphate aerosols (Mantha et al. 2012b).

The forensic examination of rock coatings is also valuable in providing an understanding of past terrestrial or planetary environments (e.g. on Mars; McLennan 2003; Milliken et al. 2008; Squyres et al. 2008), because coatings may preserve particulate mat-

ter and mineral phases, containing information about past climatic conditions, volcanic events, and biological activity.

The areas surrounding base-metal smelters are ideal locations for the study of past atmospheric particulate matter (including aerosols) because the release of vast quantities of SO_2 resulted in increased chemical weathering and ideal conditions for the formation of thick coatings on exposed rock surfaces. The chemical, mineralogical and isotopic composition of these coatings can be compared with historical data on metal emissions, metal speciation in smelter plumes, the isotopic composition of the smelted ore and smelter gases, and metal concentrations in the surrounding soils. The known point sources of SO_2 and metal particulate matter, as well as the available historical records of local emissions, represent clear advantages in the study of coatings compared to studies of coatings proximal to volcanoes (e.g. Schiffmann et al. 2006), in deserts (e.g. Perry et al. 2006) and on Antarctica (Giorgetti and Baroni 2007), which are all affected by various sources, inputs and processes potentially on a global scale.

This paper demonstrates that silica coatings in the vicinity of base-metal smelters record the mineralogical, chemical and isotopic signature of atmospheric nano- to micro-size particulate matter, as well as changes in the proportion of secondary and primary aerosols with distance from the smelter point-source.

MATERIAL AND METHODS

Black rock coatings were sampled from 30 different locations within the Greater Sudbury area (Fig. 1a), focusing on regions in the vicinity of the smelters and other areas affected by smelting emissions because of their location with respect to the dominant northeasterly wind direction. Coatings were examined using scanning electron microscopy (JEOL 6405), electron microprobe (Cameca SX-100), powder X-ray diffraction (Philips PW 1729), scanning transmission electron microscope (FEI Titan 80-300), micro-X-ray fluorescence (VESPERs beamline, Canadian Lightsource), and laser ablation-inductively coupled plasma mass

spectrometry (LA-ICP-MS; Thermo-Fisher X II Series mass spectrometer; details in Mantha et al. 2012a).

Twenty-six samples were prepared for sulphur-isotope measurements by collecting scrapings of the coatings. The $\delta^{34}\text{S}$ values of the samples were analyzed using a MAT 252 Stable Isotope Ratio Mass Spectrometer coupled with a Carlo Erba NCS 2500 Elemental Analyzer at the Queen's University Facility for Isotope Research. The $\delta^{34}\text{S}$ values of the coating material were calculated by normalizing the $^{34}\text{S}/^{32}\text{S}$ ratios in the sample to the same ratio in the Canyon Diablo Troilite (CDT) standard, and are expressed as δ values in units of permil. The scrapings also contained material from the underlying rock; however, optical microscope and scanning electron microscope (SEM) examination indicated the absence of sulphur-bearing minerals in the underlying rock (Mantha et al. 2012b).

RESULTS AND DISCUSSIONS

Black coatings occur on weathered granites, gneisses, metabasalts and gabbros throughout the Sudbury area (Mantha et al. 2012a; Fig. 1b). Their distribution within the area is consistent with a smelter impacted zone, in terms of SO_2 emissions, soil degradation and vegetation damage. The distribution of metals from a point source and subsequent environmental damage is consistently shown to be controlled, in part, by prevailing winds (Knight and Henderson 2006).

Mineralogical and Chemical Composition

Coating thickness varies from $<5\ \mu\text{m}$ to $>100\ \mu\text{m}$ and was observed to reach up to $200\ \mu\text{m}$ including accumulations of particulate matter and detrital mineral grains. The rock-coating boundary with the underlying rock is typically sharp at the micrometre scale. Feldspars and amphiboles in the underlying rock are weathered and the interstices between these minerals show a loss of mass and occasional infilling of cracks by coating material. Coatings are composed of a porous amorphous silica matrix that contains a wide variety of detrital and smelter-derived particulate matter as well as metal sulphate-rich layers (Fig. 1c).

Detrital particles originate from the underlying rock or from adjacent rocks and soils and are composed of micrometre-size feldspar, amphibole and quartz fragments (Fig. 1c) as well as nanoparticles of hematite, chlorite, and montmorillonite. Various types of smelter-derived nanoparticles occur within the silica matrix. There are spherical, spinel-type particulate matter (mainly magnetite (Fe_3O_4), cupro spinel (CuFe_2O_4), and trevorite (NiFe_2O_4); Fig. 2a), as well as angular CuO_x nanoparticles (Fig. 2b) and amorphous carbon nanoparticles (Fig. 2c). The latter are similar in appearance to coke particulate matter observed in soils surrounding a Zn-smelter in France (Sivry et al. 2010), and are commonly generated during the burning of coal for metallurgical purposes. Smelter-derived particulate matter are mainly composed of magnetite and fayalite (Fe_2SiO_4), and are locally observed with diameters greater than the average thickness of coatings in which they are embedded (Mantha et al. 2012a; Fig. 1c).

Metal-sulphate-rich layers are prominent features within the silica-rich matrix (Figs. 1c and 2d). They contain high concentrations of metal(loid)s such as Pb (max. 10 wt%), Cu (max. 2.2 wt%), As (max. 1.8 wt%) and Se (max. 0.1 wt%) and are composed of nanometre aggregates of numerous sulphates such as (synthetic) $(\text{Fe}^{3+}_4\text{SO}_4(\text{OH})_{10})$, goldichite ($\text{KFe}^{3+}(\text{SO}_4)_2(\text{H}_2\text{O})_4$), mereiterite ($\text{K}_2\text{Fe}^{2+}(\text{SO}_4)_2(\text{H}_2\text{O})_4$), guildite ($\text{CuFe}^{3+}(\text{SO}_4)_2(\text{OH})(\text{H}_2\text{O})_4$), butlerite ($\text{Fe}^{3+}(\text{SO}_4)(\text{OH})(\text{H}_2\text{O})_2$) and antlerite ($\text{Cu}_3(\text{SO}_4)(\text{OH})_4$) (Fig. 2e, f).

Model for Coatings Formation

Acidic emissions and wet precipitation, the result of vast quantities of SO_2 released from smelting, increased the chemical weathering rate of exposed rocks in the Greater Sudbury area. Non-stoichiometric dissolution of silicate minerals at the rock surface resulted in the formation of a hydrous silica gel having a lower viscosity than other solid-state weathering products. This allowed the trapping and accumulation of particulate matter (e.g. Perry et al. 2006), best observed in Figure 1c, in which a thin silica-rich layer holds a spherical particle with a significantly




Figure 2. High-Resolution Transmission Electron Microscope images (HRTEM) and Fast Fourier Transformation (FFT) patterns of former atmospheric nanoparticles within the amorphous silica matrix of the coatings. (a) Fe-rich spherical particle with spinel-type structure; (b) angular CuO_x particle; (c) amorphous carbon-rich spherical particle; (d) back-scattered electron image of metal-sulphate-rich layers (bright layers) along the atmosphere-coating and rock-coating interfaces (labelled 1 and 2, respectively); (e) HRTEM and FFT pattern of mereiterite (A) and (synthetic) $(\text{Fe}^{3+}_4\text{SO}_4(\text{OH})_{10})$ (B); (f) TEM image of antlerite particle (encircled).

larger diameter. The silica gel also contained a connected interstitial pore system that allowed the mixing of pore fluids, the *in situ* growth of minerals, and the entombment of particular matter. In these pore systems, dissolution of trapped nanoparticles by sulphuric acid produced metal(loid)-sulphate-rich pore solutions, which subsequently mixed with solutions containing elements released via the dissolution of the underlying rock (e.g. Na, K, Al, Fe). This mixing process resulted in the precipitation of K-Fe-sulphates on the atmosphere-coating and rock-coating interfaces (Mantha et al. 2012a; Fig. 2d).

Adsorbed silica species have

the ability to modify transformation processes of minerals (Cornell and Schwertmann 1996). For example, adsorbed silica species are known to slow down or even inhibit the transformation of ferrihydrite (Fe-hydroxide) and jarosite (Fe-sulphate) into the thermodynamically more stable phases, goethite and hematite. Hence, dissolved silica-species sorbed to precipitated and encapsulated precipitates of goldichite, mereiterite, guildite, butlerite and antlerite may have inhibited their transformation to thermodynamically more stable minerals. As the gel-like material further condensed and dehydrated, it began to lose its gel properties, resulting in the limited dif-

fusion of dissolved substances (including H^+) and effectively protecting nucleated and incorporated minerals. The apparent disequilibrium between many embedded nanoparticles such as clays and (hydr)oxides (stable at near neutral pH) and the Fe-Cu-sulphates (stable at low pH) indicates that prior to the hardening of the gel, trapped nanoparticles were encapsulated in pores, which prevented their equilibration with acidic metal-sulphate-bearing fluids.

This model differs from previous models for the formation of metal-sulphate- and silica-enriched coatings (Schiffman et al. 2006; Minitti et al. 2007) because it emphasizes the

initial formation of a hydrous silica gel-type layer and proposes that all subsequent processes (trapping of particulate matter, their dissolution and encapsulation) are controlled by the properties of this silica layer.

Change in Chemical Composition on the Regional Scale

The concentrations of smelter-emitted metal(loid)s (Cu, Ni, Pb, As and Se) decrease in the coatings in proportion to the distance from the smelter point-sources (Mantha et al. 2012b), which agrees with results from previous studies on the distribution of metal(loid)s in soils (Freedman and Hutchinson 1980; Hogan and Wotton 1984; Wren 2012), peat (Zoltai 1988), lake sediments (Jackson 1978), snow (Telmer et al. 2004) and humus and till sediments (Henderson et al. 1998). However, correlations between concentrations of elements in coatings from different locations only occur if these elements are also closely associated in the coatings. For example, chemical distribution maps indicate the strong association of Pb–As and Cu–Se in the coatings and in the weathered portion of the underlying rock. Similarly, 152 LA-ICP-MS depth-scan analyses indicate a close association of Pb with As and Cu with Se in all examined coatings, independent of the geographic location or the type of underlying rock (Fig. 3a, b).

Copper is one of the major elements in the metal-sulphate-rich layer (Fig. 2d), whereas Ni is typically found within spherical Fe-oxide smelter particulate matter (Mantha et al. 2012a; Lanteigne et al. 2012; Fig. 2a). Consequently, the concentrations of Ni and Cu do not correlate in all coatings of the Greater Sudbury area (Mantha et al. 2012b), which can be attributed to variations in the proportions of spherical particulate matter and metal-sulphate-rich layers with the type of rock substrate and the geographic location of the coating (Mantha et al. 2012a).

These correlations as well as differences between ratios of elements in the coatings and smelter plume (based on published emission data; Wren 2012) indicate that the elemental ratios in the coatings were mainly controlled by the occurrence of minerals




Figure 3. (a) Plot of Pb versus As (moles per 100 gr of coating) for 30 examined coating samples (a total of 152 LA-ICP-MS depth scan analyses). The uncertainty for the LA-ICP-MS data ($\pm 8\%$) is indicated for one data point; calculated linear regression lines and their equations are indicated; (b) plot of Cu versus Se (weight %) for the 30 examined coating samples; the uncertainty for the LA-ICP-MS data is given for one data point; (c) correlation between the $\delta^{34}\text{S}$ values of the coatings and their distance from the smelters at Copper Cliff (circles), Falconbridge (triangles) and Coniston (squares). The size of the symbol represents approximately the uncertainty of a data point; the average $\delta^{34}\text{S}$ value for the sulphate aerosols is indicated with a star (see text for details); (d) Correlation between distance from the Copper Cliff smelter facility and the Se:Ni ratio in coatings (for more details see Mantha et al. 2012b).

in the sulphate-rich layers and the ability of those minerals to incorporate trace metals based on their atomic size, coordination and charge (Mantha et al. 2012b).

Change in Isotope Composition on the Regional Scale

An important fingerprinting tool for identifying sources of pollutants once present in the atmosphere is the identification of isotopic compositions of anthropogenic SO_x . Several studies have examined preserved traces of atmospheric pollutants in urban areas (e.g. Norman et al. 2006; Sinha et al. 2008), and around coal-fired power plants (e.g. Forrest and Newman 1977) and smelters (Nriagu and Coker 1978), as well as to interpret the related formation of weathering crusts on building stones (Buzek and Šrámek 1985).

The sulphur isotope composi-

tion ($\delta^{34}\text{S}$) of the coatings in the Greater Sudbury area varies between $+2.6\%$ and $+5.7\%$. The higher values are associated with samples collected in closer proximity to the smelting centres, notably Copper Cliff and Coniston. A correlation between the $\delta^{34}\text{S}$ values of the coatings and their distance to the smelters was established through consideration of the prevailing southwest-to-northeast wind direction and emission data from the various smelters. Historical emission data from the smelters at Copper Cliff, Falconbridge and Coniston and the prevailing wind direction suggested that all coatings sampled west of Coniston and Falconbridge were predominantly affected by emissions from the Copper Cliff smelters, and that only coatings sampled around Coniston and Falconbridge were significantly affected by the emissions of the stacks in these

areas. In this way, distances between sample locations and the Copper Cliff smelter center were calculated for three areas west of Coniston and Falconbridge (Sudbury Centre, Sudbury South, and Copper Cliff; Fig. 1a), whereas distances to the Coniston and Falconbridge stacks were only calculated for sampling locations in their vicinity. Figure 3c indicates a good correlation between the $\delta^{34}\text{S}$ values of the coatings and their distances to the various smelters.

Coatings: Geological Records of Past Atmospheric Processes

The regional distribution of metal(loid)s and sulphur isotopes in black rock-coatings are the result of atmospheric processes such as wet and dry deposition, mixing of aerosols and scavenging effects. These processes and their preservation in the coatings reflect the speciation of metal(loid)s in the emission plume and the deposition and mixing of different sulphur-bearing atmospheric species. Information about the nature and fate of atmospheric sulphur species can be gained by comparing the $\delta^{34}\text{S}$ values of the coatings with those of the smelter plume and precipitations, and by evaluating the possibility of sulphur fractionation during oxidation of atmospheric SO_2 .

Sulphur Isotope Composition of SO_2 and Sulphates in the Smelter Plume and Wet Precipitations

Smelters in the Greater Sudbury are known to release both SO_2 gas and sulphate-bearing particulate matter (Forrest and Newman 1977; Wren 2012). Stable isotope studies of the main smelter plume (the Super Stack) documented that SO_2 in the plume displayed low positive $\delta^{34}\text{S}$ values of +0.9‰ to +1.2‰ (Nriagu and Coker 1978), which is consistent with the isotopic composition of most of the smelted Ni–Cu ores (+0.2‰ to +3.5‰) in this region (Thode et al. 1962; Schwarcz 1973). Sulphates in the smelter plume mainly occur in the form of H_2SO_4 , plus minor amounts of metal-sulphate aerosols. The proportion of these primary sulphates in the plume varies between 0.4 to 3% of total SO_2 emissions (Chan et al. 1982a, b; 1983) and their $\delta^{34}\text{S}$ values are in the

range of +4.2 to +7.6‰ (Lusis and Wiebe 1976).

Studies on the conversion of SO_2 to H_2SO_4 indicate that the oxidation rate of SO_2 is on average 1% per hour in the main smelter plume (Lusis and Wiebe 1976; Forrest and Newman 1977). These studies further indicate that the smelter plume travelled, on average, 40 km per hour. Sample locations of the black coatings were all within 40 km of the smelter centres, thus the isotopic signature of sulphates deposited in this area was significantly influenced by primary sulphates (+4.2 to +7.6‰) and their mixing with secondary sulphates (i.e. sulphates formed from the oxidation of atmospheric SO_2).

Chan et al. (1982c, d, e) and Chan and Lusis (1985) further demonstrated that sulphates and trace metals from the plume are efficiently scavenged during wet precipitation events. For example, typical scavenging coefficients for sulphates were found to be about 30 to 40% per hour, and for trace metals such as Pb, Cu, Se and As, 60 to 80% per hour (see below). The same studies also suggest that the scavenging coefficient for SO_2 was insignificantly small relative to that of sulphates, indicating that exposed rocks in the Sudbury area mainly interacted with primary and secondary sulphates scavenged via wet precipitation.

Did Fractionation of Sulphur Occur?

The fractionation of S isotopes in the atmosphere can be mediated by a gas-phase or an aqueous-phase oxidation process. The gas-phase oxidation (also known as homogeneous oxidation) occurs via OH radicals in the troposphere and stratosphere and produces sulphuric acid (H_2SO_4), which condenses to form aqueous $(\text{SO}_4)^{2-}$ and $(\text{SO}_4)^{2-}$ -bearing aerosols (Saltzman et al. 1983; Tanaka et al. 1994). The aqueous-phase oxidation (also known as heterogeneous oxidation) occurs in the aqueous component or on particle surfaces in the atmosphere. The major oxidants are H_2O_2 , O_3 and O_2 , the latter being catalyzed by Fe^{3+} and other transition metal ions (Herrmann et al. 2000).

The isotopic composition of secondary sulphates ($\delta^{34}\text{S}_{\text{H}_2\text{SO}_4}$) formed during Rayleigh distillation can be cal-

culated using the following equation:

$$\delta^{34}\text{S}_{\text{H}_2\text{SO}_4} = \alpha (\delta^{34}\text{S}_{\text{SO}_2}^0 + 10^3)^{f-1} - 10^3$$

where $\delta^{34}\text{S}_{\text{SO}_2}^0$ is the initial $\delta^{34}\text{S}$ value for SO_2 (average value of SO_2 in the plume = 1.11‰ for $f = 1$), f the fraction of SO_2 remaining (Faure 1998), and α is the fractionation factor for a gas-phase or aqueous-phase oxidation process (for details see Mantha et al. 2012b).

Using the above equation and reported fractionation factors for gas-phase or aqueous-phase oxidation processes, Mantha et al. (2012b) showed that secondary sulphates would have either large positive (>11‰) or negative (<-9‰) $\delta^{34}\text{S}$ values if the remaining fraction of SO_2 decreases in steps of 0.01 (corresponding to an oxidation rate of 1% per hour). Hence, mixing of similar proportions of primary sulphates (+4.2 to +7.6‰) with secondary sulphates containing fractionated sulphur (values of either >11‰ or <-9‰) would result in $\delta^{34}\text{S}_{\text{H}_2\text{SO}_4}$ values that are not consistent with those measured in the coatings (+2.6 to +5.7‰). As a result, sulphur fractionation during gas-phase or aqueous-phase oxidation processes was disregarded as the cause of the decrease of $\delta^{34}\text{S}$ with distance from the smelters.

Mixing of Primary and Secondary Sulphates

The content of primary sulphates in the emission plume(s) ranges between 0.4 to 3.0% (Chan et al. 1982c, d, e). Hence, an SO_2 oxidation rate of 1% per hour would result in approximately equal amounts of primary and secondary sulphates within the first two hours after emission of the plume. The average $\delta^{34}\text{S}$ for primary sulphates (5.4‰) is similar to the $\delta^{34}\text{S}$ values observed in the coatings proximal to the smelters (Fig. 3c). If gas-phase or aqueous-phase oxidation processes did not result in fractionation of sulphur isotopes, as observed in the urban areas of Calgary and Vancouver (Norman et al. 2004a, b), the $\delta^{34}\text{S}$ for the secondary sulphates would be similar to those for the SO_2 (average value = 1.11‰) from which they formed. Two processes occurred now simultaneously in the smelter plume with increasing distance to the smelter: a) Primary sulphates were scavengers by wet precipitation resulting in a decrease of sulphates

with $\delta^{34}\text{S} = 5.4\%$; and b) Oxidation of SO_2 resulted in an increase of secondary sulphates with $\delta^{34}\text{S} = 1.11\%$.

These two processes yielded to an increasing ratio of secondary ($\delta^{34}\text{S} = 1.11\%$) to primary sulphates ($\delta^{34}\text{S} = 5.4\%$) in the smelter plume and thus in a decrease in the $\delta^{34}\text{S}$ of atmospheric sulphates (and consequently of sulphur species in the coatings) with increasing distance from the smelters. This mixing model was verified through inspection of the regional distribution of trace metals in the coatings, which were found to be closely associated with primary sulphate aerosols in the smelter plume.

Deposition Rates of Sulphate Aerosols and Larger Particulate Matter

Studies of particulate matter in the Copper Cliff smelter plume found that Pb, As and Se occurred predominantly in small sulphate aerosols having average diameters $<2.5\ \mu\text{m}$, compared to Cu and Ni, which were found mainly in larger particulate matter ($>2.5\ \mu\text{m}$; Chan et al. 1982a, b). Studies on the wet and dry deposition rates of metals from a smelter plume indicate that deposition rates for the larger Fe–Ni particulate matter are similar during wet and dry conditions, whereas the rates for the smaller metal-sulphate aerosols dramatically increase during wet deposition (Chan et al. 1982c,d,e). These results indicate that during precipitation events, the travelling smelter plumes became depleted in Pb-, As- and Se-aerosols with increasing distance to the stacks. Hence, precipitation and consequently coatings closer to the smelters became more enriched in Pb-, As- and Se-bearing metal-sulphates relative to Fe–Ni-bearing particulate matter, than precipitation and coatings farther away from the smelters. Plots of Pb/Ni, As/Ni, and Se/Ni ratios in coatings versus their distance to the Copper Cliff smelter indicate that coatings closer to the smelter are indeed enriched in Pb, As and Se relative to Ni (Mantha et al. 2012b; Fig. 3d). Furthermore, the ratio of primary Pb–As–Se-bearing sulphates to secondary sulphates decreases with distance from the smelter, which is in accordance with the interpretation of the decrease in $\delta^{34}\text{S}$ with

distance from the point source.

CONCLUSIONS AND OUTLOOK

Our studies on the black coatings of the Greater Sudbury area indicate that silica coatings are remarkable geological records of nanoparticles emitted to the atmosphere as well as atmospheric processes such as mixing, deposition and scavenging. Detailed mineralogical and chemical characterization of the coatings indicated the following characteristic features of silica coatings in their gelatinous state:

- i) they contain a connected interstitial pore system that allowed the *in situ* formation of secondary minerals and the mixing of pore fluids;
- ii) they have a low viscosity that allowed the trapping and entombment of nano- to micro-size particulate matter; and
- iii) their nanometre-size pores and adsorbed silica species allowed the encapsulation and preservation of nanoparticles and soluble acidic and basic Fe–Cu-sulphates.

On a regional scale, the coatings display a decrease in metal concentrations, an increase in the ratio of secondary to primary sulphates, and a decrease in the $\delta^{34}\text{S}$ of primary and secondary atmospheric sulphates with increasing distance from the smelter point-sources. The coatings also record:

- i) higher scavenging and thus higher deposition rates of primary sulphates containing Pb, As and Se by wet precipitations; and
- ii) similar deposition rates of larger Ni-bearing particulate matter under both wet and dry conditions.

A comparison of emission data and elemental ratios between Pb, As, Cu and Ni ratios in the coatings showed that the ratios are mainly controlled by the occurrence of minerals in the sulphate-rich layers and their ability to incorporate trace metals according to their atomic size, coordination and charge (Mantha et al. 2012b).

Another interesting aspect is that silica-rich coatings on Mars, which are believed to be products of the weathering of the underlying rock, contain sulphates and amorphous silica as their main constituents (Tosca et al.

2005; Tosca and McLennan 2006; Hurowitz et al. 2006). The present study has shown that silica and Fe-sulphate coatings preserve the isotopic signature of sulphur involved in their formation as well as differences between wet deposition rates for atmospheric sulphate species and particular matter. Similarly, silica coatings on Mars may contain information with respect to atmospheric sources and processes.

ACKNOWLEDGEMENTS

This work was supported by a NSERC Discovery grant to MS, and an Ontario Graduate Scholarship to NM. KK acknowledges financial support from NSERC, CFI and OCE. We would like to thank K. Klassen for her help at the Queen's University Stable Isotope Lab. MH acknowledges NSF and EPA under NSF Cooperative Agreement EF-0830093. Any opinions, findings, conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the NSF or the EPA. This work has not been subjected to EPA review and no official endorsement should be inferred. We would like to thank technical staff members at Laurentian University, Ontario GEOLABS, Queen's University Stable Isotope Lab and at the Nanoscale Characterization and Fabrication Laboratory, ICTAS, Virginia Tech for their assistance. We also want to thank Editor Brendan Murphy and an anonymous reviewer for their comments.

REFERENCES

- Buzek, F. and Šrámek, J., 1985, Sulphur isotopes in the study of stone monument conservation: Studies in Conservation, v. 30, p. 171-176, <http://dx.doi.org/10.2307/1506039>.
- Chan, W.H and Lusic, M.A., 1985, Post-superstack smelter emissions and their fate in the atmosphere: An overview of the Sudbury environment study results: Water, Air and Soil Pollution, v. 26, p. 43-58, <http://dx.doi.org/10.1007/BF00299487>.
- Chan, W.H., Vet, R.J., Skelton, G.B., and Lusic, M.A., 1982a, Size distribution and emission rate measurements of particulates in the 93m Falconbridge smelter stack plume, 1979: Report

- ARB-TDA-57-80, Air Resources Branch, Ontario Ministry of the Environment.
- Chan, W.H., Vet, R.J., Lusi, M.A., and Skelton, B.G., 1982b, Size distribution and emission rate measurements of particulates in the Inco 381 m chimney and iron ore recovery plant stack fumes, 1979-80: Report ARB-TDA-62-80, Air Resources Branch, Ontario Ministry of the Environment.
- Chan, W.H., Vet, R.J., Ro, C.U., Tang, A.J.S., and Lusi, M.A., 1982c, Precipitation quality and wet deposition in the Sudbury basin: Sudbury environmental cumulative precipitation network results: Report ARB-04-82-ARSP, Air Resources Branch, Ontario Ministry of the Environment.
- Chan, W.H., Vet, R.J., Ro, C.U., Tang, A.J.S., and Lusi, M.A., 1982d, An analysis of the impact of smelter emissions on precipitation quality and wet deposition in the Sudbury area: Sudbury environmental study event precipitation network results: Report ARB-05-82-ARSP, Air Resources Branch, Ontario Ministry of the Environment.
- Chan, W.H., Vet, R.J., Ro, C.U., Tang, A.J.S., and Lusi, M.A., 1982e, An analysis of the impact of smelter emissions on atmospheric dry deposition in the Sudbury area: Sudbury environmental study event precipitation network results: Report ARB-12-81-ARSP, Air Resources Branch, Ontario Ministry of the Environment.
- Chan, W.H., Vet, R.J., Lusi, M.A., and Skelton, G.B., 1983, Airborne particulate size distribution measurements in nickel smelter plumes: *Atmospheric Environment*, v. 17, p. 1173-1181, [http://dx.doi.org/10.1016/0004-6981\(83\)90340-2](http://dx.doi.org/10.1016/0004-6981(83)90340-2).
- Cornell, R.M., and Schwertmann, U., 1996, *The Iron Oxides: Structure, Properties, Reactions, Occurrences and Uses*: VCH Verlagsgesellschaft, New York, USA, 703 p.
- Curtiss, B., Adams, J.B., and Ghorso, M.S., 1985, Origin, development and chemistry of silica-alumina rock coatings from the semiarid regions of the island of Hawaii: *Geochimica et Cosmochimica Acta*, v. 49, p. 49-56, [http://dx.doi.org/10.1016/0016-7037\(85\)90190-5](http://dx.doi.org/10.1016/0016-7037(85)90190-5).
- Dorn, R., 1998, *Rock coatings*: Amsterdam, Elsevier, 429 p.
- Durocher, J.L., and Schindler, M., 2011, Iron-hydroxide, iron-sulphate and hydrous silica coatings in acid-mine tailings facilities: A comparative study of their trace-element composition: *Applied Geochemistry*, v. 26, p. 1337-1352, <http://dx.doi.org/10.1016/j.apgeochem.2011.05.007>.
- Faure, G., 1998, *Principles and Applications of Geochemistry*, 2nd edition: Prentice-Hall Inc., New Jersey, 600 p.
- Fisk, E.P., 1971, Desert Glaze: *Journal of Sedimentary Petrology*, v. 41, p. 1136-1137.
- Forrest, J., and Newman, L., 1977, Oxidation of sulfur dioxide in the Sudbury smelter plume: *Atmospheric Environment*, v. 11, p. 517-520, [http://dx.doi.org/10.1016/0004-6981\(77\)90068-3](http://dx.doi.org/10.1016/0004-6981(77)90068-3).
- Freedman, B., and Hutchinson, T.C., 1980, Pollutant inputs from the atmosphere and accumulations in soils and vegetation near a nickel-copper smelter at Sudbury, Ontario, Canada: *Canadian Journal of Botany*, v. 58, p. 108-132.
- Friedmann, E.I., and Weed, R., 1987, Microbial trace-fossil formation, biogenous, and abiotic weathering in the Antarctic cold desert: *Science*, v. 236, p. 703-705, <http://dx.doi.org/10.1126/science.11536571>.
- Giorgetti, G., and Baroni, C., 2007, High-resolution analysis of silica and sulphate-rich rock varnishes from Victoria Land (Antarctica): *European Journal of Mineralogy*, v. 19, p. 381-389, <http://dx.doi.org/10.1127/0935-1221/2007/0019-1725>.
- Gundermann, D.G., and Hutchinson, T.C., 1995, Changes in soil chemistry 20 years after the closure of a nickel-copper smelter near Sudbury, Ontario, Canada: *Journal of Geochemical Exploration*, v. 52, p. 231-236, [http://dx.doi.org/10.1016/0375-6742\(94\)00026-8](http://dx.doi.org/10.1016/0375-6742(94)00026-8).
- Henderson, P.J., McMartin, I., Hall, G.E., Percival, J.B., and Walker, D.A., 1998, The chemical and physical characteristics of heavy metals in humus and till in the vicinity of the base metal smelter at Flin Flon, Manitoba, Canada: *Environmental Geology*, v. 34, p. 39-58, <http://dx.doi.org/10.1007/s002540050255>.
- Herrmann, H., Ervens, B., Jacobi, H.-W., Wolke, R., Nowacki, P., and Zellner, R., 2000, CAPRAM2.3: A chemical aqueous phase radical mechanism for tropospheric chemistry: *Journal of Atmospheric Chemistry*, v. 36, p. 231-284, <http://dx.doi.org/10.1023/A:1006318622743>.
- Hoar, K., Nowinski, P., Hodge, V., and Cizdziel, J., 2011, Rock varnish: A passive forensic tool for monitoring recent air pollution and source identification: *Nuclear Technology*, v. 175, p. 351-359.
- Hodge, V.F., Farmer, D.E., Diaz, T., and Orndorff, R.L., 2005, Prompt detection of alpha particles from ²¹⁰Po: Another clue to the origin of rock varnish?: *Journal of Environmental Radioactivity*, v. 78, p. 331-342, <http://dx.doi.org/10.1016/j.jenvrad.2004.05.009>.
- Hogan, G.D., and Wotton, D.L., 1984, Pollutant distribution and effects in forests adjacent to smelters: *Journal of Environmental Quality*, v. 13, p. 377-382, <http://dx.doi.org/10.2134/jecq1984.00472425001300030011x>.
- Hurowitz, J.A., McLennan, S.M., Tosca, N.J., Arvidson, R.E., Michalski, J.R., Ming, D.W., Schröder, C., and Squyres, S.W., 2006, In situ and experimental evidence for acidic weathering of rocks and soils on Mars: *Journal of Geophysical Research*, v. 111, E02S19, <http://dx.doi.org/10.1029/2005JE002515>.
- Jackson, T.A., 1978, The biogeochemistry of heavy metals in polluted lakes and streams at Flin Flon, Canada, and a proposed method for limiting heavy-metal pollution of natural waters: *Environmental Geology*, v. 2, p. 173-189, <http://dx.doi.org/10.1007/BF02430671>.
- Knight, R.D., and Henderson, P.J., 2006, Smelter dust in humus around Rouyn-Noranda, Quebec: *Geochemistry: Exploitation, Environment, Analysis*, v. 6, p. 203-214.
- Langworthy, K.A., Krinsley, D.H., and Dorn, R.I., 2010, High resolution transmission electron microscopy evaluation of silica glaze reveals new textures: *Earth Surface Processes and Landforms*, v. 35, p. 1615-1620, <http://dx.doi.org/10.1002/esp.2061>.
- Lanteigne, S., Schindler, M., McDonald, A.M., Skeries, K., Abdu, Y., Mantha, N.M., Murayama, M., Hawthorne, F.C., and Hochella, M.F., Jr., 2012, Mineralogy and weathering of smelter-derived spherical particles in soils: Implications for the mobility of Ni and Cu in the surficial environment: *Water Air Soil Pollution*, <http://dx.doi.org/10.1007/s11270-012-1135-3>.
- Lusi, M.A., and Wiebe, H.A., 1976, The rate of oxidation of sulfur dioxide in the plume of a nickel smelter stack: *Atmospheric Environment*, v. 10, p. 793-798,

- 6981(76)90133-5.
- Mantha, N.M., Schindler, M., Murayama, M., and Hochella, M.F., 2012a, Silica and sulfate-bearing rock coatings in smelter areas: Products of chemical weathering and atmospheric pollution I. Formation and mineralogical composition: *Geochimica et Cosmochimica Acta*, v. 85, p. 254-274, <http://dx.doi.org/10.1016/j.gca.2012.01.033>.
- Mantha, N.M., Schindler, M., and Kyser, T.K., 2012b, Silica- and sulfate-bearing rock coatings in smelter areas: Part II. Forensic tools for atmospheric metal(loid)- and sulfur-isotope compositions: *Geochimica et Cosmochimica Acta*, v. 90, p. 221-241, <http://dx.doi.org/10.1016/j.gca.2012.05.013>.
- McLennan, S.M., 2003, Sedimentary silica on Mars: *Geology*, v. 31, p. 315-318, [http://dx.doi.org/10.1130/0091-7613\(2003\)031<0315:SSOM>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2003)031<0315:SSOM>2.0.CO;2).
- Milliken, R.E., Swayze, G.A., Arvidson, R.E., Bishop, J.L., Clark, R.N., Ehlmann, B.L., Green, R.O., Grotzinger, J.P., Morris, R.V., Murchie, S.L., Mustard, J.F., and Weitz, J., 2008, Opaline silica in young deposits on Mars: *Geology*, v. 36, p. 847-850, <http://dx.doi.org/10.1130/G24967A.1>.
- Miniti, M.E., Weitz, C.M., Lane, M.D., and Bishop, J.L., 2007, Morphology, chemistry, and spectral properties of Hawaiian rock coatings and implications for Mars: *Journal of Geophysical Research: Planets*, v. 112, E05015, <http://dx.doi.org/10.1029/2006JE002839>.
- Nord, A.G., Svärdh, A., and Tronner, K., 1994, Air pollution levels reflected in deposits on building stone: *Atmospheric Environment*, v. 28, p. 2615-2622, [http://dx.doi.org/10.1016/1352-2310\(94\)90435-9](http://dx.doi.org/10.1016/1352-2310(94)90435-9).
- Norman, A.-L., Belzer, W., and Barrie, L., 2004a, Insights into the biogenic contribution to total sulphate in aerosol and precipitation in the Fraser Valley afforded by isotopes of sulphur and oxygen: *Journal of Geophysical Research: Atmospheres*, v. 109, D05311, <http://dx.doi.org/10.1029/2002JD003072>.
- Norman, A.L., Krouse, H.R., and MacLeod, J., 2004b, Apportionment of pollutant S in urban airshed: Calgary, Canada; a case study: *In* Borrego, C. and Incecik, S., eds., *Air Pollution Modeling and its Application*: Kluwer, New York, p. 107-125.
- Norman, A.-L., Anlauf, K., Hayden, K., Thompson, B., Brook, J.R., Li, S.-M., and Bottenheim, J., 2006, Aerosol sulphate and its oxidation on the Pacific NW coast: S and O isotopes in PM_{2.5}: *Atmospheric Environment*, v. 40, p. 2676-2689, <http://dx.doi.org/10.1016/j.atmosenv.2005.09.085>.
- Nriagu, J.O., and Coker, R.D., 1978, Isotopic composition of sulphur in atmospheric precipitation around Sudbury, Ontario: *Nature*, v. 274, p. 883-885, <http://dx.doi.org/10.1038/274883a0>.
- Perry, R.S., Lynne, B., Sephton, M., Kolb, V., Perry, C., and Staley, J., 2006, Baking black opal in the desert sun: The importance of silica in desert varnish: *Geology*, v. 34, p. 537-540, <http://dx.doi.org/10.1130/G22352.1>.
- Saltzman, E.S., Brass, G.W., and Price, D.A., 1983, The mechanism of sulfate aerosol formation: Chemical and sulfur isotopic evidence: *Geophysical Research Letters*, v. 10, p. 513-516, <http://dx.doi.org/10.1029/GL010i007p00513>.
- Schiffman, P., Zierenberg, R., Marks, N., Bishop, J., and Dyar, M., 2006, Acid-fog deposition at Kilauea volcano: A possible mechanism for the formation of siliceous-sulfate rock coatings on Mars: *Geology*, v. 34, p. 921-924, <http://dx.doi.org/10.1130/G22620A.1>.
- Schindler, M., Durocher, J.L., Abdu, Y., and Hawthorne, F.C., 2009, Hydrous silica coatings: Occurrence, speciation of metals, and environmental significance: *Environmental Science and Technology*, v. 43, p. 8775-8780, <http://dx.doi.org/10.1021/es9018817>.
- Schindler, M., Fayek, M., and Hawthorne, F.C., 2010, Uranium-rich opal from the Nopal I uranium deposit, Peña Blanca, Mexico: Evidence for the uptake and retardation of radionuclides: *Geochimica et Cosmochimica Acta*, v. 74, p. 187-202, <http://dx.doi.org/10.1016/j.gca.2009.09.017>.
- Schwarcz, H.P., 1973, Sulfur isotope analyses of some Sudbury, Ontario, ores: *Canadian Journal of Earth Sciences*, v. 10, p. 1444-1459, <http://dx.doi.org/10.1139/e73-135>.
- Sinha, B.W., Hoppe, P., Huth, J., Foley, S., and Andreae, M.O., 2008, Sulfur isotope analyses of individual aerosol particles in the urban aerosol at a central European site (Mainz, Germany): *Atmospheric Chemistry and Physics*, v. 8, p. 7217-7238, <http://dx.doi.org/10.5194/acp-8-7217-2008>.
- Sivry, Y., Munoz, M., Sappin-Didier, V., Riotte, J., Denaix, L., de Parseval, P., Destrigneville, C., and Dupre, B., 2010, Multimetallic contamination from Zn-ore smelter: solid speciation and potential mobility in riverine floodbank soils of the upper Lot River (SW France): *European Journal of Mineralogy*, v. 22, p.679-691, <http://dx.doi.org/10.1127/0935-1221/2010/0022-2060>.
- Smith, B.J., and Prikryl, R., 2007, Diagnosis decay: the value of medical analogy in understanding the weathering of building stones. *In* Prikryl, R. and Smith, B.J., eds., *Building Stone Decay: From Diagnosis to Conservation*: Journal of the Geological Society of London, v. 271, p. 1-8.
- Smith, B.J., and Whalley, W.B., 1988, A note on the characteristics and possible origins of desert varnishes from southeast Morocco: *Earth Surfaces Processes and Landforms*, v. 13, p. 251-258, <http://dx.doi.org/10.1002/esp.3290130306>.
- Squyres, S.W., and 17 others, 2008, Detection of silica-rich deposits on Mars: *Science*, v. 320, p. 1063-1067, <http://dx.doi.org/10.1126/science.1155429>.
- Tanaka, N., Rye, D.M., Xiao, Y., and Lasaga, A.C., 1994, Use of stable sulfur isotope systematics for evaluating oxidation reaction pathways and in-cloud scavenging of sulfur dioxide in the atmosphere: *Geophysical Research Letters*, v. 21, p. 1519-1522, <http://dx.doi.org/10.1029/94GL00893>.
- Telmer, K., Bonham-Carter, G.F., Kliza, D.A., and Hall, G.E., 2004, The atmospheric transport and deposition of smelter emissions: Evidence from the multi-element geochemistry of snow, Quebec, Canada: *Geochimica et Cosmochimica Acta*, v. 68, p. 2961-2980, <http://dx.doi.org/10.1016/j.gca.2003.12.022>.
- Thode, H.G., Dunford, H.B., and Shima, M., 1962, Sulfur isotope abundances in rocks of the Sudbury district and their geological significance: *Economic Geology*, v. 57, p. 565-578, <http://dx.doi.org/10.2113/gsecongeo.57.4.565>.
- Tosca, N.J., and McLennan, S.M., 2006, Chemical divides and evaporite assemblages on Mars: *Earth and Planetary Science Letters*, v. 241, p. 21-31,

- <http://dx.doi.org/10.1016/j.epsl.2005.10.021>.
- Tosca, N.J., McLennan, S.M., Clark, B.C., Grotzinger, J.P., Hurowitz, J.A., Knoll, A.H., Schröder, C., and Squyres, S.W., 2005, Geochemical modeling of evaporation processes on Mars: Insight from the sedimentary record at Meridiani Planum: *Earth and Planetary Science Letters*, v. 240, p. 122-148, <http://dx.doi.org/10.1016/j.epsl.2005.09.042>.
- Wayne, D.M., Diaz, T.A., Fairhurst, R.J., Orndorff, R.L., and Pete, D.V., 2006, Direct major- and trace-element analyses of rock varnish by high resolution laser ablation inductively-coupled plasma mass spectrometry (LA-ICPMS): *Applied Geochemistry*, v. 21, p. 1410-1431, <http://dx.doi.org/10.1016/j.apgeochem.2006.04.005>.
- Weed, R., and Norton, S.A., 1991, Siliceous crusts, quartz rinds and biotic weathering of sandstones in the cold desert of Antarctica, *In* Berthelin, J., *ed.*, *Diversity of Environmental Biogeochemistry*, *Developments in Geochemistry*, v. 6, p. 327-339.
- Whalley, W.B., Gellatly, A.F., Gordon, J.E., and Hansom, J.D., 1990, Ferromanganese rock varnish in north Norway: A subglacial origin: *Earth Surface Processes and Landforms*, v. 15, p. 265-275, <http://dx.doi.org/10.1002/esp.3290150308>.
- Wren, C., 2012, Risk assessment and environmental management: A case study in Sudbury, Ontario, Canada: *Progress in Environmental Sciences*, v. 1, p. 1-450.
- Zoltai, S.C., 1988, Distribution of base metals in peat near a smelter at Flin Flon, Manitoba: *Water, Air, and Soil Pollution*, v. 37, p. 217-228, <http://dx.doi.org/10.1007/BF00226493>.

Received June 2012

Accepted as revised October 2012