

Gravel Cones and Depressions: Sedimentary Indicators of Intense Winds

Nicholas Eyles

Volume 12, numéro 2, august 1976

URI : https://id.erudit.org/iderudit/ageo12_2rep03

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer cet article

Eyles, N. (1976). Gravel Cones and Depressions: Sedimentary Indicators of Intense Winds. *Atlantic Geology*, 12(2), 75–76.

GRAVEL CONES AND DEPRESSIONS: SEDIMENTARY INDICATORS OF INTENSE WINDS

NICHOLAS EYLES¹

Department of Geology, Memorial University of Newfoundland,
St. John's, Newfoundland, Canada

This note describes groups of unusual gravel cones and depressions forming on an extensive beach which is part of a breached, bay-mouth bar across Holyrood Pond, Avalon Peninsula, southeastern Newfoundland (Fig. 1).

The cones are flat topped with approximately 30° side slopes, up to 0.3 m high, and triangular in profile (Fig. 2). They are graded from pebbles at the base to pea gravel at the apex. The gravel is set in a matrix of sand (Fig. 3). Inverted gravel cones, pebble-lined depressions, also occur but less frequently. The cones and depressions are thought to be generated by intense, short-lived cyclic winds popularly called "wind devils".

The beach upon which the cones occur is subject to steady summer onshore winds from the south-southwest characterized by strong gusts. The nearest station for which wind speed data are available is Argentia airport, 65 kms to the northwest. Average wind speeds during the summer range from 19.3 to 27.2 kms/hr., with gusts reaching 137.6 kms/hr. (Canada Dept. of Transport 1968, for the period 1955-1964). The beach sediments consist of sand and rounded to discoidal-shaped pebbles which are derived from nearby coastal cliffs of till (Eyles and Slatt, In press). Larger pebbles tend to be discoidal and occur as well imbricated pebble pavements behind prominent storm beaches. Smaller pebbles are rounded and lack imbrication. Steady




Fig. 1. Location of Holyrood Pond.

¹Present address:
School of Environmental Sciences,
University of East Anglia, University Village
Norwich, Great Britain



Fig. 2. Newly formed gravel cone. Note key case for scale. In the foreground a gravel cone is being buried by sand.




Fig. 3. A mature gravel cone exhibiting flat top and graded sides.



Fig. 4. The beach strand. Groups of orientated gravel cones and sand ridges form in response to strong, gusty, unidirectional winds.

winds and an abundant sand supply have resulted in a number of small-scale aeolian forms on the beach surface, as well as various stages of burial of the pebble pavement (Fig. 4).

The cones probably are initially established from random groups of pebbles; local concentrations of pebbles in the beach are uncovered by the removal of the sand matrix. Once established, they will act as loci for the formation of wind devils, and the pebbles become transported in a circular fashion. The gradation in particle size of the cones reflects the wind-velocity gradient within the wind devil. Cone height and width are also controlled by wind velocity since the intensity of flow and the ability to transport pebbles will decrease as the area of the cone is increased. For this reason cones wider than 0.5 m at the base were not found.

Initial depressions on the beach also become the foci of development of wind devils. As sand is removed by wind scour, small-scale pebble lags develop. These depressions appear to be self-maintaining for the same reasons as described above for the cones. The maximum depth and width noted are 15 cm, and the pebbles exhibit imbrication toward the centre. Cones and depressions are seen to be destroyed during periods of heavy rain when pebbles and sand may flow as a slurry.

Importantly, these constructional and destructional features are observed to be buried by sand moving onshore during more normal periods of wind velocity (Fig. 3), and thus may be preserved in the geologic record. Their presence can be used as diagnostic criteria of sedimentary environments over which strong winds and gusts blow. For example, aside from their occurrence on a beach, similar forms have been seen to occur as strong winds blow off modern valley glaciers onto outwash plains (R.M. Slatt pers. comm. 1976). A literature search indicates that such forms have not been previously described.

ACKNOWLEDGEMENTS

The writer wishes to thank Dr. R.M. Slatt of the Department of Geology, Arizona State University for critical discussion, and financial support through his National Research Council of Canada operating grant.

REFERENCES

- Canada Department of Transport, 1968, Wind, in Climatic Normals, Vol. 5, pp. 551-552, Meteorology Branch for the period 1955-1964.
- Eyles, N. and R.M. Slatt, Ice-marginal sedimentary, glaciectonic and morphological features of Pleistocene drift; an example from Newfoundland. Quaternary Research, In Press.