

# The Rockslide -Debris Avalanche of the May 18, 1980, Eruption of Mount St. Helens — 10th Anniversary Perspectives

Harry Glicken

Volume 17, numéro 3, septembre 1990

URI : [https://id.erudit.org/iderudit/geocan17\\_3art06](https://id.erudit.org/iderudit/geocan17_3art06)

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)

1911-4850 (numérique)

[Découvrir la revue](#)

Citer cet article

Glicken, H. (1990). The Rockslide -Debris Avalanche of the May 18, 1980, Eruption of Mount St. Helens — 10th Anniversary Perspectives. *Geoscience Canada*, 17(3), 150–153.

Résumé de l'article

The massive rockslide-debris avalanche of the May 18, 1980, eruption of Mount St. Helens began with a retrogressive failure triggered by the 08:32 PDT earthquake. It depressurized the volcano's magmatic and hydrothermal system and produced a hummocky deposit with a volume of 2.5 km<sup>3</sup>. Detailed work provides a comprehensive understanding of a previously poorly understood type of event.

The deposit consists of relatively intact pieces (block facies) of the pre-1980 mountain and mixed material (mixed facies) that is primarily rocks from the pre-1980 Mount St. Helens and the 1980 cryptodome. Travel paths of rockslide blocks are interpreted from a geologic map of the deposit. The material was fractured and dilated during the rockslide, after which grain-to-grain dispersive stress facilitated flow. During transport, the dilated material mixed but significant fine material was not produced.

- Endo, E.T., Malone, S.D., Noson, L.L. and Weaver, C.S., 1981, Locations, magnitudes, and statistics of the March 20-May 18 earthquake sequence, in Lipman, P.W. and Mullineaux, D.R., eds., *The 1980 eruptions of Mount St. Helens*, Washington: United States Geological Survey, Professional Paper 1250, p. 93-107.
- Jonientz-Trisler, C. and Zollweg, J.E., 1987, Sub-epidicial seismicity at Mt. St. Helens prior to the 1980 eruption: Hawaii Symposium on How Volcanoes Work, Hilo, Hawaii, January 19-25, Abstract Volume, p. 126.
- Lipman, P.W., Moore, J.G. and Swanson, D.A., 1981, Bulging of the north flank before the May 18 eruption — geodetic data, in Lipman, P.W. and Mullineaux, D.R., eds., *The 1980 eruptions of Mount St. Helens*, Washington: United States Geological Survey, Professional Paper 1250, p. 143-155.
- Lipman, P.W. and Mullineaux, D.R., 1981, eds., *The 1980 eruptions of Mount St. Helens*, Washington: United States Geological Survey, Professional Paper 1250, 844 p.
- Malone, S.D., Endo, E.T., Weaver, C.S. and Ramey, J.W., 1981, Seismic monitoring for eruption prediction, 1981, in Lipman, P.W. and Mullineaux, D.R., eds., *The 1980 eruptions of Mount St. Helens*, Washington: United States Geological Survey, Professional Paper 1250, p. 803-813.
- Malone, S.D. and Frank, D., 1975, Increased heat emission from Mount Baker, Washington, 1975: EOS, v. 56, p. 679-685.
- Malone, S.D. and Pavlis G.L., 1983, Velocity structure and relocation of earthquakes at Mount St. Helens (abstract): EOS, v. 64, p. 895.
- Moran, S.C. and Malone, S.D., 1989, Pre-1980 Seismicity at Mt. St. Helens: Is the past the key to the present? (abstract): American Geophysical Union, Pacific Northwest Region, 36th annual meeting, Portland State University, Proceedings.
- Qamar, A., St. Lawrence, W., Moore, J.N. and Kendrick, G., 1983, Seismic signals preceding the Explosive eruption of Mount St. Helens, Washington, on 18 May 1980: Bulletin of the Seismological Society of America, v. 16, p. 1797-1813.
- Scandone, R. and Malone, S.D., 1985, Magma supply, magma discharge and the readjustment of the feeding system of Mount St. Helens during 1980: Journal of Volcanology and Geothermal Research, v. 23, p. 239-262.
- Shemeta, J.E. and Weaver, C.S., 1986, Seismicity accompanying the May 18, 1980 eruption of Mount St. Helens, Washington, in Keller, S.A.C., ed., *Mount St. Helens: Five years later*: Eastern Washington University Press, Cheney, WA, p. 44-58.
- Swanson, D.A., Casadevall, T.J., Dzurisin, D., Malone, S.D., Newhall, C.G. and Weaver, C.S., 1983, Predicting eruptions at Mount St. Helens, June 1980 through December, 1982: Science, v. 221, p. 1369-1376.
- Weaver, C.S., Grant, W.C. and Shemeta, J., 1987, Local crustal extension at Mount St. Helens Washington: Journal of Geophysical Research, v. 92, p. 10,170-10,178.
- Weaver, C.S. and Malone, S.D., 1976, Mount St. Helens seismic events: volcanic earthquakes or glacial noises?: Geophysical Research Letters, v. 3, p. 197-200.
- Weaver, C.S., Norris, R.D. and Jonientz-Trisler, C., 1990, Results of seismological monitoring in the Cascade Range, 1960-1989: earthquakes, eruptions, avalanches and other curiosities: Geoscience Canada, v. 17, p. 158-162.



## The Rockslide – Debris Avalanche of the May 18, 1980, Eruption of Mount St. Helens — 10th Anniversary Perspectives

Harry Glicken  
*Department of Geological Sciences  
 University of California  
 Santa Barbara, California 93106*

### Summary

The massive rockslide-debris avalanche of the May 18, 1980, eruption of Mount St. Helens began with a retrogressive failure triggered by the 08:32 PDT earthquake. It depressurized the volcano's magmatic and hydrothermal system and produced a hummocky deposit with a volume of 2.5 km<sup>3</sup>. Detailed work provides a comprehensive understanding of a previously poorly understood type of event.

The deposit consists of relatively intact pieces (block facies) of the pre-1980 mountain and mixed material (mixed facies) that is primarily rocks from the pre-1980 Mount St. Helens and the 1980 cryptodome. Travel paths of rockslide blocks are interpreted from a geologic map of the deposit. The material was fractured and dilated during the rockslide, after which grain-to-grain dispersive stress facilitated flow. During transport, the dilated material mixed but significant fine material was not produced.

### Introduction

One of the most important events of the May 18, 1980, eruption of Mount St. Helens was the rockslide-debris avalanche. The rockslide was triggered by a M=5.1 earthquake at 08:32 PDT, which depressurized the volcano's magmatic and hydrothermal system, resulting in the lateral blast. Eyewitness photographs document that the rockslide began with detachment of at least three slide blocks which accelerated to a maximum of 70 m·s<sup>-1</sup>. The slide blocks broke up into smaller pieces to become a flowing debris avalanche, moving at an average rate of 35 m·s<sup>-1</sup> (Voight, 1981; Voight *et al.*, 1983). The resulting hummocky deposit has a volume of

2.5 km<sup>3</sup> (Figure 1). The Mount St. Helens rockslide-debris avalanche is the largest known mass movement in historic times.

In the ten years since the eruption, detailed field, laboratory, and modelling work has provided a comprehensive understanding of a previously poorly-understood type of event. The work includes analyses of the eyewitness photographs (Voight, 1981), the stability of the pre-eruption mountain (Voight *et al.*, 1983), the geology and emplacement of the deposit (Glicken, 1986, in press-a), the stability of debris-avalanche dams formed by the deposit (Glicken *et al.*, 1989a; Glicken and Voight, in press; Meyer *et al.*, 1985; Meyer *et al.*, 1986) and microscopic characteristics of the deposit with implications for particle-particle interactions (Glicken *et al.*, 1989b). Results of the Mount St. Helens work continue to be applied to the study of volcanic debris avalanches, associated eruptions, and hazards around the world (*e.g.*, Boudon *et al.*, 1987; summarized in Siebert *et al.*, 1987).

### Stability Model

A static mathematical model of the stability of the pre-eruption mountain indicates that the mountain was stable under reasonable assumptions of cohesion and water table conditions (Voight *et al.*, 1983). Reduction of cohesion, resulting primarily from intrusion of the March-May 1980 cryptodome and associated hydrothermal fluids, and dynamic loading resulting from the M=5.1 earthquake were required for failure.

### Composition of the Deposit

To analyze the debris avalanche deposit, it is necessary to define terms rigorously (Glicken, in press-b). Two different kinds of particles compose the deposit: clasts, which are rocks that would not break if passed through a sieve or immersed in water (hard rocks), and debris-avalanche blocks, which are unconsolidated or semi-consolidated (relatively soft) pieces of the pre-1980 mountain transported relatively intact. The parts of the deposit composed of debris-avalanche blocks are called block facies; the parts of the deposit composed of a blended mix of rock types from the pre-1980 mountain, juvenile material (cryptodome) and, locally, material from the underlying terrain are called mixed facies (previously referred to as matrix facies; *e.g.*, Ul, 1983).

In debris-avalanche blocks (debris blocks for brevity), recognizable structures or stratigraphy preserved from the pre-eruption Mount St. Helens are locally present (Figure 2). However, the clasts within the debris blocks are shattered, so the deposit is finer grained than the comparable material of the pre-1980 mountain.

The eastern half of the deposit is composed almost entirely of block facies (Figures 1 and 4), while the western part consists of primarily mixed facies with some debris



**Figure 1** Photograph of May 18, 1980, debris avalanche deposit, with Mount St. Helens in the background. Photograph taken summer 1980, before fluvial erosion. Note correlation of light and dark rocks of debris blocks in deposit with rocks of the crater wall.



**Figure 2** Canyon of North Fork Toutle River in 1984, showing debris block with preserved, faulted, lava flow stratigraphy. Debris-avalanche deposit is covered by blast deposit. Canyon is 50 m deep.

blocks (Figure 4). The block facies is divided into five lithologic units based on rock types of the pre-eruption Mount St. Helens. The juvenile material in the mixed facies indicates that the mixed facies was derived from the explosions of the cryptodome as well as from disaggregation and mixing of debris blocks.

**Relationship of Slide Blocks to Debris Avalanche Deposit**

The geologic cross-section of the pre-eruption mountain, with the slide blocks superimposed (Figure 3), together with the geologic map of the deposit (Figure 4) can be used to interpret the travel paths of the three slide blocks. Slide block I was a simple slide, triggered by the earthquake, and was deposited in Spirit Lake, on the ridge just north of the mountain, and flowed as a debris avalanche down the valley of the North Fork Toutle River. The exploding cryptodome burst through slide block II to produce the "blast surge" (Fisher *et al.*, 1987) that travelled faster than, and overtook, slide block I; the remainder of slide block II travelled down the valley, pushing slide block I into the margins of the valley. Slide block III failed as the cryptodome was still exploding. The continuing explosions disrupted large segments of the moving slide block, breaking it into debris blocks, and disaggregating debris blocks. Exploding fragments of juvenile dacite of the

cryptodome mixed with the disaggregating blocks. Slide block III, therefore, consisted of a travelling mass of debris blocks engulfed by, and intermixed with, disaggregated, mixed material. The denser parts of slide block III were deposited as block facies in the

eastern part of the North Fork Toutle River valley, while most of the disaggregated, mixed parts continued to flow westward to produce the bulk of the mixed facies of the western part of the deposit. The cryptodome continued to depressurize after slide block




Figure 3 N-S cross-section of Mount St. Helens just before eruption, showing slide blocks I, II, and III visible in eyewitness photographs. Geology of pre-1980 eruption mountain from C.A. Hopson, summarized in Glicken (1986, in press-a).




Figure 4 Generalized map of debris avalanche deposit. Interpretations of resting places of slide blocks shown by roman numerals. III (II) indicates primarily slide block III with subordinate volume of slide block II. Generalized from 1:12,000 map of Glicken (1986). H stands for Harrys Ridge.




Figure 5 Summary of sedimentary processes in rockslide-debris avalanche transport.

III left the crater (minutes after 08:32 PDT), resulting in a juvenile dacite-rich pyroclastic flow deposit that rests on top of the debris-avalanche deposit as far as 20 km from the crater (Criswell, 1987). After deposition, major lahars were produced by dewatering of water-saturated parts of the debris-avalanche deposit.

**Sedimentary Processes**

Sedimentary processes of the rockslide-debris avalanche (how the material moved, broke up, and was deposited) are summarized in Figure 5. The mean density of the pre-1980 cone was determined by laboratory specific gravity measurements of individual clasts and in-place sand-cone density to be about 20% greater than the mean density of the avalanche deposit measured by the sand-cone density technique (Glicken, 1986, in press-a). The density of the deposit does not decrease with distance, suggesting that debris blocks were fractured and dilated at the mountain, rather than during transport.

Recent work (Komorowski *et al.*, in prep.) shows that waves of compression and rarefaction propagated from the base of the moving rockslide fractured and dilated the material. The material remained dilated because of grain-to-grain dispersive stress, which facilitated flow. The fracturing resulted in jigsaw cracks that are visible at the micron scale in SEM images (Glicken *et al.*, 1989b), but, unlike other debris avalanches, are rarely seen in exposures in the field. Various grain-size parameters show that during transport, the dilated material mixed together, but significant amounts of fine material were not produced.

**Conclusions**

Mount St. Helens provided a serendipitous confluence of pre-eruption studies, observations and photographs made during the early moments of sliding of the rockslide blocks,

and an opportunity to recognize and analyze each specific component of the debris-avalanche deposit. Each has been vital in gaining an understanding of the mechanisms of failure, transport, and deposition of the immense rockslide-debris avalanche of May 18, 1980. The processes are more common at volcanoes than had been realized previously, and application of the work of Mount St. Helens contributes to improved assessments of hazards at volcanoes throughout the world.

**Acknowledgements**

Thanks to the U.S. Geological Survey for supporting most of this work, and to C. Dan Miller, Cathie Hickson, and Don Peterson for constructive reviews.

**References**

Boudon, G., Semet, M.P. and Vincent, P.M., 1987, Magma and hydrothermally-driven sector collapses: the 3100 and 11,000 year b.p. eruptions of la Grande Decouverte (la Soufrière) volcano, Guadeloupe, French West Indies: *Journal of Volcanology and Geothermal Research*, v. 33, p. 317-323.

Criswell, C.W., 1987, Chronology and pyroclastic stratigraphy of the May 18, 1980, eruption of Mount St. Helens, Washington: *Journal of Geophysical Research*, v. 92, p. 10,237-10,266.

Fisher, R.V., Glicken, H. and Hoblitt, R.P., 1987, May 18, 1980, Mount St. Helens deposits in South Coldwater Creek, Washington: *Journal of Geophysical Research*, v. 92, no. B10, p. 10,267-10,283.

Glicken, H., 1986, Rockslide-debris avalanche of May 18, 1980, Mount St. Helens Volcano, Washington, Ph.D. Dissertation, University of California, Santa Barbara, California, 304 p.

Glicken, H., in press-a, Rockslide-debris avalanche of May 18, 1980, Mount St. Helens Volcano, Washington: United States Geological Survey, Professional Paper 1488, 304 p.

Glicken, H., in press-b, Sedimentary architecture of large volcanic debris avalanches, in Fisher, R.V. and Smith, G.A., eds., *Sedimentation in Volcanic Settings: Society of Economic Paleontologists and Mineralogists Special Publication*, in press.

Glicken, H., Komorowski, J.-C. and Sheridan, M., 1989b, Particle-particle interaction in the Mount St. Helens debris avalanche: *Scanning Electron Microscope Approach: EOS*, v. 70, p. 1423.

Glicken, H., Meyer, W. and Sabol, M.A., 1989a, Geology and ground-water hydrology of Spirit Lake Blockage, Mount St. Helens, Washington, with implications for lake retention: United States Geological Survey, Bulletin 1789, 33 p.

Glicken, H., and Voight, B., in press, Geology of the North Fork Toutle River Blockages, in Schuster, R.L. and Meyer, W., eds., *Major Lakes Impounded by the May 18, 1980, Eruption of Mount St. Helens, and the Stability of Their Blockages: United States Geological Survey, Professional Paper*, in press.

Komorowski, J.-C., Glicken, H. and Sheridan, M.F., in prep., Secondary electron imagery of micro-cracks and hackly fractures in sand-size clasts from the 1980 Mount St. Helens debris avalanche deposit: implications for particle-particle interactions.

Meyer, W., Sabol, M.A., Glicken, H.X. and Voight, B., 1985, The effects of gravitational and earthquake forces on the stability of the South Fork Castle Creek Blockage in the Mount St. Helens area, Washington: United States Geological Survey, Professional Paper 1345, 42 p.

Meyer, W., Sabol, M.A. and Schuster, R.L., 1986, Landslide-dammed lakes at Mount St. Helens, Washington, in Schuster, R.L., ed., *Landslide Dams: Processes, Risk, and Mitigation: American Society of Civil Engineers, Geotechnical Special Publication* no. 3, p. 21-41.

Siebert, L., Glicken, H. and Ui, T., 1987, Volcanic hazards of Bezymianny- and Bandai-type eruptions: *Bulletin of Volcanology*, v. 49, p. 435-459.

Ui, T., 1983, Volcanic debris avalanche deposits — identification and comparison with non-volcanic debris stream deposits: *Journal of Volcanology and Geothermal Research*, v. 18, p. 135-150.

Voight, B., 1981, Time scale for the first moments of the May 18 eruption, in Lipman, P.W. and Mullineaux, D.R., eds., *The 1980 Eruptions of Mount St. Helens*, Washington: United States Geological Survey, Professional Paper 1250, p. 69-86.

Voight, B., Janda, R.J., Glicken, H. and Douglass, P.M., 1983, Nature and mechanics of the Mount St. Helens rockslide-avalanche of 18 May 1980: *Geotechnique*, v. 33, p. 243-273.