

Caribbean Geology: Extended and Subsided Continental Crust Sharing History with Eastern North America, the Gulf of Mexico, the Yucatán Basin and Northern South America

K. H. James

Volume 40, numéro 1, 2013

URI : https://id.erudit.org/iderudit/geocan40_1gd01

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)

1911-4850 (numérique)

[Découvrir la revue](#)

Citer ce document

James, K. H. (2013). Caribbean Geology:: Extended and Subsided Continental Crust Sharing History with Eastern North America, the Gulf of Mexico, the Yucatán Basin and Northern South America. *Geoscience Canada*, 40(1), 3–8.

GEOSCIENCE DEBATE

Caribbean Geology: Extended and Subsided Continental Crust Sharing History with Eastern North America, the Gulf of Mexico, the Yucatán Basin and Northern South America

K. H. James

Honorary Departmental Fellow
Institute of Geography and Earth Sciences
Aberystwyth University, Wales, UK,
SY23 3DB
E-mail: khj@aber.co.uk

THE PROBLEM

The geology of Middle America is complicated by its dispersal over many geographic elements (Fig. 1). Some are well studied (Gulf of Mexico and Colombia – Venezuela – Trinidad hydrocarbon provinces). Others remain poorly known because of lack of study, difficult access, tropical vegetation/weathering or large submarine extension. The area is largely submarine but there is no clear evidence of ‘oceanic spreading’ (ridges, magnetic anomalies) apart from the centre of the Cayman Trough. So, is the area composed of allochthonous oceanic crust or *in situ* continental crust, extended in place between the Americas?

THE PARADIGM

Popular understanding (e.g. Pindell and Kennan 2009) is that Jurassic oceanic Caribbean crust formed in the Pacific

(Fig. 2). It thickened over a hotspot/mantle plume (Galapagos, or Sala y Gomez, or both; Révillon et al. 2000), or above a ‘slab gap’ as it overrode the Proto-Caribbean (Pindell and Kennan 2009) forming an ‘oceanic plateau’ or large igneous province. Drifting eastwards relative to the Americas, it collided with a linear, west-facing intra-oceanic volcanic arc, reversing its subduction polarity and driving it between the Americas to collide progressively with North and South America during the Late Cretaceous – Eocene, dying in the process. The Lesser Antilles, formed in the Eocene after subduction jump from the Aves Ridge, are seen as the remnants of this arc. The Chortis and Maya blocks are seen to have rotated clockwise or anticlockwise by up to 180° from the Gulf of Mexico or southern Mexico.

ANALOGUES

The Caribbean Plate does not reveal classic oceanic spreading ridges or magnetic anomalies; except for the centre of the Cayman Trough (north – south anomalies; 300 km of Oligocene – Recent east – west spreading, 100 km north – south opening), the normal signature of oceanic origins is lacking. However, the Scotia Plate in the South Atlantic Ocean, and the Banda Sea Plate in Indonesia, which do carry ridges and identified magnetic stripes, are remarkably similar to the Caribbean and their geology is highly relevant (Fig. 10 in James 2005).

The Scotia and Banda Sea plates formed in place (Barker 2001; Honthaas et al. 1998). They lie, like the

Caribbean Plate, between sinistrally-offset, major continental blocks to the north and south. Each is around 3000 km long and 700–800 km wide. Each has a curved volcanic arc in the east. Northwest-trending volcanic arcs in the west follow the continental Sumatra/Java block on the western edge of the Banda region and the Chortis Block of the Caribbean Plate, while the Scotia Plate is bounded by the shallow (700 m), northwest-trending Shackleton Fracture Zone, built of continental slivers (Livermore et al. 2004), has active volcanoes in the Drake Passage. Fractures in the oceans east and west of the plates diverge towards them, indicating extension.

Magnetic data show that extension in the Scotia Plate distributed continental blocks that originally connected South America and Antarctica (Fig. 3; Barker 2001). The North and South Scotia ridges carry continental rocks and elevated parts of the Scotia Sea may be continental crust thinned by extension. The Neogene Banda Sea also opened in an extensional setting (Hall 1997). Dredged sedimentary and metamorphic rocks show that internal ridges are continental slivers.

Is there evidence that the Caribbean Plate carries continental crust? James (2009a) presented a comprehensive synthesis of data (regional tectonic fabric (Fig. 3), crustal thicknesses, gravity and magnetic data, stratigraphy, paleontology, highly silicic chemistry, ancient zircon grains in Cretaceous arc rocks) from the whole of Middle America; a separate article (James 2009b) interpreted these data. They all converge to show a regional



Figure 1. Middle America. Intracontinental Gulf of Mexico is surrounded by southern North America and the Florida – Bahamas, Tehuantepec (T) and Campeche (Maya) platforms. Cuba (C), with basement and Mesozoic carbonate cover related to the Florida – Bahamas platform, bounds the Yucatán Basin to the north, and the Cayman Ridge separates it from the Cayman Trough, a 1200 km long pull-apart structure in the sinistral boundary between North America and the Caribbean. The Chortis Block (CB), with its marine extension, the Nicaragua Rise (NR), limited to the south by the Hess Escarpment (HE), forms about a third of the Caribbean Plate and is the only place where continental crust is generally recognized. The Chorotega and Chocó blocks (Chr, Chc) link the Chortis Block to South America; for the most part, they comprise volcanic arc and accreted oceanic rocks. Crustal thickness (up to 45 km), gravity data and high silica chemistry indicate continental roots. The active Lesser Antilles volcanic arc is generally seen as the remnant of a larger, Cretaceous arc (Fig. 2) whose extinct parts occur in the Greater Antilles, northern South America and the Aruba – Blanquilla (AB) islands. Jamaica (J), Hispaniola (H), Puerto Rico (PR) and the northern Virgin Islands are large, thick (30 km), mostly submerged blocks separated by narrow deeps. Gravity data and high silica content again indicate continental roots. The Beata and Aves Ridges (BR, AR) separate the Colombian – Venezuelan and the Venezuelan – Grenada basins. Sinistral and dextral strike-slip motion on the wide (several hundred kms) northern and southern plate boundaries accommodate westward movement of North and South America relative to the Caribbean. The Pacific Cocos Plate converges northeastwards with the Chocó Block/northwestern South America, driving them over the western Caribbean and narrowing the area (James 2007a). DSDP: Deep Sea Drilling Project.

history of continental extension during Jurassic–Cretaceous divergence of North and South America.

TECTONIC SETTING AND FABRIC

As spreading occurred in the Central Atlantic, the westerly convex Mid-Atlantic Ridge moved away from Africa and extended N-S over

Caribbean latitudes. The Caribbean Sea lies in an area of divergence between the central and northern South Atlantic (Figs. 3, 4, 5 in Fairhead and Wilson 2004; Figs. 3, 4B in Fairhead and Wilson 2005). Fracture patterns indicate that much of this extension took place in the Cretaceous and it is noteworthy that basalt extrusion occurred in the

Caribbean at 120, 80–90 and 76 Ma.

EVOLUTION

Following development of the Central Atlantic Magmatic Province in the Late Triassic, extension resulted in sinistral offset of North America from South America along N60°W fractures/intra-continental faults, while the South Atlantic remained stable until the late Early Cretaceous. N 35° E Palaeozoic compressional structures along eastern North America (Appalachians), through Middle America (Cuba, Maya and Chortis blocks; Fig. 1) and into northern South America (Perijá, Maracaibo, Mérida, Espino grabens; Fig. 3) were reactivated as Triassic – early Jurassic rifts and dextral faults. N 60° E normal faults (e.g. Hess Escarpment; Fig. 1) and the Florida Arch formed and sinistral slip occurred along the northern Caribbean boundary. This early ‘Cayman’ displacement of around 600 km – the offset between the Maya and Chortis blocks recorded by their eastern margins and Jurassic grabens – did not affect the southern boundary. Rifting continued in the Jurassic – Cretaceous, evolving to (listric) fault block rotation accommodating wedges of continental red beds, marine source rocks and salt.

Subsidence of extended continental crust was matched by thick Cretaceous carbonate platform growth along eastern North America (up to 9 km in the Bahamas), around the Gulf of Mexico and in the Caribbean. Correlation of sections in the north of Cuba and Hispaniola with the Bahamas shows that these areas are in place, not allochthonous. Figure 4 shows the resultant geology of present-day offshore North America farther north. It serves as a template for Caribbean geology, as indicated by seismic data, but here we have to add the Cretaceous basalt extrusions mentioned earlier.

Regional Late Cretaceous–Paleogene compression occurred around the Caribbean, culminating violently with Middle Eocene emplacement of major allochthons northward onto Cuba (e.g. 1000 km-long serpentinite sheets), and southward in Venezuela (the 250 x 25 x 5 km Villa de Cura volcanic complex). Regional occurrence of associated deposits




Figure 2. Model for Caribbean Plate (CAP) migration (red arrows) from the Pacific. Central America, outlined, was not present until Maya and Chortis rotated from the Gulf of Mexico and southwest Mexico (black arrow) and, together with intra-oceanic volcanic arc and oceanic rocks of the Chorotega and Chocó blocks (Fig. 1), accreted to the western tail of the Caribbean Plate. Note impossible bending of a linear volcanic arc, which must be rooted in crust, into an extreme curve. Note also that in the Lesser Antilles (Fig. 1) supposedly Eocene – Recent in age, there are Lower Cretaceous arc rocks in the northern islands (Bouysse et al. 1983). Brm – Barremian; Tur – Turonian; Mas – Maastrichtian; MEoc – Middle Eocene (orange line); MMio – Middle Miocene (yellow line).

(wildfysch) and of overlying Middle Eocene shallow marine carbonate units again demonstrates the internal coherence and autochthonous nature of Caribbean elements.

In the late Eocene – Oligocene the Caribbean plate began east-west extension, with sinistral and dextral movement occurring along its northern and southern boundaries. This was a second sinistral event (the ‘Central Cayman’) on the northern boundary. It did not extend as far as the Pacific Plate and so the upper Cretaceous to Recent section in the Gulf of Tehuantepec is undisturbed. It resulted in some 300 km of east – west pull-apart extension between the Greater and Leeward Antillean islands. Sinistral and dextral displacement along the northern and southern Caribbean boundaries continues today, generating eastward-migrating uplift and complementary subsidence (spectacularly seen along northern Venezuela, where the -200 mgal gravity anomaly over the Maturín Basin, in the east, is the world’s largest negative anomaly at sea level).

In the southwestern Caribbean, convergence of the Pacific Chocó area drives northwestern South America and southern Central America to the northeast. The result is northward movement of the Panama arc and the Maracaibo Block over the southwest Caribbean, driving the South Caribbean Deformed Belt northwards. The +200 mgal gravity anomaly associated with the 5800m high Sierra Nevada de Santa Marta in northwest Colombia, the world’s highest elevation adjacent to sea floor (3000 m deep), demonstrates the dynamic nature of this area. Contrary to models of southward Caribbean subduction (there is no volcanism and GPS data show that the Caribbean Plate is moving eastward), the area is delaminating and moving over the Caribbean, which has continental composition (Ceron-Abril 2008). As mentioned above, Fairhead and Wilson (2004) used gravity, magnetic data and fractures to show that divergence occurs between the central and northern South Atlantic – there is no convergence over the Caribbean area.




Figure 3. Tectonic fabric. Middle America shows a regional structural pattern that demonstrates shared tectonic history. There is no indication in the Caribbean Plate of the radial fabric expected above plumes. Jurassic grabens crossing the Maya (RHF – Rio Honda Fault) and Chortis (GF – Guayape Fault) blocks remain parallel to each other and to rifts in North (BRR – Blue Ridge Rift) and South America (PG – Perija Graben, MG – Merida Graben, EG – Espino Graben), demonstrating that the Maya and Chortis blocks have not rotated.

THE CARIBBEAN ‘OCEANIC PLATEAU’

The western part of the Venezuela Basin crust is up to 20 km thick and is generally interpreted as an ‘oceanic plateau’ or large igneous province. The Kerguelen, Iceland, Jan Mayen, Ontong Java, Rockall, and Seychelles ‘oceanic plateaus’ are known, from ancient zircon grains and dredged rocks, to be underpinned by continental crust (e.g. Foulger et al. 2005; Schaltegger et al. 2002).

Magnetic lineations over the area 1) trend northeast – southwest, 2) conform to regional tectonic fabric (Fig. 3), 3) reflect structure, not sea-floor spreading (Donnelly 1989; Diebold et al. 1999), and 4) extend the Triassic – Jurassic rift trends of North and South America. Seismic data show buried highs and ridges flanked by wedges of dipping reflections extending from 20 km to over 100 km. Pacific ‘oceanic’ models see these as volcanic flows and seamounts (Diebold et al. 1999).

Five Deep Sea Drilling Project holes have sampled the Caribbean




Figure 4. Cross-section (see inset for location) offshore eastern North America, based on seismic and borehole data (after Benson and Doyle 1988; Manspeizer 1988). The section shows rifted basement, with red beds, followed by sediment wedges (blue), salt (pink, cross-hatched), limestones (blue, brick), pale blue wedges of Middle Jurassic – Early Cretaceous sediments, followed by Upper Cretaceous – Paleogene (green) and Neogene (yellow) sediments.




Figure 5. *In situ* interpretation (James 2007b) of seismic line 1293 (location in Fig. 3) over the Caribbean ‘plateau’. See Diebold et al. 1999 (Figs. 2, 15) for original seismic line and ‘oceanic’ interpretation of 40 km-wide highs of vertical dykes flanked by volcanic flows, with local seamounts. The latter are shown here as salt diapirs (inset compares seismic over the Caribbean diapir, left, with the drilled Challenger Knoll salt dome, right, of the Gulf of Mexico) and ‘volcanic wedges’ are seaward-dipping reflections recording continental extension (cf. Fig. 4). M-U Cret: Middle to Upper Cretaceous; U-Jur – L-Cret: Middle to Upper Jurassic to Lower Cretaceous; M-U Jur: Middle to Upper Jurassic.

‘plateau’, terminating after a few metres in basalts or diorite sills dated 88–90 Ma (Saunders et al. 1973; Kerr et al. 2003). The basalt is vesicular and smooth with “morphology reminiscent

of continental basalts” (Diebold et al. 1999); overlying sediments are shallow water in origin. The interpretation of Caribbean ‘plateau’ seismic data in the light of extended continental crust,

such as seen offshore Europe and along eastern North America, appears in Figure 5 (James 2007b).

THE STATE OF PLAY

Pacific models become ever more complex and numerous. There are at least six different models for opening of both the Grenada and Yucatán basins. There is disagreement over the existence and timing of the proposed Cretaceous arc polarity reversal and the number and polarity of Cretaceous arcs in the Caribbean. The presence of ancient zircon grains in Cretaceous arc rocks is explained by subduction of continental crust by the migrating/colliding arc and by (never illustrated) resurfacing in Cuba and along northern Venezuela. Blueschist is supposed to have formed in similar settings, yet it is interbedded with coeval sedimentary rocks in Cuba and Venezuela and is absent from the active subduction zones of the Lesser Antilles and Central America, in existence since at least the Early Cretaceous.

The new 'Pirate' model (Keppie 2013) proposed that the Caribbean has grown in the northwest and southwest by capturing microplates from northwest South America and the Gulf of Mexico. This model appears to regard the bulk of the Caribbean Plate to have formed in place (Fig. 4 in Keppie 2013) and so is a variant of the *in situ* model. The main debate remains 'Pacific' or '*in situ*'. As for 'capture' of northwest South America, Ceron-Abril (2008) showed how the area is delaminating and overriding the Caribbean continental crust. Jurassic rifts that parallel the regional tectonic fabric of Middle America deny rotation of the Chortis Block from the Gulf of Mexico.

THE POINT OF THIS PAPER

The main thrust of this article is to support the *in situ*, extended continental origin of the Caribbean Plate and to counter the 'Pacific paradigm'. However, its lessons extend globally.

Many assumptions preface work in the oceans: "The ocean basins provide a unique opportunity to investigate magmatic processes and mantle composition. The absence of continental crust as a potential contaminant provides an untarnished sample of the deep Earth" (Saunders and Norry 1989). "Intra-oceanic arcs are important for geochemical studies of basalts because contamination by continent

cannot have occurred" (Leat and Larter 2003).

Assumptions that continental crust is absent from the oceans are not justified. Many samples of continental rocks have been recovered from the oceans, such as those recovered by Woods Hole and the ODP (Ewing 1949; Ewing et al. 1948; Vasiliev and Yano 2007). The Lesser Antilles arc has continental thickness, produces rocks with silica content up to 76%, including quartz diorite and granite on La Désirade, and rounded quartz grains. Basalt compositions are directly comparable in chemical composition and mineralogy with those from continental margin orogenic belts (Lewis 1971). Seismic and gravity data show continental thicknesses and densities in the Central America arc, which produces high silica volcanic rocks and continental xenoliths. Original continental crust is present below both arcs. The northern Izu-Ogasawara 'intra-oceanic island arc' is underlain by continental crust (granite and andesite; Suyehiro et al. 1996). 'Subduction factories' of continental crust (Tatsumi and Kosigo 2003) and complex derivation of andesites (the 'andesite problem') from basalt (Takahashi et al. 2007) below 'intra-oceanic arcs' are not necessary.

ENDPOINT

While the Caribbean is held to have Pacific, oceanic origins, nobody will look for hydrocarbons there. The area lies between the prolific oil provinces of the Gulf of Mexico and northern South America. Oil occurrences/shows occur on surrounding islands and in Central America.

ACKNOWLEDGEMENTS

I am grateful for the invitation from Geoscience Canada to contribute to scientific debate on the origin of the Caribbean Plate. I thank Fraser Keppie and the editors for constructive comments.

REFERENCES

Barker, P.F., 2001, Scotia Sea regional tectonic evolution: implications for mantle flow and palaeocirculation: *Earth-Science Reviews*, v. 55, p. 1–39, [http://dx.doi.org/10.1016/S0012-8252\(01\)00055-1](http://dx.doi.org/10.1016/S0012-8252(01)00055-1).

- Benson, R.N., and Doyle, R.G., 1988, Early Mesozoic rift basins and the development of the United States middle Atlantic continental margin, *in* Manzpeizer, W., *ed.*, Triassic–Jurassic rifting, continental breakup, and the origin of the Atlantic Ocean and passive margins: Elsevier, Amsterdam, p. 99–127.
- Bouysse, P., Schmidt-Effing, R., and Westercamp, D., 1983, La Desirade Island (Lesser Antilles) revisited: Lower Cretaceous radiolarian cherts and arguments against an ophiolitic origin for the basal complex: *Geology*, v.11, p. 244–247, [http://dx.doi.org/10.1130/0091-7613\(1983\)11<244:LDILAR>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(1983)11<244:LDILAR>2.0.CO;2).
- Ceron-Abril, J.F., 2008, Crustal structure of the Colombian Caribbean Basin and margins: Unpublished Ph.D. thesis, University of South Carolina, Columbia, 165 p.
- Diebold, J.B., Driscoll, N.W., and the EW-9501 Science Team, 1999, New insights on the formation of the Caribbean Basalt Province revealed by multichannel seismic images of volcanic structures in the Venezuelan Basin, *in* Mann, P. *ed.*, Caribbean Basins. Sedimentary Basins of the World, 4 (Series Editor: K.J. Hsü): Elsevier, Amsterdam, p. 561–589.
- Donnelly, T.W., 1989, Geologic history of the Caribbean and Central America, *in* Bally, A.W., and Palmer A.R., *eds.*, The Geology of North America – An Overview: The Geology of North America, v. A: Geological Society of America, Boulder, Colorado, p. 299–321.
- Ewing, M., 1949, New Discoveries on the Mid Atlantic Ridge: *National Geographic Magazine*, September, v. XCVI No. 5, p. 611–640.
- Ewing, M., Walker, J., Henry, T.R., Llocke, J.N., Watson, D. Culver, W.R., Vosburg, F.G., Stewart, B.A., and Wharton, C.H., 1948, Exploring the Mid Atlantic Ridge: *National Geographic Magazine*, September, v. XCIV No. 3, p. 275–294.
- Fairhead, J.D., and Wilson, M., 2004, Seafloor spreading and deformation processes in the South Atlantic Ocean: Are hot spots needed?: *MantlePlume.org*, University of Durham, U.K. Available from <http://www.mantleplumes.org/SAtlantic.html>.
- Fairhead, J.D., and Wilson, M., 2005, Plate tectonic processes in the South Atlantic Ocean: Do we need deep mantle plumes?, *in* Foulger, G.R., Natland, J.H., Presnall, D.C., and Ander-

- son D.L., *eds.*, Plates, plumes, and paradigms: Geological Society of America Special Paper, 388, p. 537–553, <http://dx.doi.org/10.1130/0-8137-2388-4.537>.
- Foulger, G. R., Natland, J.H., and Anderson, D.L., 2005, Genesis of the Iceland melt anomaly by plate tectonic processes, *in* Foulger, G.R., Natland, J.H., Presnall, D.C., and Anderson, D.L., *eds.*, Plates, plumes and paradigms: Geological Society of America Special Paper 388, p. 595–625, <http://dx.doi.org/10.1130/0-8137-2388-4.595>.
- Hall, R., 1997, Cenozoic plate tectonic reconstruction of SE Asia, *in* Fraser, A.J. Matthews, S.J., and Murphy, R.W., *eds.*, Petroleum Geology of Southeast Asia: Geological Society of London Special Publication, 126, p. 11–23.
- Honthaas, C., Réhault, J-P., Maury, R.C., Bellon, H., Hémond, C., Malod, J-A., Cornée, J-J., Villeneuve, M., Cotten, J., Burhanuddin, S., Guillou H., and Arnaud, N., 1998, A Neogene back-arc origin for the Banda Sea basins: Geochemical and geochronological constraints from the Banda ridges (East Indonesia): Tectonophysics, v. 298, p. 297–317, [http://dx.doi.org/10.1016/S0040-1951\(98\)00190-5](http://dx.doi.org/10.1016/S0040-1951(98)00190-5).
- James, K.H., 2005, Arguments for and against the Pacific origin of the Caribbean Plate and arguments for an in situ origin: Transactions of the 16th Caribbean Geological Conference, Barbados, Geological Society of Jamaica, Caribbean Journal of Earth Sciences, v. 39, p. 47–67. Available from <http://caribjes.com/>.
- James, K.H., 2007a, Structural Geology: From local elements to regional synthesis, *in* Bundschuh, J., and Alvarado, G.E., *eds.*, Central America: Geology, Resources and Hazards: Taylor and Francis/Balkema, The Netherlands, Chapter 11, p. 277–321.
- James, K.H., 2007b, The Caribbean Ocean plateau: MantlePlume.org, University of Durham, U.K. Available from <http://www.mantleplumes.org/Caribbean.html>.
- James, K.H., 2009a, In-situ origin of the Caribbean: Discussion of data, *in* James, K.H., Lorente, M.A., and Pindell, J., *eds.*, Origin and evolution of the Caribbean Plate: Geological Society of London Special Publications, 328, p. 75–124.
- James, K.H., 2009b, In-situ origin of the Caribbean: Interpretation of data, *in* James, K.H., Lorente, M.A., and Pindell, J., *eds.*, Origin and evolution of the Caribbean Plate, Geological Society of London Special Publications, 328, p. 125–36.
- James, K.H., 2010, Observations on new magnetic map from the Commission for the Geological Map of the World: New Concepts in Global Tectonics Newsletter, Issue 57, p. 14–26. Available from <http://www.ncgt.org/newsletter.php>.
- Keppie, F., 2013, The rationale and essential elements for the new Pirate model of Caribbean tectonics: Geoscience Canada, v.40, p. 9–16, <http://dx.doi.org/10.12789/geocanj.2013.40.002>.
- Kerr, A.C., White, R.V., Thompson, P.M.E., Tarney, J., and Saunders, A.D., 2003, No Oceanic Plateau - No Caribbean Plate? The Seminal Role of Oceanic Plateau(s) in Caribbean Plate Evolution, *in* Bartolini, C., Buffler, R.T., and Blickwede, J., *eds.*, The Gulf of Mexico and Caribbean Region: Hydrocarbon Habitats, Basin Formation and Plate Tectonics: American Association of Petroleum Geology Memoir, 79, p. 126–168.
- Leat, P.T., and Larter, R.D., 2003, Intra-oceanic subduction systems: Introduction, *in* Larter, R.D., and Leat, P.T., *eds.*, Intra-Oceanic Subduction Systems: Tectonic and Magmatic Processes: Geological Society of London Special Publication, 219, p. 1–17.
- Lewis, J.F., 1971, Composition, origin, and differentiation of basalt magma in the Lesser Antilles, *in* Donnelly, T.W., *ed.*, Caribbean Geophysical, Tectonic and Petrologic Studies: Geological Society of America Memoir 130, p.159–179.
- Livermore, R., Eagles, G., Morris, P., and Maldonado, A., 2004, Shackleton Fracture Zone: No barrier to early circum-polar ocean circulation: Geology, v. 32, p. 797–800, <http://dx.doi.org/10.1130/G20537.1>.
- Manspeizer, W., 1988, Triassic-Jurassic rifting and opening of the Atlantic: An Overview, *in* Manspeizer, W., *ed.*, Triassic-Jurassic rifting, continental breakup, and the origin of the Atlantic Ocean and passive margins: Elsevier, Amsterdam, p. 41–79.
- Pindell, J., and Kennan, L., 2009, Tectonic evolution of the Gulf of Mexico, Caribbean and northern South America in the mantle reference frame: an update (abstract): Workshop, Cardiff, Wales, UK, Circum-Caribbean and North Andean tectonomagmatic evolution: impacts on palaeoclimate and resource formation, Abstracts, p. 35.
- Révilion, S., Hallot, E., Arndt, N.T., Chauvel, C., and Duncan, R.A., 2000, A Complex History for the Caribbean Plateau: Petrology, Geochemistry, and Geochronology of the Beata Ridge, South Hispaniola: The Journal of Geology, v. 108, p. 641–661, <http://dx.doi.org/10.1086/317953>.
- Saunders, A.D., and Norry, M.J., 1989, Introduction, *in* Saunders, A. D. and Norry, M.J., *eds.*, Magmatism in the Ocean Basins: Geological Society of London Special Publication, 42, p. vii–viii.
- Saunders, J.B., Edgar, N.T., Donnelly, T.W., and Hay, W.W., 1973, Cruise synthesis, *in*: Edgar, N.T. and others, *eds.*, Initial Reports of the Deep Sea Drilling Project: U.S. Government Printing Office, Washington, D.C., v.15, p. 1077–1111, <http://dx.doi.org/10.2973/dsdp.proc.15.140.1973>.
- Schaltegger, U., Amundsen, H., Jamtveit, B., Frank, M., Griffin, W.L., Grönvold, K., Trönnnes, R., Torsvik, T., 2002, Contamination of OIB by underlying ancient continental lithosphere: U–Pb and Hf isotopes in zircons question EM1 and EM2 mantle components (abstract): Goldschmidt Conference Abstracts 2002, Geochimica et Cosmochimica Acta 66, Supplement 1, p. A673, [http://dx.doi.org/10.1016/S0016-7037\(02\)01012-8](http://dx.doi.org/10.1016/S0016-7037(02)01012-8).
- Suyehiro, K., Takahashi, N., Ariie, Y., Yokoi, Y., Hino, R., Shinohara, M., Kanazawa, T., Hirata, N., Tokuyama H., and Taira, A., 1996, Continental crust, crustal underplating, and Low-*Q* upper mantle below an oceanic island arc: Science, v. 272, p. 390–392, <http://dx.doi.org/10.1126/science.272.5260.390>.
- Takahashi, N., Kodaira, S., Klemperer, S.L., Tatsumi, Y., Kaneda, Y., and Suyehiro, K., 2007, Crustal structure and evolution of the Mariana intra-oceanic island arc: Geology, v. 35, p. 203–206, <http://dx.doi.org/10.1130/G23212A.1>.
- Tatsumi, Y., and Kosigo, T., 2003, The subduction factory: its role in the evolution of the Earth's crust and mantle, *in* Larter, R.D., and Leat, P.T., *eds.*, Intra-Oceanic Subduction Systems: Tectonic and Magmatic Processes: Geological Society of London Special Publication, 219, p. 55–80.
- Vasiliev, B.I., and Yano, T., 2007, Ancient and continental rocks discovered in the ocean floors: New Concepts in Global Tectonics Newsletter, Issue 43, p. 3–17. Available from <http://www.ncgt.org/newsletter.php>.

Received March 2012

Accepted as revised February 2013