

Logan Medallist 7. Appinite Complexes, Granitoid Batholiths and Crustal Growth: A Conceptual Model

J. Brendan Murphy, William J. Collins et Donnelly B. Archibald

Volume 49, numéro 3-4, 2022

URI : <https://id.erudit.org/iderudit/1095214ar>
DOI : <https://doi.org/10.12789/geocanj.2022.49.191>

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)
1911-4850 (numérique)

[Découvrir la revue](#)

Citer cet article

Murphy, J., Collins, W. & Archibald, D. (2022). Logan Medallist 7. Appinite Complexes, Granitoid Batholiths and Crustal Growth: A Conceptual Model. *Geoscience Canada*, 49(3-4), 237-249.
<https://doi.org/10.12789/geocanj.2022.49.191>

Résumé de l'article

Les corps d'appinite sont une suite de roches plutoniques, de composition ultramafique à felsique, qui se caractérisent par de la hornblende idiomorphe comme minéral mafique dominant dans toutes les lithologies et par des textures spectaculairement diverses, y compris des fabriques magmatiques planaires et linéaires, des pegmatites mafiques et de nombreuses preuves de « mingling », mélange hétérogène, des compositions mafiques et felsiques de même âge. Ces caractéristiques suggèrent une cristallisation à partir d'un magma anormalement riche en eau qui, selon un nombre limité d'études isotopiques, possède à la fois des composants mantelliques et météoriques.

Les corps d'appinite se présentent généralement sous la forme de petits complexes (~ 2 km de diamètre) mis en place à la périphérie des plutons granitoïdes et généralement adjacents aux principales failles crustales profondes qu'ils exploitent préférentiellement lors de leur ascension. Plusieurs études soulignent la relation entre l'intrusion d'appinite, le plutonisme granitoïde et l'arrêt de la subduction. Cependant, des données géochronologiques récentes suggèrent une relation génétique de plus longue durée entre la génération d'appinite et de magma granitoïde et la subduction.

L'appinite peut représenter des aliquotes de magma basaltique hydraté dérivées de sous-plaques mafiques à fractionnement variable qui ont été initialement mises en place lors d'une subduction prolongée adjacente au Moho, déclenchant la génération de magma granitoïde volumineux par fusion partielle dans la zone MASH sus-jacente. Le magma mafique hydraté de cette sous-plaque peut avoir remonté et s'être accumulé et différencié à des niveaux crustaux moyens à supérieurs (environ 3 à 6 kbar, 15 km de profondeur) et avoir cristallisé dans des conditions de saturation en eau. Le magma granitoïde s'est mis en place par impulsions lorsque des contraintes transitoires ont activé des structures favorablement orientées qui sont devenues des conduits pour le transport du magma. L'ascension du magma mafique tardif, cependant, est entravée par les barrières rhéologiques créées par les corps magmatiques granitoïdes structurellement sus-jacents. Le magma qui forme des complexes d'appinite a échappé à ces barrières rhéologiques car il a exploité préférentiellement les failles crustales profondes qui délimitaient le système plutonique. Dans ce scénario, les complexes d'appinite peuvent être une connexion directe à la sous-plaque mafique et ainsi ses composants les plus mafiques peuvent fournir des informations sur les processus qui génèrent des batholites granitoïdes et, plus généralement, sur la croissance crustale dans les systèmes d'arc.

GAC MEDALLIST SERIES



Logan Medallist 7. Appinite Complexes, Granitoid Batholiths and Crustal Growth: A Conceptual Model

J. Brendan Murphy¹, William J. Collins² and Donnelly B. Archibald¹

¹Department of Earth Sciences, St. Francis Xavier University
P.O. Box 5000, Antigonish, Nova Scotia, B2G 2W5, Canada
E-mail: bmurphy@stfx.ca

²Earth Dynamics Research Group, The Institute for Geoscience Research (ITGeR), School of Earth and Planetary Sciences
Curtin University, Perth, Western Australia, Australia

J. Brendan Murphy received the 2022 GAC Logan Medal; this paper is also part of the Journal's Igneous Rock Associations Series number 29.

SUMMARY

Appinite bodies are a suite of plutonic rocks, ranging from ultramafic to felsic in composition, that are characterized by idiomorphic hornblende as the dominant mafic mineral in all lithologies and by spectacularly diverse textures, including planar and linear magmatic fabrics, mafic pegmatites and widespread evidence of mingling between coeval mafic and felsic compositions. These features suggest crystallization from anomalously water-rich magma which, according to limited isotopic studies, has both mantle and meteoric components.

Appinite bodies typically occur as small (~2 km diameter) complexes emplaced along the periphery of granitoid plutons and commonly adjacent to major deep crustal faults, which they preferentially exploit during their ascent. Several studies emphasize the relationship between intrusion of appinite, granitoid plutonism and termination of subduction. However, recent geochronological data suggest a more long-lived genetic relationship between appinite and granitoid magma generation and subduction.

Appinite may represent aliquots of hydrous basaltic magma derived from variably fractionated mafic underplates that were originally emplaced during protracted subduction adjacent to the Moho, triggering generation of voluminous granitoid magma by partial melting in the overlying MASH zone. Hydrous mafic magma from this underplate may have ascended, accumulated, and differentiated at mid-to-upper crustal levels (ca. 3–6 kbar, 15 km depth) and crystallized under water-saturated conditions. The granitoid magma was emplaced in pulses when transient stresses activated favourably oriented structures which became conduits for magma transport. The ascent of late mafic magma, however, is impeded by the rheological barriers created by the structurally overlying granitoid magma bodies. Magma that forms appinite complexes evaded those rheological barriers because it preferentially exploited the deep crustal faults that bounded the plutonic system. In this scenario, appinite complexes may be a direct connection to the mafic underplate and so its most mafic components may provide insights into processes that generate granitoid batholiths and, more generally, into crustal growth in arc systems.

RÉSUMÉ

Les corps d'appinite sont une suite de roches plutoniques, de composition ultramafique à felsique, qui se caractérisent par de la hornblende idiomorphe comme minéral mafique dominant dans toutes les lithologies et par des textures spectaculairement diverses, y compris des fabriques magmatiques planaires et linéaires, des pegmatites mafiques et de nombreuses preuves de « mingling », mélange hétérogène, des compositions mafiques et felsiques de même âge. Ces caractéristiques suggèrent une cristallisation à partir d'un magma anormalement riche en eau qui, selon un nombre limité d'études isotopiques, possède à la fois des composants mantelliques et météoriques.

Les corps d'appinite se présentent généralement sous la forme de petits complexes (~ 2 km de diamètre) mis en place à la périphérie des plutons granitoïdes et généralement adja-

cents aux principales failles crustales profondes qu'ils exploitent préférentiellement lors de leur ascension. Plusieurs études soulignent la relation entre l'intrusion d'appinite, le plutonisme granitoïde et l'arrêt de la subduction. Cependant, des données géochronologiques récentes suggèrent une relation génétique de plus longue durée entre la génération d'appinite et de magma granitoïde et la subduction.

L'appinite peut représenter des aliquotes de magma basaltique hydraté dérivées de sous-plaques mafiques à fractionnement variable qui ont été initialement mises en place lors d'une subduction prolongée adjacente au Moho, déclenchant la génération de magma granitoïde volumineux par fusion partielle dans la zone MASH sus-jacente. Le magma mafique hydraté de cette sous-plaque peut avoir remonté et s'être accumulé et différencié à des niveaux crustaux moyens à supérieurs (environ 3 à 6 kbar, 15 km de profondeur) et avoir cristallisé dans des conditions de saturation en eau. Le magma granitoïde s'est mis en place par impulsions lorsque des contraintes transitoires ont activé des structures favorablement orientées qui sont devenues des conduits pour le transport du magma. L'ascension du magma mafique tardif, cependant, est entravée par les barrières rhéologiques créées par les corps magmatiques granitoïdes structurellement sus-jacents. Le magma qui forme des complexes d'appinite a échappé à ces barrières rhéologiques car il a exploité préférentiellement les failles crustales profondes qui délimitaient le système plutonique. Dans ce scénario, les complexes d'appinite peuvent être une connexion directe à la sous-plaque mafique et ainsi ses composants les plus mafiques peuvent fournir des informations sur les processus qui génèrent des batholites granitoïdes et, plus généralement, sur la croissance crustale dans les systèmes d'arc.

Traduit par la Traductrice

INTRODUCTION

Mafic magma bodies are under-represented compared to intermediate–felsic magmas in continental arc systems, which are typically dominated by composite granitoid batholiths that are the end-product of a sequence of subduction-related processes that transfer energy and mass from the mantle to the crust (e.g. Pearce et al. 1984; Pitcher 1997; Ducea 2001). Coeval mafic rocks exposed at the same structural level as the granitoid batholiths typically occur as small plutons (1–2 km in diameter) and syn-plutonic dykes that are peri-batholithic, i.e. preferentially located along the batholith periphery (Bowes and Wright 1967; Pitcher and Berger 1972; Ratcliffe et al. 1982; Fowler and Henney 1996; Pitcher 1997; Clarke et al. 1997). However, most internal domains of batholiths preserve field evidence of interaction between coeval granitoid and mafic magmas, such as the presence of mafic enclaves (e.g. Vernon 1984; Barbarin and Didier 1992; Clarke et al. 2000; Chen et al. 2018), as well as petrographic and geochemical evidence for processes such as mingling and mixing, which can produce rocks of intermediate compositions (e.g. Chappell 1996; Tate et al. 1997; Baxter and Feely 2002; Miller et al. 2009; Muir et al. 2014). These overall relationships imply a genetic linkage

between the felsic–intermediate magmas that dominate the batholiths and coeval mafic magmas.

Appinite complexes are hornblende-rich plutonic rocks, predominantly mafic in composition, that typically occur as small bodies around the periphery of large, composite granitoid batholiths (Murphy 2013). Field relationships (e.g. Pitcher and Berger 1972) and geochronological studies (e.g. Archibald et al. 2021) indicate emplacement of appinite bodies and granitoid batholiths are broadly coeval. Their hornblende-rich mineralogy, together with their spectacularly diverse array of textures, even on a hand-specimen scale, suggests crystallization from anomalously water-rich magma (e.g. Pitcher and Berger 1972; Bowes and McArthur 1976; Pitcher and Hutton 2003; Murphy 2013, 2020). Although data are very limited, recently published O- and H-isotope data from magnesio-hornblende in an appinite body (Greendale Complex, Nova Scotia, Canada) identified a mantle component to the water incorporated in the hornblende crystal structure of mafic–ultramafic appinite (Cawood et al. 2021), suggesting a connection to the mantle processes that may have stimulated the generation of coeval granitoid magma. However, the potential importance of appinite in understanding the origin of batholiths in continental arcs (and by implication, the generation of continental crust) is often overlooked, possibly because of its subordinate volume relative to the adjacent batholith.

The tectonic setting of appinite emplacement has been largely inferred from regional studies which have emphasized the close spatial and temporal association between intrusion of appinite complexes and termination of subduction following accretional or collisional orogenesis (Atherton and Ghani 2002; Neilson et al. 2009; Granja Dorilêo Leite et al. 2021; Yuan et al. 2022). Such studies imply that appinite may be an important indicator of the tectonic processes responsible for the generation of granitoid batholiths, and by implication, the crustal growth which primarily occurs in continental arc environments (Cawood et al. 2013; Hawkesworth et al. 2013).

The purpose of this article is to highlight the potential genetic connection between appinite suite rocks and mantle-derived mafic magmas that underplate the crust and trigger the formation of granitoid magma. We also review recent geochronological evidence that the relationship between appinitic and granitoid magmas may have initiated during the subduction cycle and so may be of longer duration than hitherto realized, thereby constraining the tectonic setting of both magma generation in the lower crust and its subsequent emplacement in mid-to-shallow crustal levels.

GEOLOGICAL CONTEXT

Arc granitoid rocks commonly occur as trans-crustal composite batholiths that reflect transport of magma from the mantle and lower crust in incremental batches over timescales ranging from thousands to tens of millions of years (Petford et al. 1993, 2000; Paterson and Vernon 1995; Pitcher 1997; Glazner et al. 2004; Miller et al. 2007; Clemens and Stevens 2012; Schoene et al. 2012; Miles and Woodcock 2016; Schaltegger et al. 2019; Smith et al. 2019; Collins et al. 2020, 2021; Archibald




Figure 1. Concentration of arc magma bodies above a zone of fluid flux melting of mantle lithosphere and mafic underplating beneath the Moho (based on Hyndman et al. 2005; Weaver et al. 2011; Grove et al. 2012; Collins et al. 2020). Hydrous fluxing from the subducting slab metasomatizes the lithospheric mantle and results in the generation of water-rich mafic magmas which underplate the Moho. As they cool and fractionate, they exsolve water which initiates melting in the overlying (MASH zone) crust (diagram modified after Grove et al. 2012; Collins et al. 2020). Appinite intrusions preferentially occur along or adjacent to major deep crustal faults that bound the plutonic system. The evolution of the region above the MASH zone is expanded in Figure 4. MASH, Melting, Assimilation, Silicification, Hybridization; LAB, Lithosphere–Asthenosphere Boundary; SCLM, Subcontinental Lithospheric Mantle; TBL, Thermal Boundary Layer.

et al. 2021; Bickerton et al. 2022). A wealth of experimental, theoretical and geochemical studies indicates these processes occur during ongoing subduction by (i) melting of a lithospheric mantle wedge metasomatized by hydrous fluids and by silicic melts rising from the subducting slab and (ii) melting reactions in the subducted slab and its carapace of sedimentary

rocks at depths between 100 and 150 km (e.g. Spandler and Pirard 2013; Zhu et al. 2021).

Although the details are controversial, melting of the metasomatized mantle wedge produces hydrous to super-hydrous (> 8 wt.% H₂O) mafic magmas that rise and congregate in the vicinity of the mantle–crust boundary (Fig. 1) where they

underplate, assimilate and mix with overlying continental crust (e.g. Hildreth and Moorbath 1988; Castro 2020; Collins et al. 2020, 2021). A combination of magma, heat and fluids emanating from this mafic underplate generates a crustal regime above the underplate dominated by melting, assimilation, storage and homogenization, known as a MASH zone (Hildreth and Moorbath 1988; Annen et al. 2006). Geophysical data beneath modern arcs indicate that (i) such MASH zones typically occur at depths of ~25–30 km beneath magmatic arcs (Whitney 1988; Daczko et al. 2002; Miller et al. 2009), (ii) may be hydrous to super-hydrous (> 8 wt.% H₂O) (e.g. Bedrosian et al. 2018; Gavrilenko et al. 2019; Müntener et al. 2021) and (iii) are commonly overlain, at mid-crustal levels (ca. 6 kbar), by H₂O-rich (up to 10 wt.%) magma which stalls at these depths due to a combination of decompression-induced crystallization and fractionation, which increases magma viscosity (Lauzonier et al. 2017). Exsolution of fluids as a result of water-saturation raises the solidus temperature and so induces rapid crystallization.

MASH zones are viewed as sites where granitoid magmas originate and gestate, prior to their emplacement as composite batholiths in middle-to-upper crust (Annen et al. 2006; Jackson et al. 2018). Precise U–Pb geochronological studies indicate that gestation may be up to 20 m.y. in duration (e.g. Memeti et al. 2010; Schoene et al. 2012; Miles and Woodcock 2016; Schaltegger et al. 2019; Archibald et al. 2021; Bickerton et al. 2022). The transition from gestation to emplacement is likely triggered by a range of variables, including changes in tectonic setting (e.g. Ringwood et al. 2021) that either initiate or reactivate favourably oriented crustal structures which then become conduits for magma transport (e.g. Vigneresse 1995; Cruden 1998; Petford et al. 2000; Archibald et al. 2021).

Models to explain the generation of the felsic magmas that dominate granitoid batholiths generally fall into three end-member categories, though they are not mutually exclusive: (i) fractionation of a mafic parent (e.g. Fowler et al. 2001, 2008; Lee and Bachmann 2014; Jagoutz and Klein 2018; Müntener and Ulmer 2018; Ulmer et al. 2018; Granja Dorilêo Leite et al. 2021), (ii) relatively low temperature (< 850°C) water-fluxed partial melting of lower crust induced by heat and fluids rising from the mafic underplate (Castro 2020; Collins et al. 2020, 2021), and (iii) fluid-absent partial melting of lower crustal rocks at higher temperatures (≥ 850°C; Thompson 1982; Clemens 1998; Brown 2007).

Fractionation and water-fluxed models both require mafic magma to be more voluminous at depth than is represented at the crustal level of batholith emplacement. Exposed trans-crustal arc sections (e.g. Sierra Nevada and Woolley Creek batholiths, California, Ague 1997, Saleeby et al. 2003, Barnes et al. 2016; Fiordland, New Zealand, Daczko et al. 2002; Sierra Valle Fertil complex, Argentina, Walker et al. 2015) are characterized by lower crustal mafic rocks and ultramafic cumulates, which likely represent vestiges of the mafic underplate. In the fluid-absent model, water is provided only from the breakdown of hydrous minerals and the process is also known as dehydration melting (e.g. Thompson 1982) or hydrate-breakdown melting (Brown 2007). As recent geochronological stud-

ies have shown that many granitoid batholiths are assembled episodically over tens of millions of years (Miles and Woodcock 2018; Clemens et al. 2020; Archibald et al. 2021; Bickerton et al. 2022), these models should be considered end-members in a scenario in which water activity in the source rocks can vary over time within an evolving tectonic setting (Collins et al. 2021).

Characteristics of the Appinite Suite

Appinite suite rocks (see Murphy 2013, 2020 for details) were first defined in the Scottish Highlands (Bailey and Maufe 1916) as the plutonic equivalent of lamprophyre, with which they are commonly spatially and temporally associated. Although predominantly mafic, the appinite suite ranges from ultramafic to felsic in composition. The suite's unifying characteristics include (i) idiomorphic hornblende as the dominant mafic mineral in ultramafic to felsic rocks, and (ii) a spectacularly diverse array of textures, even on a hand-specimen scale, varying from coarse mafic pegmatite to fine grained “salt-and-pepper” hornblende gabbro and diorite (Fig. 2).

Ultramafic rocks have affinities with lamprophyre intrusions, and on IUGS classifications range from hornblendite to olivine–pyroxene hornblendite to hornblende peridotite. Volumetrically dominant mafic to intermediate rocks have a simple mineralogy (hornblende, plagioclase) and are classified as hornblende gabbro and hornblende diorite, respectively. Despite their mineralogical simplicity, geochemical analyses indicate that mafic to intermediate rocks range from high-K shoshonitic to low-K calcalkaline compositions. Intriguingly, high-K shoshonitic rocks are associated with high-K, Ba–Sr rich granite with adakitic affinities whereas granite associated with low-K calc-alkaline mafic compositions are also low-K calc-alkaline, implying a genetic relationship between coeval mafic and felsic compositions (Murphy 2020; Archibald and Murphy 2021; Archibald et al. 2022).

Taken together, these mineralogical and textural features suggest appinite bodies crystallize from anomalously water-rich mafic magma (e.g. Bowes and Wright 1967; Pitcher and Berger 1972; Pitcher and Hutton 2003) in which the stability field of hornblende is expanded relative to olivine, pyroxene and plagioclase (Moore and Carmichael 1998; Grove et al. 2003; Krawczynski et al. 2012; Loucks 2014; Fig. 3). The low viscosity of water-rich mafic magma promotes the local growth of pegmatitic textures dominated by hornblende with subordinate biotite (Fig. 2; Murphy 2013).

In addition to their peri-batholithic location, appinite bodies also preferentially occur adjacent to major fault zones, which act as conduits that facilitate their ascent to higher structural levels (Hutton 1988; Murphy and Hynes 1990; Rogers and Dunning 1991). Appinitic rocks also occur within subhorizontal mafic sheeted complexes interlayered with migmatite at the base of some plutons, which can be interpreted as active extensional detachments (Richards and Collins 2004). In some complexes, appinite bodies also exhibit locally developed planar and linear fabrics (e.g. Murphy and Hynes 1990). The subvertical planar fabrics reflect multiple cycles of magma injection and crystallization along the extensional plane of the




Figure 2. a) Example of lamprophyric sheets and pods (dark colour) enclosed by hornblende diorite (pale colour). The pods are lamprophyric sheets dismembered by later intrusions within the Greendale Complex. Inset shows hornblende poikilitically enclosing olivine and clinopyroxene in lamprophyre. The water in hornblende in the lamprophyre has a mantle water component (see Cawood et al. 2021); b) Example of textural variability of appinite suite rocks, Greendale Complex, Nova Scotia. The mineralogy is dominated by hornblende and plagioclase with minor accessory phases such as apatite, titanite, and zircon. Coarse hornblende growing perpendicular to the margins of previously injected sheets likely grew in situ (i.e. at the depth of emplacement of the Greendale Complex). The Al content of the finer grained hornblende indicates it grew at depth and was entrained in the magma as it ascended; c) Example of layering with “stacked log” hornblende growing perpendicular to the layer margins, indicating dilation during emplacement (see Murphy and Hynes 1990); d) Mafic pegmatite, dominated by idiomorphic and zoned hornblende, plagioclase, in a groundmass of hornblende, plagioclase ± quartz ± K-feldspar; e) example of mingling between coeval mafic and felsic phases; f) Example of structural control on the intrusion of felsic magma within a local “pull-apart” structure. Note the pegmatitic texture.

the depth of emplacement of the Greendale Complex). The Al content of the finer grained hornblende indicates it grew at depth and was entrained in the magma as it ascended; c) Example of layering with “stacked log” hornblende growing perpendicular to the layer margins, indicating dilation during emplacement (see Murphy and Hynes 1990); d) Mafic pegmatite, dominated by idiomorphic and zoned hornblende, plagioclase ± quartz ± K-feldspar; e) example of mingling between coeval mafic and felsic phases; f) Example of structural control on the intrusion of felsic magma within a local “pull-apart” structure. Note the pegmatitic texture.




Figure 3. Summary diagram (after Loucks 2014) showing the effect of dissolved wt.% H₂O (grey contours) on the sequence of crystallization from an andesite melt in the upper crust (composition, upper left). Each black square represents experimental results of Moore and Carmichael (1998), in which the mineral products were identified and the dissolved H₂O was determined from quenched glass. The diagram highlights the dramatic change in the sequence of crystallization with increasing dissolved H₂O, especially affecting hornblende and plagioclase, the two dominant minerals in mafic appinite.

instantaneous strain ellipsoid associated with strike-slip motion on the bounding faults during their emplacement. The linear fabrics range from orientations perpendicular to planar fabrics, consistent with extension during magma emplacement, to orientations parallel to planar fabrics, where hornblende was entrained by magma flow.

Tectonic Setting

Several studies have emphasized the close spatial and temporal association between intrusion of appinite complexes and termination of subduction (Atherton and Ghani 2002; Neilson et al. 2009; Granja Dorilêo Leite et al. 2021; Yuan et al. 2022). These interpretations are largely based on regional syntheses in which a combination of field observations and geochronological studies indicate that emplacement of appinite complexes occurred either late- or after collision-related deformation. For

example, according to Atherton and Ghani (2002), appinite emplacement occurred in the aftermath of the ca. 430–420 Ma closure of the Iapetus oceanic tract by collision between Ganderia–Avalonia and Laurentia. In this scenario, subduction of Laurentia continental crust following collision-initiated slab break-off with asthenospheric upwelling advecting heat and causing melting in the overlying mantle lithosphere which produced a mafic underplate. When the slab is detached and sinks into the mantle, rapid uplift occurs and high temperatures cause partial melting of the underplate to form granitic magma, which is emplaced in the upper crust. Similarly, appinite and coeval Ba–Sr granite of the West Kunlun orogen (northwestern margin of Tibetan Plateau; Ye et al. 2008) were emplaced during slab break-off after ca. 440 Ma closure of the Proto-Tethys Ocean (Wang et al. 2014). The advent of discrimination diagrams that identify “slab failure” granite plu-




Figure 4. Conceptual model explaining the peri-batholithic position of appinite complexes relative to granitoid batholiths and the mafic underplate in a transtensional setting (note increasing horizontal distance between faults from stages I to III). Appinite complexes are preferentially located adjacent to deep crustal faults. Stage I. Early phase of hydrous mafic magmatism includes the formation of the mafic underplate, which becomes part of the crust. Magmas are derived from metasomatized spinel- or garnet-bearing lithospheric mantle, or asthenospheric mantle. Magma derived from melting of spinel lherzolite mantle predominantly has low-K calc-alkalic affinity. Magma derived from garnet lherzolite has shoshonitic affinity. Stage II. Felsic magma masses form by fractionation of a mafic parent, by anatexis (fluid flux melting) of the lower crust and/or by melting of the mafic underplate. Felsic magma rises to the mid- and upper crust where it can become (i) neutrally buoyant, (ii) impeded by resistant lithologies, (iii) water-saturated, or (iv) trapped in extensional decollement structures that provide accommodation space for repeated injections of magma (e.g. Richards and Collins 2004). Stage III. Felsic magma bodies spread out laterally forming a rheological barrier that impedes the subsequent rise of mafic magma. Early mafic intrusions are entrained as enclaves or are hybridized. Only the appinite bodies intruding along the periphery of the system are preserved intact. The long-dashed line in Stage III shows the typical exposure level of the batholiths and appinite complexes. The short-dashed line is the Moho. Scheme modified from Hildreth and Moorbath (1988), Richard and Collins (2004) and Murphy (2020).

tons (Hildebrand and Whalen 2017; Hildebrand et al. 2018) supports models in which significant volumes of granitoid and related appinitic plutonism are late- to post-tectonic and related to slab failure (e.g. Archibald and Murphy 2021).

However, recent studies that determined the precise crystallization age of the appinite, as well as the ages of antecrysts in the coeval batholiths, suggest the association between appinite and granitoid batholiths was more long-lived, and initiated during subduction that preceded slab-breakoff. For example, recent age data show that mafic and felsic magma generation associated with the Donegal Composite Batholith and adjacent appinite intrusions (NW Ireland) overlapped for at least 15 m.y. (ca. 431–416 Ma, Murphy et al. 2019; Archibald et al. 2021). Similarly, in Avalonia of northern Nova Scotia, late Neoproterozoic emplacement of appinite and granitoid plutons overlapped for about 30 m.y. (ca. 632–602 Ma; Keppie et al. 1990; Pe-Piper et al. 1996, 2010; Murphy et al. 1997a, b; Pe-Piper and Piper 2018; White et al. 2021, 2022).

These age relationships suggest that the parental magma for appinite, like that for granite, may gestate near the base of the crust or uppermost mantle for up to 20 m.y., and that emplacement at middle to upper crustal levels occurs in specific time intervals when transient stresses act on favourably oriented structures (Archibald et al. 2021). In this context, slab failure and associated asthenospheric upwelling may provide the final impetus of energy and mass transfer from the mantle to the crust but the apparent temporal association of magmatism with subduction termination may be because the latest appinite intrusions are the best preserved.

In summary, granitoid batholiths and coeval peri-batholithic mafic intrusions are the end-product of magma bodies produced throughout subduction. If so, interpretations that relate

appinite magma genesis solely to subduction termination are an artefact of preservation potential within this dynamic and evolving magmatic system. Compositions representative of the mafic underplate that triggered granitoid magma formation may be preserved in specific regimes within appinite complexes whose emplacement spans the longevity of arc magmatism.

CONCEPTUAL MODEL FOR COEVAL APPINITE SUITE AND GRANITOID COMPLEXES

Recent high spatial resolution geochronological studies of both antecrystic and autocrystic zircon domains in granitoid batholiths have documented that (i) subduction beneath continental arcs generates granitoid magma continuously over millions of years, and (ii) batholiths are composite bodies constructed incrementally and in discrete time intervals when magma is transported from the lower crustal MASH zone and emplaced episodically in the middle or upper crust (see Glazner et al. 2004; Miles and Woodcock 2018; Archibald et al. 2021; Bickerton et al. 2022 and references therein). As emplacement is likely facilitated by transient stresses that reactivate brittle structures (Pitcher 1997; Cruden 1998), many composite batholiths are bounded by major, deeply penetrating crustal faults. These studies imply underplating of similar longevity for processes that trigger the formation of granitoid magma, i.e. the repeated intrusion into the lower crust by H₂O-rich mafic magmas (Fig. 4), an interpretation supported by field evidence and geochronological studies (e.g. Archibald et al. 2021). Recent petrological studies show that mafic arc magmas near the base of the crust in Kamchatka (1100–1050°C, 25–30 km depth) were “super-hydrous”, containing up to 14 wt.% H₂O (Goltz et al. 2020). Modeling shows that (i) this

hydrous mafic underplate solidifies as hornblende gabbro with a pyroxenitic residue and (ii) as each successive underplate cools, the emanating heat and exsolved fluids trigger fluid-fluxed crustal melting that characterizes the overlying MASH zone (Collins et al. 2020).

Felsic magma, formed by a combination of fluid-fluxed melting and fractionation of the cooling mafic underplate, eventually rises to a crustal level where it either becomes neutrally buoyant, is impeded by resistant lithologies, is water-saturated, or becomes trapped in active subhorizontal decollements whose motion provides the accommodation space for repeated injections of magma (e.g. Richards and Collins 2004; Fig. 4). At this juncture, the felsic magma migrates laterally, thereby forming a rheological barrier that impedes the subsequent rise of mafic magma above the same structural level, except along the periphery where mafic magma can exploit the crustal faults that bound the system.

Early mafic magma masses from the underplate that intrude the crust are likely to accumulate and differentiate at mid-crustal levels (ca. 3–6 kbar, 10–18 km depth). At such depths, ascending hydrous magma becomes water-saturated, which induces crystallization, and the resulting increase in viscosity inhibits further ascent (Laumonier et al. 2017). More generally, mafic magma masses that intrude early in this evolutionary history would have been engulfed by the subsequent emplacement of voluminous felsic magma, especially if such emplacement preferentially occurred where motion on active faults provide accommodation space (e.g. local extensional, transtensional or pull-apart regimes, Fig. 4). Early infusions of magma from the mafic underplate therefore have low preservation potential, occurring only as mafic xenoliths or enclaves, or possibly as one end-member of intermediate (andesitic) magmas that reflect two-stage hybridization with felsic magma at various crustal levels (e.g. Muir et al. 2014; Li et al. 2021).

Later mafic infusions into granitic magma would also have low preservation potential. Their vestiges typically occur as enclaves (e.g. pillows) and distended syn-plutonic dykes that exhibit visible evidence of magma mingling and limited evidence of mixing, resulting in rocks with intermediate compositions (Collins et al. 2000; Chen et al. 2018). As the granite solidifies beyond the particle locking threshold (~72–75% solidification, Vigneresse et al. 1996), the latest mafic intrusions may be preserved as discrete syn- to late-plutonic dykes ranging from diabase to lamprophyre in composition.

In this overall scenario, appinite bodies may represent aliquots of hydrous basaltic magma derived from a variably fractionated mafic underplate that preferentially exploited either faults located along the periphery of the magmatic system, or late-stage fractures within the essentially solidified granitic crystal mush. Their peri-batholithic location implies that the appinite magma had more limited interaction with coeval granitoid magma and so appinite complexes, preferentially emplaced along faults which define the periphery of the system, may develop as the hydrous magma decompresses and fractionates upon ascent (e.g. McCarthy and Müntener 2016). This interpretation is supported by recent $\delta^{18}\text{O}$ and δD isotopic studies of hornblende in ultramafic appinite of the Neo-

proterozoic Greendale Complex, Nova Scotia (Cawood et al. 2021), which imply growth of a generation of hornblende from magma with a significant component of mantle-derived water ($\delta^{18}\text{O}$, 4.7–6.8‰; $\delta\text{D} < -90$ ‰). This generation of hornblende preferentially occurs in lamprophyric sheets and pods (sheets dismembered by later intrusions) within the complex where they poikilitically enclose olivine and clinopyroxene.

Although a study of the aureole of the Greendale Complex indicates emplacement at 3–5 kbar (Abad et al. 2011), hornblende with mantle water isotopic signatures has an Al content (e.g. high total Al, high Al^{IV}) consistent with crystallization between 5 and 8 kbar (Murphy et al. 2012; Pe-Piper and Piper 2018; Murphy 2020; Cawood et al. 2021). More generally, experiments suggest crystallization conditions of at least 1000°C at $P > 6$ kbar for the beginning of hornblende crystallization under water-saturated conditions (Krawczynski et al. 2012). Such depths imply water contents of ~10 wt.% or greater for such mid- to lower crustal appinitic magma (Krawczynski et al. 2012). Isotopic data imply that mantle water dissolved in the magma was captured by growing hornblende which was then entrained and transported by mafic magma to shallower crustal levels by exploiting the Hollow–Greendale fault system. On the other hand, hornblende in mafic to intermediate appinite, including grains that grew in situ across the vein walls and those that exhibit exquisite porphyritic textures, have $\delta^{18}\text{O}$ (0.9 to 4.6‰) and δD (–106 to –64‰) indicating mantle water became mixed with meteoric fluids as the magma ascended. This mixing could be explained by assimilation of rocks that had previously been altered by meteoric fluids but is also compatible with recent studies (e.g. Bindeman et al. 2008; Diamond et al. 2018) of active geothermal systems that imply meteoric water can penetrate along deeply penetrating crustal faults to depths of at least 9 km where these faults are involved in seismic events. The textures indicate rapid growth of hornblende and plagioclase, and suggest the magma was likely water-saturated, implying water contents between 7 and 10 wt.% H_2O (Müntener et al. 2021).

DISCUSSION

The origin of arc-related granitic batholiths and their genetic relationship with coeval mafic magma bodies is an enduring controversy in geology. Irrespective of whether coeval mafic magmatism is primarily a source of heat and/or fluids for crustal anatexis, or is a parental magma guiding crystal fractionation, a common factor is that these models each require more voluminous mafic magma underplated at depth than is reflected by the comparatively minor volume typically exposed at the crustal level of the batholiths. Indeed, seismic reflection data beneath the Scottish Caledonides, where appinite was originally defined, are interpreted to represent invasion of the lower crust by voluminous mafic magma (Hynes and Snyder 1995). Similarly, seismic data of the crust beneath the Taupo Volcanic Zone, New Zealand, a region of voluminous rhyolitic eruptions with one of the highest heat flow zones on Earth, are consistent with voluminous mafic intrusions and/or underplated mafic crust at depths between 16 and 30 km (Har-

riation and White 2006). As such underplated material is predominantly mafic in composition, it becomes part of the crust and so represents the transfer of mass from the mantle to the crust, implying the processes involved are important in understanding mechanisms of crustal growth.

Mafic and felsic magmas produced continuously during protracted, steady-state, subduction congregate and gestate near the Moho. Although magma production is semi-continuous, magma emplacement in the middle or upper crust is episodic, and occurs during favourable changes to the stress regime within the crust (e.g. Miles and Woodcock 2018) and/or build-up of fluid pressure (Karlstrom et al. 2010). Such episodes may reflect any number of discrete tectonic events (e.g. slab roll-back, terrane accretion, oceanic plateau subduction, slab failure) that either modify the subduction zone geometry or terminate subduction. According to geodynamic models (e.g. Currie et al. 2004), the back arc region of continental arcs is anomalously hot because of small-scale asthenospheric convection driven by the reduction in mantle viscosity, which reflects the influx of water derived from the dehydrating subducting slab (Hyndman 2015). In that scenario, mafic magma can be generated in both the lithospheric and asthenospheric mantle, especially the mantle wedge, and so may exhibit a wide range in radiogenic isotopic compositions.

The dominance of felsic relative to mafic rocks at mid-crustal levels may be because felsic magma forms a rheological barrier impeding the ascent of the mafic magmatic underplate, except along the deeply penetrating faults that bound the system where dyke complexes comprised of appinite and coeval lamprophyre bodies preferentially occur (Fig. 4). As such, although their composition may be modified during their ascent, appinite complexes (especially their most mafic components) may provide a window into the composition of the mafic underplate in arc systems (Cawood et al. 2021).

Despite their mineralogical simplicity, geochemical analyses of appinite complexes indicate that mafic to intermediate rocks vary from high-K shoshonitic to low-K calc-alkaline compositions. This contrasting geochemistry likely reflects differences in the volume and composition of subduction-derived fluids and melts that contaminated the lithospheric mantle source as well as the greater depth of shoshonitic magma formation compared to low-K calc-alkaline (garnet lherzolite and spinel lherzolite mantle sources, respectively) (e.g. Peate et al. 1997; Scarrow et al. 2008; Müller and Groves 2019).

Geochemical data are consistent with a genetic connection between mafic and felsic magmas. For example, in the Scottish and Irish Caledonides, high-K shoshonitic appinite bodies are associated with Ba–Sr rich syenite and granite with adakitic affinities to the north of the Great Glen Fault, whereas appinite and coeval granite plutons have low-K calc-alkaline compositions to the south of the Great Glen Fault (Archibald et al. 2022). Although they share similar depletions in HREE and HFS elements (e.g. Ta, Nb, Ti), suggesting mafic melt generation in the garnet peridotite stability field (> 70 km depth), the high Ba–Sr granite suites contrast with low-K calc-alkalic granite suites in that they are strongly LREE enriched, and lack a

significant Eu anomaly. However, the origins of each of these suites are controversial in their own right. According to fractionation models (Fowler and Henney 1996; Fowler et al. 2001, 2008), the high Ba–Sr syenite and granitoid compositions reflect 50% fractionation of olivine, calcic clinopyroxene, biotite, apatite and titanite from a mafic parent. The mafic magmas were themselves formed by melting of a mantle wedge metasomatized by Ba–Sr rich fluids and melts derived from subducted sedimentary rocks (Fowler et al. 2008). The high Ba and Sr contents, together with the lack of a Eu anomaly in both mafic and felsic rocks, is consistent with experimental studies indicating hydrous arc magmas are characterized by a delay in plagioclase crystallization relative to olivine, clinopyroxene and hornblende (Fig. 3; Blatter et al. 2013; Loucks 2014; Nandedkar et al. 2014; Yanagida et al. 2018). Indeed, water-saturated experiments on primitive high magnesian andesite suggest plagioclase crystallizes only at 3–4 kbar and 950°C (Krawczynski et al. 2012), which explains the high Ba–Sr contents with no Eu anomaly of many lamprophyric rocks. The high LREE/HREE signatures require melting occurred at > 1 GPa in the garnet stability field (see below).

The geochemical variability of appinite also may reflect the depth and extent of partial melting as well as composition of the lithospheric mantle source of the mafic underplate. Appinite bodies with shoshonitic tendencies are highly enriched in LREE relative to HREE, reflecting low-degree melting of a deep (> 2 GPa) garnet lherzolite lithospheric mantle source. In addition, subduction-induced metasomatism produces domains in the lithospheric mantle wedge that are enriched in volatile-bearing phases, such as amphibole and phlogopite, which renders those domains more susceptible to melting (e.g. Francis and Ludden 1990; Scarrow et al. 2008; Ghent et al. 2019). Mantle domains anomalously enriched in phlogopite may in turn reflect metasomatism by K-rich fluids and felsic melts produced by dehydration and/or melting of subducted K-rich protoliths (e.g. pelite; Mallik et al. 2015). Usually, enrichment at these depths occurs because the K- and other LILE-components are retained in phengite, which destabilizes at much greater depth than amphibole (e.g. Spandler and Pirard 2012). Melting of a metasomatized garnet lherzolite mantle at depths > 70 km would produce residual garnet and yield a K- and LREE-enriched magma and, hence, a mafic underplate with a shoshonitic composition. On the other hand, experiments by Codillo et al. (2018) showed that melting of a spinel lherzolite mantle contaminated by fluid, melt or buoyant diapirs derived from a serpentinite-dominated oceanic lithosphere or subducted mélange, can generate magma of both tholeiitic and low-K calc-alkaline affinities at depths less than 70 km.

CONCLUSION

Appinite complexes may represent aliquots of hydrous magmas derived from the mafic underplate that are preferentially emplaced into deep crustal faults along the periphery of coeval granitoid complexes. If so, detailed geochemical and isotopic studies of appinite complexes, especially their most mafic components, may provide insights into the mantle and subduc-



tion dynamics, water content, and conditions of mafic melt generation responsible for continental growth and the generation of composite granitoid batholiths.

ACKNOWLEDGEMENTS

We are grateful to Michelle DeWolfe and Derek Thorkelson for constructive reviews, to editor Andy Kerr for the invitation to submit, to Rob Raeside for his diligent proof-reading, and to Randy Corney for excellent technical assistance. JBM thanks Duncan Keppie for introducing him to appinite (it seems like at least two ice ages ago), and to Duncan and Andrew Hynes for stimulating that interest. JBM and DBA are grateful to NSERC Canada for ongoing research support. WJC acknowledges funding from the James Chair of Pure and Applied Sciences at St. Francis Xavier University. In addition to Duncan and Andrew and co-authors Bill and Donnelly, JBM acknowledges the profound influence of many long-term friends and research collaborators including Damian Nance, Jarda Dostal, Rob Strachan, Cecilio Quesada, Stephen Johnston, Gabi Gutiérrez-Alonso, Ulf Linnemann, Georgia Pe-Piper, David Piper, Sergei Pisarevsky, Zheng-Xiang Li, David Evans, Richard D'Lemos, Sandra Barr, John Waldron, Peter Cawood, Cees van Staal, Shoufa Lin, Jamie Braid and generations of undergraduate and graduate students as well as post-doctoral fellows.

REFERENCES

- Abad, I., Murphy, J.B., Nieto, F., Gutiérrez-Alonso, G., and Walsh, E., 2011, Distinction between deformation and contact metamorphism by very low-grade metamorphism indicators in the Georgeville Group (Nova Scotia, Canada), *in* Aerden, D.G.A.M., and Johnson, S.E., eds., The interrelationship between deformation and metamorphism: University of Granada, Granada, Spain, p. 38–49.
- Ague, J.J., 1997, Thermodynamic calculation of emplacement pressures for batholithic rocks, California: Implication for the aluminum-in-hornblende barometer: *Geology*, v. 25, p. 563–566, [https://doi.org/10.1130/0091-7613\(1997\)025<0563:TCOEPF>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0563:TCOEPF>2.3.CO;2).
- Annen, C., Blundy, J.D., and Sparks, R.S.J., 2006, The genesis of intermediate and silicic magmas in deep crustal hot zones: *Journal of Petrology*, v. 47, p. 505–539, <https://doi.org/10.1093/ptrology/egi084>.
- Archibald, D.B., and Murphy, J.B., 2021, A slab failure origin for the Donegal composite batholith, Ireland as indicated by trace-element geochemistry, *in* Murphy, J.B., Strachan, R.A., and Quesada, C., eds., Pannotia to Pangaea: Neoproterozoic and Paleozoic Orogenic Cycles in the Circum-Atlantic Region: Geological Society, London, Special Publications, v. 503, p. 347–370, <https://doi.org/10.1144/SP503-2020-6>.
- Archibald, D.B., Macquarrie, L.M.G., Murphy, J.B., Strachan, R.A., McFarlane, C.R.M., Button, M., Larson, K.P., and Dunlop, J., 2021, The construction of the Donegal composite batholith, Irish Caledonides: Temporal constraints from U–Pb dating of zircon and titanite: *Geological Society of America Bulletin*, v. 133, p. 2335–2354, <https://doi.org/10.1130/B35856.1>.
- Archibald, D.B., Murphy, J.B., Fowler, M.B., Strachan, R.A., and Hildebrand, R.S., 2022, Testing petrogenetic models for contemporaneous mafic and felsic to intermediate magmatism within the “Newer Granite” suite of the Scottish and Irish Caledonides, *in* Kuiper, Y.D., Murphy, J.B., Nance, R.D., Strachan, R.A., and Thompson, M.D., eds., New Developments in the Appalachian-Caledonian-Variscan Orogen: Geological Society of America Special Papers, v. 554, p. 375–400, [https://doi.org/10.1130/2021.2554\(15\)](https://doi.org/10.1130/2021.2554(15)).
- Atherton, M.P., and Ghani, A.A., 2002, Slab breakoff: A model for Caledonian, late granite syn-collisional magmatism in the orthotectonic (metamorphic) zone of Scotland and Donegal, Ireland: *Lithos*, v. 62, p. 65–85, [https://doi.org/10.1016/S0024-4937\(02\)00111-1](https://doi.org/10.1016/S0024-4937(02)00111-1).
- Bailey, E.B., and Maufe, H.B., 1916, The geology of Ben Nevis and Glen Coe and the surrounding country: *Geological Society of Scotland Memoirs*, v. 53, p. 1–247.
- Barbarin, B., and Didier, J., 1992, Genesis and evolution of mafic microgranular enclaves through various types of interaction between coexisting felsic and mafic magmas: *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, v. 83, p. 145–153, <https://doi.org/10.1017/S0263593300007835>.
- Barnes, C.G., Ernst, W.G., Berry, R., and Tsujimori, T., 2016, Petrology and geochemistry of an upper crustal pluton: a view into crustal-scale magmatism during arc to retro-arc transition: *Journal of Petrology*, v. 57, p. 1361–1388, <https://doi.org/10.1093/ptrology/egw043>.
- Baxter, S., and Feely, M., 2002, Magma mixing and mingling textures in granitoids: examples from the Galway Granite, Connemara, Ireland: *Mineralogy and Petrology*, v. 76, p. 63–74, <https://doi.org/10.1007/s007100200032>.
- Bedrosian, P.A., Peacock, J.R., Bowles-Martinez, E., Schultz, A., and Hill, G.J., 2018, Crustal inheritance and a top-down control on arc magmatism at Mount St. Helens: *Nature Geoscience*, v. 11, p. 865–870, <https://doi.org/10.1038/s41561-018-0217-2>.
- Bickerton, L., Kontak, D.J., Murphy, J.B., Kellett, D.A., Samson, I.M., Marsh, J.H., Dunning, G., and Stern, R., 2022, The age and origin of the South Mountain Batholith, (Nova Scotia, Canada) as constrained by zircon U–Pb geochronology, geochemistry, and O–Hf isotopes: *Canadian Journal of Earth Sciences*, v. 59, p. 418–454, <https://doi.org/10.1139/cjes-2021-0097>.
- Bindeman, I.N., Brooks, C.K., McBirney, A.R., and Taylor, H.P., 2008, The low- $\delta^{18}\text{O}$ late-stage ferrodiorite magmas in the Skaergaard Intrusion: result of liquid immiscibility, thermal metamorphism, or meteoric water incorporation into magma?: *Journal of Geology*, v. 116, p. 571–586, <https://doi.org/10.1086/591992>.
- Blatter, D.L., Sisson, T.W., and Hankins, W.B., 2013, Crystallization of oxidized, moderately hydrous arc basalt at mid- to lower-crustal pressures: Implications for andesite genesis: *Contributions to Mineralogy and Petrology*, v. 166, p. 861–886, <https://doi.org/10.1007/s00410-013-0920-3>.
- Bowes, D.R., and McArthur, A.C., 1976, Nature and genesis of the appinitic suite: *Krystalinikum*, v. 12, p. 31–46.
- Bowes, D.R., and Wright, A.E., 1967, The explosion breccia pipes near Kentallen, Scotland, and their geological setting: *Earth and Environmental Transactions of the Royal Society of Edinburgh*, v. 67, p. 109–143, <https://doi.org/10.1017/S0080456800023954>.
- Brown, M., 2007, Crustal melting and melt extraction, ascent and emplacement in orogens: mechanisms and consequences: *Journal of the Geological Society*, v. 164, p. 709–730, <https://doi.org/10.1144/0016-76492006-171>.
- Castro, A., 2020, The dual origin of I-type granites: the contribution from experiments, *in* Janoušek, V., Bonin, B., Collins, W.J., Farina, F., and Bowden, P., eds., Post-Archean Granitic Rocks: Petrogenetic Processes and Tectonic Environments: Geological Society, London, Special Publications, v. 491, p. 101–145, <https://doi.org/10.1144/SP491-2018-110>.
- Cawood, I.P., Murphy, J.B., McCarthy, W.J., and Boyce, A.J., 2021, O and H isotopic evidence for a mantle source of water in appinite magma: An example from the late Neoproterozoic Greendale Complex, Nova Scotia: *Lithos*, v. 386–387, 105997, <https://doi.org/10.1016/j.lithos.2021.105997>.
- Cawood, P.A., Hawkesworth, C.J., and Dhuime, B., 2013, The continental record and the generation of continental crust: *Geological Society of America Bulletin*, v. 125, p. 14–32, <https://doi.org/10.1130/B30722.1>.
- Chappell, B.W., 1996, Magma mixing and the production of compositional variation within granite suites: Evidence from the granites of southeastern Australia: *Journal of Petrology*, v. 37, p. 449–470, <https://doi.org/10.1093/ptrology/37.3.449>.
- Chen, S., Fan, S., Yang, L., Zhang, Y., Zhang, L., Liu, L., and Zhang, T., 2018, The characteristics, origin, and significance of mafic microgranular enclaves in the granitoids from the Bailingshan complex, Eastern Tianshan, NW China: *Geological Journal*, v. 53, p. 87–96, <https://doi.org/10.1002/gj.3232>.
- Clarke, D.B., MacDonald, M.A., and Tate, M.C., 1997, Late Devonian mafic-felsic magmatism in the Meguma Zone, Nova Scotia, *in* Sinha, A.K., Whalen, J.B., and Hogan, J.P., eds., The Nature of Magmatism in the Appalachian Orogen: Geological Society of America Memoirs, v. 191, p. 107–127, <https://doi.org/10.1130/0-8137-1191-6.107>.
- Clarke, D.B., Fallon, R., and Heaman, L.M., 2000, Interaction among upper crustal, lower crustal, and mantle materials in the Port Mouton pluton, Meguma Lithotectonic Zone, southwest Nova Scotia: *Canadian Journal of Earth Sciences*, v. 37, p. 579–600, <https://doi.org/10.1139/e99-124>.
- Clemens, J.D., 1998, Observations on the origins and ascent mechanisms of granitic magmas: *Journal of the Geological Society*, v. 155, p. 843–851, <https://doi.org/10.1144/gsjgs.155.5.0843>.
- Clemens, J.D., and Stevens, G., 2012, What controls chemical variation in granitic magmas?: *Lithos*, v. 134–135, p. 317–329, <https://doi.org/10.1016/j.lithos.2012.01.001>.
- Clemens, J.D., Stevens, G., and Bryan, S.E., 2020, Conditions during the formation of granitic magmas by crustal melting – Hot or cold; drenched, damp or dry?: *Earth-Science Reviews*, v. 200, 102982, <https://doi.org/10.1016/j.earscirev.2019.102982>.
- Codillo, E.A., Le Roux, V., and Marschall, H.R., 2018, Arc-like magmas generated by mélange-peridotite interaction in the mantle wedge: *Nature Communications*, v. 9, 2864, <https://doi.org/10.1038/s41467-018-05313-2>.
- Collins, W.J., Richards, S.W., Healy, B.E., and Ellison, P.I., 2000, Origin of heterogeneous mafic enclaves by two-stage hybridisation in magma conduits (dykes) below and in granitic magma chambers: *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, v. 91, p. 27–45,

- <https://doi.org/10.1017/S0263593300007276>.
- Collins, W.J., Murphy, J.B., Johnson, T.E., and Huang H.-Q., 2020, Critical role of water in the formation of continental crust: *Nature Geoscience*, v. 13, p. 331–338, <https://doi.org/10.1038/s41561-020-0573-6>.
- Collins, W.J., Murphy, J.B., Blereau, E., and Huang, H.-Q., 2021, Water availability controls crustal melting temperatures: *Lithos*, v. 402–403, 106351, <https://doi.org/10.1016/j.lithos.2021.106351>.
- Cruden, A.R., 1998, On the emplacement of tabular granites: *Journal of the Geological Society*, v. 155, p. 853–862, <https://doi.org/10.1144/gsjgs.155.5.0853>.
- Currie, C.A., Wang, K., Hyndman, R.D., and He, J., 2004, The thermal effects of steady-state slab-driven mantle flow above a subducting plate: The Cascadia subduction zone and backarc: *Earth and Planetary Science Letters*, v. 223, p. 35–48, <https://doi.org/10.1016/j.epsl.2004.04.020>.
- Daczko, N.R., Clarke, G.L., and Klepeis, K.A., 2002, Kyanite-paragonite-bearing assemblages, northern Fiordland, New Zealand: rapid cooling at the lower crustal root to a Cretaceous magmatic arc: *Journal of Metamorphic Geology*, v. 20, p. 887–902, <https://doi.org/10.1046/j.1525-1314.2002.00421.x>.
- Diamond, L.W., Wanner, C., and Waber, H.N., 2018, Penetration depth of meteoric water in orogenic geothermal systems: *Geology*, v. 46, p. 1063–1066, <https://doi.org/10.1130/G45394.1>.
- Ducea, M., 2001, The California arc: thick granitic batholiths, eclogitic residues, lithospheric-scale thrusting, and magmatic flare-ups: *GSA Today*, v. 11, p. 4–10, [https://doi.org/10.1130/1052-5173\(2001\)011<0004:TCATGB>2.0.CO;2](https://doi.org/10.1130/1052-5173(2001)011<0004:TCATGB>2.0.CO;2).
- Fowler, M.B., 1988, Ach'uaie hybrid apinitic pipes: Evidence for mantle-derived shoshonitic parent magmas in Caledonian granite genesis: *Geology*, v. 16, p. 1026–1030, [https://doi.org/10.1130/0091-7613\(1988\)016<1026:AUHAPE>2.3.CO;2](https://doi.org/10.1130/0091-7613(1988)016<1026:AUHAPE>2.3.CO;2).
- Fowler, M.B., and Henney, P.J., 1996, Mixed Caledonian apinitic magmas: implications for lamprophyre fractionation and high Ba-Sr granite genesis: *Contributions to Mineralogy and Petrology*, v. 126, p. 199–215, <https://doi.org/10.1007/s004100050244>.
- Fowler, M.B., Henney, P.J., Darbyshire, D.P.F., and Greenwood, P.B., 2001, Petrogenesis of high Ba-Sr granites: the Rogart pluton, Sutherland: *Journal of the Geological Society*, v. 158, p. 521–534, <https://doi.org/10.1144/jgs.158.3.521>.
- Fowler, M.B., Kocks, H., Darbyshire, D.P.F., and Greenwood, P.B., 2008, Petrogenesis of high Ba-Sr plutons from the Northern Highlands Terrane of the British Caledonian Province: *Lithos*, v. 105, p. 129–148, <https://doi.org/10.1016/j.lithos.2008.03.003>.
- Francis, D., and Ludden, J., 1990, The mantle source for olivine nephelinite, basanite, and alkaline olivine basalt at Fort Selkirk, Yukon, Canada: *Journal of Petrology*, v. 31, p. 371–400, <https://doi.org/10.1093/petrology/31.2.371>.
- Gavrilenko, M., Krawczynski, M., Ruprecht, P., Li, W., and Catalano, J.G., 2019, The quench control of water estimates in convergent margin magmas: *American Mineralogist*, v. 104, p. 936–948, <https://doi.org/10.2138/am-2019-6735>.
- Ghent, E.D., Edwards, B.R., and Russell, J.K., 2019, Pargasite-bearing vein in spinel lherzolite from the mantle lithosphere of the North America Cordillera: *Canadian Journal of Earth Sciences*, v. 56, p. 870–885, <https://doi.org/10.1139/cjes-2018-0239>.
- Glazner, A.F., Bartley, J.M., Coleman, D.S., Gray, W., and Taylor, R.Z., 2004, Are plutons assembled over millions of years by amalgamation from small magma chambers?: *GSA Today*, v. 14, p. 4–11, [https://doi.org/10.1130/1052-5173\(2004\)014<0004:APAOMO>2.0.CO;2](https://doi.org/10.1130/1052-5173(2004)014<0004:APAOMO>2.0.CO;2).
- Goltz, A.E., Krawczynski, M.J., Gavrilenko, M., Gorbach, N.V., and Ruprecht, P., 2020, Evidence for superhydrous primitive arc magmas from mafic enclaves at Shiveluch volcano, Kamchatka: *Contributions to Mineralogy and Petrology*, v. 175, 115, <https://doi.org/10.1007/s00410-020-01746-5>.
- Granja Doriléo Leite, A.F., Fuck, R.A., Dantas, E.L., and Ruiz, A.S., 2021, Appinitic and high Ba-Sr magmatism in central Brazil: Insights into the late accretion stage of West Gondwana: *Lithos*, v. 398–399, 106333, <https://doi.org/10.1016/j.lithos.2021.106333>.
- Grove, T.L., Elkins-Tanton, L.T., Parman, S.W., Chatterjee, N., Müntener, O., and Gaetani, G.A., 2003, Fractional crystallization and mantle-melting controls on calc-alkaline differentiation trends: *Contributions to Mineralogy and Petrology*, v. 145, p. 515–533, <https://doi.org/10.1007/s00410-003-0448-z>.
- Grove, T.L., Till, C.B., and Krawczynski, M.J., 2012, The role of H₂O in subduction zone magmatism: *Annual Review of Earth and Planetary Sciences*, v. 40, p. 413–439, <https://doi.org/10.1146/annurev-earth-042711-105310>.
- Harrison, A., and White, R.S., 2006, Lithospheric structure of an active backarc basin: the Taupo Volcanic Zone, New Zealand: *Geophysical Journal International*, v. 167, p. 968–990, <https://doi.org/10.1111/j.1365-246X.2006.03166.x>.
- Hawkesworth, C., Cawood, P., and Dhuime, B., 2013, Continental growth and the crustal record: *Tectonophysics*, v. 609, p. 651–660, <https://doi.org/10.1016/j.tecto.2013.08.013>.
- Hildebrand, R.S., and Whalen, J.B., 2017, The tectonic setting and origin of Cretaceous batholiths within the North American Cordillera: The case for slab failure magmatism and its significance for crustal growth: *Geological Society of America Special Papers*, v. 532, 113 p., <https://doi.org/10.1130/2017.2532>.
- Hildebrand, R.S., Whalen, J.B., and Bowring, S.A., 2018, Resolving the crustal composition paradox by 3.8 billion years of slab failure magmatism and collisional recycling of continental crust: *Tectonophysics*, v. 734–735, p. 69–88, <https://doi.org/10.1016/j.tecto.2018.04.001>.
- Hildreth, W., and Moorhead, S., 1988, Crustal contributions to arc magmatism in the Andes of Central Chile: *Contributions to Mineralogy and Petrology*, v. 98, p. 455–489, <https://doi.org/10.1007/BF00372365>.
- Hutton, D.H.W., 1988, Igneous emplacement in a shear-zone termination: The biotite granite at Strontian, Scotland: *Geological Society of America Bulletin*, v. 100, p. 1392–1399, [https://doi.org/10.1130/0016-7606\(1988\)100<1392:IEIASZ>2.3.CO;2](https://doi.org/10.1130/0016-7606(1988)100<1392:IEIASZ>2.3.CO;2).
- Hyndman, R.D., 2015, Tectonic consequences of a uniformly hot backarc and why is the Cordilleran mountain belt high?: *Geoscience Canada*, v. 42, p. 383–402, <https://doi.org/10.12789/geocanj.2015.42.078>.
- Hyndman, R.D., Currie, C.A., and Mazzotti, S.P., 2005, Subduction zone backarcs, mobile belts, and orogenic heat: *GSA Today*, v. 15, p. 4–10, [https://doi.org/10.1130/1052-5173\(2005\)15<4:SZBMBA>2.0.CO;2](https://doi.org/10.1130/1052-5173(2005)15<4:SZBMBA>2.0.CO;2).
- Hynes, A., and Snyder, D.B., 1995, Deep-crustal mineral assemblages and potential for crustal rocks below the Moho in the Scottish Caledonides: *Geophysical Journal International*, v. 123, p. 323–339, <https://doi.org/10.1111/j.1365-246X.1995.tb06857.x>.
- Jackson, M.D., Blundy, J., and Sparks, R.S.J., 2018, Chemical differentiation, cold storage and remobilization of magma in the Earth's crust: *Nature*, v. 564, p. 405–409, <https://doi.org/10.1038/s41586-018-0746-2>.
- Jagoutz, O., and Klein, B., 2018, On the importance of crystallization-differentiation for the generation of SiO₂-rich melts and the compositional build-up of arc (and continental) crust: *American Journal of Science*, v. 318, p. 29–63, <https://doi.org/10.2475/01.2018.03>.
- Karlstrom, L., Dufek, J., and Manga, M., 2010, Magma chamber stability in arc and continental crust: *Journal of Volcanology and Geothermal Research*, v. 190, p. 249–270, <https://doi.org/10.1016/j.jvolgeores.2009.10.003>.
- Keppie, J.D., Dallmeyer, R.D., and Murphy, J.B., 1990, Tectonic implications of ⁴⁰Ar/³⁹Ar hornblende ages from late Proterozoic–Cambrian plutons in the Avalon Composite Terrane, Nova Scotia, Canada: *Geological Society of America Bulletin*, v. 102, p. 516–528, [https://doi.org/10.1130/0016-7606\(1990\)102<0516:TIOAAH>2.3.CO;2](https://doi.org/10.1130/0016-7606(1990)102<0516:TIOAAH>2.3.CO;2).
- Krawczynski, M.J., Grove, T.L., and Behrens, H., 2012, Amphibole stability in primitive arc magmas: effects of temperature, H₂O content, and oxygen fugacity: *Contributions to Mineralogy and Petrology*, v. 164, p. 317–339, <https://doi.org/10.1007/s00410-012-0740-x>.
- Laumonier, M., Gaillard, F., Muir, D., Blundy, J., and Unsworth, M., 2017, Giant magmatic water reservoirs at mid-crustal depth inferred from electrical conductivity and the growth of the continental crust: *Earth and Planetary Science Letters*, v. 457, p. 173–180, <https://doi.org/10.1016/j.epsl.2016.10.023>.
- Lee, C.-T.A., and Bachmann, O., 2014, How important is the role of crystal fractionation in making intermediate magmas? Insights from Zr and P systematics: *Earth and Planetary Science Letters*, v. 393, p. 266–274, <https://doi.org/10.1016/j.epsl.2014.02.044>.
- Li, R., Collins, W.J., Yang, J.H., Blereau, E., and Wang, H., 2021, Two-stage hybrid origin of Lachlan S-type magmas: A re-appraisal using isotopic microanalysis of lithic inclusion minerals: *Lithos*, v. 402–403, 106378, <https://doi.org/10.1016/j.lithos.2021.106378>.
- Loucks, R.R., 2014, Distinctive composition of copper-ore-forming arc magmas: *Australian Journal of Earth Sciences*, v. 61, p. 5–16, <https://doi.org/10.1080/08120099.2013.865676>.
- Mallik, A., Nelson, J., and Dasgupta, R., 2015, Partial melting of fertile peridotite fluxed by hydrous rhyolitic melt at 2–3 GPa: implications for mantle wedge hybridization by sediment melt and generation of ultrapotassic magmas in convergent margins: *Contributions to Mineralogy and Petrology*, v. 169, 48, <https://doi.org/10.1007/s00410-015-1139-2>.
- Mallik, A., Dasgupta, R., Tsuno, K., and Nelson, J., 2016, Effects of water, depth and temperature on partial melting of mantle-wedge fluxed by hydrous sediment-melt in subduction zones: *Geochimica et Cosmochimica Acta*, v. 195, p. 226–243, <https://doi.org/10.1016/j.gca.2016.08.018>.
- McCarthy, A., and Müntener, O., 2016, Comb layering monitors decompressing and fractionating hydrous mafic magmas in subvolcanic plumbing systems (Fisher Lake, Sierra Nevada, USA): *Journal of Geophysical Research: Solid Earth*, v. 121, p. 8595–8621, <https://doi.org/10.1002/2016JB013489>.
- Memeti, V., Paterson, S., Matzel, J., Mundil, R., and Okaya, D., 2010, Magmatic lobes



- as “snapshots” of magma chamber growth and evolution in large, composite batholiths: an example from the Tuolumne intrusion, Sierra Nevada, California: *Geological Society of America Bulletin*, v. 122, p. 1912–1931, <https://doi.org/10.1130/B30004.1>.
- Miles, A.J., and Woodcock, N.H., 2018, A combined geochronological approach to investigating long lived granite magmatism, the Shap granite, UK: *Lithos*, v. 304–307, p. 245–257, <https://doi.org/10.1016/j.lithos.2018.02.012>.
- Miles, A.J., Woodcock, N.H., and Hawkesworth, C.J., 2016, Tectonic controls on post-subduction granite genesis and emplacement: The late Caledonian suite of Britain and Ireland: *Gondwana Research*, v. 39, p. 250–260, <https://doi.org/10.1016/j.gr.2016.02.006>.
- Miller, J.S., Matzel, J.E.P., Miller, C.F., Burgess, S.D., and Miller, R.B., 2007, Zircon growth and recycling during the assembly of large, composite arc plutons: *Journal of Volcanology and Geothermal Research*, v. 167, p. 282–299, <https://doi.org/10.1016/j.jvolgeores.2007.04.019>.
- Miller, R.B., Paterson, S.R., and Matzel, J.P., 2009, Plutonism at different crustal levels: Insights from the ~5–40 km (paleodepth) North Cascades crustal section, Washington, in Miller, R.B., and Snoke, A.W., eds., *Crustal Cross Sections from the Western North American Cordillera and Elsewhere: Implications for Tectonic and Petrologic Processes*: Geological Society of America Special Papers, v. 456, p. 1–25, [https://doi.org/10.1130/2009.2456\(05\)](https://doi.org/10.1130/2009.2456(05)).
- Moore, G., and Carmichael, I.S.E., 1998, The hydrous phase equilibria (to 3 kbar) of an andesite and basaltic andesite from western Mexico: Constraints on water content and conditions of phenocryst growth: *Contributions to Mineralogy and Petrology*, v. 130, p. 304–319, <https://doi.org/10.1007/s004100050367>.
- Muir, D.D., Blundy, J.D., Rust, A.C., and Hickey, J., 2014, Experimental constraints on dacite pre-eruptive magma storage conditions beneath Uturuncu Volcano: *Journal of Petrology*, v. 55, p. 749–767, <https://doi.org/10.1093/petrology/egu005>.
- Müller, D., and Groves, D.I., 2019, *Potassic igneous rocks and associated gold-copper mineralization* (5th edition): *Mineral Resource Reviews*, Springer Cham, 398 p., <https://doi.org/10.1007/978-3-319-92979-8>.
- Müntener, O., and Ulmer, P., 2006, Experimentally derived high-pressure cumulates from hydrous arc magmas and consequences for the seismic velocity structure of lower arc crust: *Geophysical Research Letters*, v. 33, L21308, <https://doi.org/10.1029/2006GL027629>.
- Müntener, O., Ulmer, P., and Blundy, J.D., 2021, Superhydrous arc magmas in the Alpine context: *Elements*, v. 17, p. 35–40, <https://doi.org/10.2138/gselements.17.1.35>.
- Murphy, J.B., 2013, Appinite suites: A record of the role of water in the genesis, transport, emplacement and crystallization of magma: *Earth-Science Reviews*, v. 119, p. 35–59, <https://doi.org/10.1016/j.earscirev.2013.02.002>.
- Murphy, J.B., 2020, Appinite suites and their genetic relationship with coeval voluminous granitoid batholiths: *International Geology Review*, v. 62, p. 683–713, <https://doi.org/10.1080/00206814.2019.1630859>.
- Murphy, J.B., and Hynes, A.J., 1990, Tectonic control on the origin and orientation of igneous layering: an example from the Greendale Complex, Antigonish Highlands, Nova Scotia, Canada: *Geology*, v. 18, p. 403–406, [https://doi.org/10.1130/0091-7613\(1990\)0182.3.CO;2](https://doi.org/10.1130/0091-7613(1990)0182.3.CO;2).
- Murphy, J.B., Hynes, A.J., and Cousens, B.L., 1997a, Tectonic influence on late Proterozoic Avalonian magmatism: An example from the Greendale Complex, Antigonish Highlands, Nova Scotia, Canada, in Sinha, A.K., Whalen, J.B., and Hogan, J.P., eds., *The Nature of Magmatism in the Appalachian Orogen*: Geological Society of America Memoirs, v. 191, p. 255–274, <https://doi.org/10.1130/0-8137-1191-6.255>.
- Murphy, J.B., Keppie, J.D., Davis, D., and Krogh, T.E., 1997b, Regional significance of new U–Pb age data for Neoproterozoic igneous units in Avalonian rocks of northern mainland Nova Scotia, Canada: *Geological Magazine*, v. 134, p. 113–120, <https://doi.org/10.1017/S0016756897006596>.
- Murphy, J.B., Blais, S.A., Tubrett, M., McNeil, D., and Middleton, M., 2012, Microchemistry of amphiboles near the roof of a mafic magma chamber: Insights into high level melt evolution: *Lithos*, v. 148, p. 162–175, <https://doi.org/10.1016/j.lithos.2012.06.012>.
- Murphy, J.B., Nance, R.D., Gabler, L.B., Martell, A., and Archibald, D.A., 2019, Age, geochemistry and origin of the Ardara appinite plutons, Northwest Donegal, Ireland: *Geoscience Canada*, v. 46, p. 31–48, <https://doi.org/10.12789/geocanj.2019.46.144>.
- Nandedkar, R.H., Ulmer, P., and Müntener, O., 2014, Fractional crystallization of primitive, hydrous arc magmas: An experimental study at 0.7 GPa: *Contributions to Mineralogy and Petrology*, v. 167, 1015, <https://doi.org/10.1007/s00410-014-1015-5>.
- Neilson, J.C., Kokelaar, B.P., and Crowley, Q.G., 2009, Timing, relations and cause of plutonic and volcanic activity of the Siluro–Devonian post-collision magmatic episode in the Grampian Terrane, Scotland: *Journal of the Geological Society*, v. 166, p. 545–561, <https://doi.org/10.1144/0016-76492008-069>.
- Paterson, S.R., and Vernon, R.H., 1995, Bursting the bubble of ballooning plutons: A return to nested diapirs emplaced by multiple processes: *Geological Society of America Bulletin*, v. 107, p. 1356–1380, [https://doi.org/10.1130/0016-7606\(1995\)107<1356:BTBOBP>2.3.CO;2](https://doi.org/10.1130/0016-7606(1995)107<1356:BTBOBP>2.3.CO;2).
- Pe-Piper, G., and Piper, D.J.W., 2018, The Jeffers Brook diorite–granodiorite pluton: Style of emplacement and role of volatiles at various crustal levels in Avalonian appinites, Canadian Appalachians: *International Journal of Earth Sciences*, v. 107, p. 863–883, <https://doi.org/10.1007/s00531-017-1536-z>.
- Pe-Piper, G., Piper, D.J.W., and Koukouvelas, I., 1996, Precambrian plutons of the Cobequid Highlands, Nova Scotia, Canada, in Nance, R.D., and Thompson, M.D., eds., *Avalonian and Related Peri-Gondwana Terranes of the Circum-North Atlantic*: Geological Society of America Special Papers, v. 304, p. 121–132, <https://doi.org/10.1130/0-8137-2304-3.121>.
- Pe-Piper, G., Piper, D.J.W., and Tsikouras, B., 2010, The late Neoproterozoic Frog Lake hornblende gabbro pluton, Avalon Terrane of Nova Scotia: evidence for the origins of appinites: *Canadian Journal of Earth Sciences*, v. 47, p. 103–120, <https://doi.org/10.1139/E09-077>.
- Pearce, J.A., Harris, N.B.W., and Tindle, A.G., 1984, Trace element discrimination diagrams for the tectonic interpretation of granitic rocks: *Journal of Petrology*, v. 25, p. 956–983, <https://doi.org/10.1093/petrology/25.4.956>.
- Peate, D.W., Pearce, J.A., Hawkesworth, C.J., Colley, H., Edwards, C.M.H., and Hirose, K., 1997, Geochemical variations in Vanuatu arc lavas: The role of subducted material and a variable mantle wedge composition: *Journal of Petrology*, v. 38, p. 1331–1358, <https://doi.org/10.1093/petroj/38.10.1331>.
- Petford, N., Kerr, R.C., and Lister, J.R., 1993, Dike transport of granitoid magmas: *Geology*, v. 21, p. 845–848, [https://doi.org/10.1130/0091-7613\(1993\)021<0845:DTOGM>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0845:DTOGM>2.3.CO;2).
- Petford, N., Cruden, A.R., McCaffrey, K.J.W., and Vigneresse, J.-L., 2000, Granite magma formation, transport and emplacement in the Earth’s crust: *Nature*, v. 408, p. 669–673, <https://doi.org/10.1038/35047000>.
- Pitcher, W.S., 1997, *The Nature and Origin of Granite*, (2nd edition): Springer Dordrecht, 395 p., <https://doi.org/10.1007/978-94-011-5832-9>.
- Pitcher, W.S., and Berger, A.R., 1972, The appinite suite: Basic rocks genetically associated with granite, in Pitcher, W.S., and Berger, A.R., *The Geology of Donegal, A Study of Granite Emplacement and Unroofing*: John Wiley and Sons Ltd, Chichester, Sussex, p. 143–168.
- Pitcher, W.S., and Hutcheon, D.H.W., 2003, *A Master Class Guide to the Granites of Donegal: Geological Survey of Ireland*, Dublin, 97 p.
- Ratliff, N.M., Armstrong, R.L., Mose, D.G., Seneschal, R., Williams, N., and Baia-monte, M.J., 1982, Emplacement history and tectonic significance of the Cortlandt complex, related plutons, and dike swarms in the Taconide Zone of southeastern New York based on K–Ar and Rb–Sr investigations: *American Journal of Science*, v. 282, p. 358–390, <https://doi.org/10.2475/ajs.282.3.358>.
- Richards, S.W., and Collins, W.J., 2004, Growth of wedge-shaped plutons at the base of active half grabens: *Royal Society of Edinburgh Transactions, Earth Sciences*, v. 95, p. 309–317, <https://doi.org/10.1017/S0263593304000252>.
- Ringwood, M.F., Schwartz, J.J., Turnbull, R.E., and Tulloch, A.J., 2021, Phanerozoic record of mantle-dominated arc magmatic surges in the Zealandia Cordillera: *Geology*, v. 49, p. 1230–1234, <https://doi.org/10.1130/G48916.1>.
- Rogers, G., and Dunning, G.R., 1991, Geochronology of appinitic and related granitic magmatism in the W Highlands of Scotland: Constraints on the timing of transcurrent fault movement: *Journal of the Geological Society*, v. 148, p. 17–27, <https://doi.org/10.1144/gsjgs.148.1.0017>.
- Saleeby, J., Ducea, M., and Clemens-Knott, D., 2003, Production and loss of high-density batholithic root, southern Sierra Nevada, California: *Tectonics*, v. 22, 1064, <https://doi.org/10.1029/2002TC001374>.
- Scarrow, J.H., Bea, F., Montero, P., and Molina, J.F., 2008, Shoshonites, vaugnerites and potassic lamprophyres: Similarities and differences between ‘ultra’-high-K rocks: *Earth and Environmental Sciences Transactions of the Royal Society of Edinburgh*, v. 99, p. 159–175, <https://doi.org/10.1017/S1755691009008032>.
- Schaltegger, U., Nowak, A., Ulianov, A., Fisher, C.M., Gerdes, A., Spikings, R., Whitehouse, M.J., Bindeman, I., Hanchar, J.M., Duff, J., Vervoort, J.D., Shel-drake, T., Caricchi, L., Brack, P., and Müntener, O., 2019, Zircon petrochronology and ⁴⁰Ar/³⁹Ar thermochronology of the Adamello Intrusive Suite, N. Italy: Monitoring the growth and decay of an incrementally assembled magmatic system: *Journal of Petrology*, v. 60, p. 701–722, <https://doi.org/10.1093/petrology/egz010>.
- Schoene, B., Schaltegger, U., Brack, P., Latkoczy, C., Stracke, A., and Günther, D., 2012, Rates of magma differentiation and emplacement in a ballooning pluton recorded by U–Pb TIMS-TEA, Adamello batholith, Italy: *Earth and Planetary Science Letters*, v. 355–356, p. 162–173, <https://doi.org/10.1016/>

- j.epsl.2012.08.019.
- Smith, W.D., Darling, J.R., Bullen, D.S., Lasalle, S., Pereira, I., Moreira, H., Allen, C.J., and Tapster, S., 2019, Zircon perspectives on the age and origin of evolved S-type granites from the Cornubian Batholith, southwest England: *Lithos*, v. 336–337, p. 14–26, <https://doi.org/10.1016/j.lithos.2019.03.025>.
- Spandler, C., and Pirard, C., 2013, Element recycling from subducting slabs to arc crust: a review: *Lithos*, v. 170–171, p. 208–223, <https://doi.org/10.1016/j.lithos.2013.02.016>.
- Tate, M.C., Clarke, D.B., and Heaman, L.M., 1997, Progressive hybridisation between Late Devonian mafic-intermediate and felsic magmas in the Meguma Zone of Nova Scotia, Canada: *Contributions to Mineralogy and Petrology*, v. 126, p. 401–415, <https://doi.org/10.1007/s004100050259>.
- Thompson, A.B., 1982, Dehydration melting of pelitic rocks and the generation of H₂O-undersaturated granitic liquids: *American Journal of Science*, v. 282, p. 1567–1595, <https://doi.org/10.2475/ajs.282.10.1567>.
- Ulmer, P., Kaegi, R., and Müntener, O., 2018, Experimentally derived intermediate to silica-rich arc magmas by fractional and equilibrium crystallization at 1•0 GPa: an evaluation of phase relationships, compositions, liquid lines of descent and oxygen fugacity: *Journal of Petrology*, v. 59, p. 11–58, <https://doi.org/10.1093/ptrology/egy017>.
- Vernon, R.H., 1984, Microgranitoid enclaves in granites—globules of hybrid magma quenched in a plutonic environment: *Nature*, v. 309, p. 438–439, <https://doi.org/10.1038/309438a0>.
- Vigneresse, J.L., 1995, Control of granite emplacement by regional deformation: *Tectonophysics*, v. 249, p. 173–186, [https://doi.org/10.1016/0040-1951\(95\)00004-7](https://doi.org/10.1016/0040-1951(95)00004-7).
- Vigneresse, J.L., Barbey, P., and Cuney, M., 1996, Rheological transitions during partial melting and crystallization with application to felsic magma segregation and transfer: *Journal of Petrology*, v. 37, p. 1579–1600, <https://doi.org/10.1093/ptrology/37.6.1579>.
- Walker Jr., B.A., Bergantz, G.W., Otamendi, J.E., Ducea, M.N., and Cristofolini, E.A., 2015, A MASH zone revealed: The mafic complex of the Sierra Valle Fértil: *Journal of Petrology*, v. 56, p. 1863–1896, <https://doi.org/10.1093/ptrology/egv057>.
- Wang, C., Liu, L., Xiao, P.-X., Cao, Y.-T., Yu, H.-Y., Meert, J.G., and Liang, W.-T., 2014, Geochemical and geochronologic constraints for Paleozoic magmatism related to the orogenic collapse in the Qimantagh–South Altyn region, northwestern China: *Lithos*, v. 202–203, p. 1–20, <https://doi.org/10.1016/j.lithos.2014.05.016>.
- Weaver, S.L., Wallace, P.J., and Johnston, A.D., 2011, A comparative study of continental vs. intraoceanic arc mantle melting: Experimentally determined phase relations of hydrous primitive melts: *Earth and Planetary Science Letters*, v. 308, p. 97–106, <https://doi.org/10.1016/j.epsl.2011.05.040>.
- White, C.E., Barr, S.M., Hamilton, M.A., and Murphy, J.B., 2021, Age and tectonic setting of Neoproterozoic granitoid rocks, Antigonish Highlands, Nova Scotia, Canada: Implications for Avalonia in the northern Appalachian orogen: *Canadian Journal of Earth Sciences*, v. 58, p. 396–412, <https://doi.org/10.1139/cjes-2020-0110>.
- White, C.E., Barr, S.M., Crowley, J.L., van Rooyen, D., and MacHattie, T.G., 2022, U–Pb zircon ages and Sm–Nd isotopic data from the Cobequid Highlands, Nova Scotia, Canada: New contributions to understanding the Neoproterozoic geologic history of Avalonia, in Kuiper, Y.D., Murphy, J.B., Nance, R.D., Strachan, R.A., and Thompson, M.D., eds., *New Developments in the Appalachian-Caledonian-Variscan Orogen*: Geological Society of America Special Papers, v. 554, p. 135–172, [https://doi.org/10.1130/2021.2554\(07\)](https://doi.org/10.1130/2021.2554(07)).
- Whitney, J.A., 1988, The origin of granite: The role and source of water in the evolution of granitic magmas: *Geological Society of America Bulletin*, v. 100, p. 1886–1897, [https://doi.org/10.1130/0016-7606\(1988\)100<1886:TOOGTR>2.3.CO;2](https://doi.org/10.1130/0016-7606(1988)100<1886:TOOGTR>2.3.CO;2).
- Yanagida, Y., Nakamura, M., Yasuda, A., Kuritani, T., Nakagawa, M., and Yoshida, T., 2018, Differentiation of a hydrous arc magma recorded in melt inclusions in deep crustal cumulate xenoliths from Ichinomegata Maar, NE Japan: *Geochemistry, Geophysics, Geosystems*, v. 19, p. 838–864, <https://doi.org/10.1002/2017GC007301>.
- Ye, H.-M., Li, X.-H., Li, Z.-X., and Zhang, C.-L., 2008, Age and origin of high Ba–Sr appinitic–granites at the northwestern margin of the Tibet Plateau: Implications for early Paleozoic tectonic evolution of the Western Kunlun orogenic belt: *Gondwana Research*, v. 13, p. 126–138, <https://doi.org/10.1016/j.gr.2007.08.005>.
- Yuan, L., Zhang, X., and Yang, Z., 2022, The timeline of prolonged accretionary processes in eastern Central Asian Orogenic Belt: Insights from episodic Paleozoic intrusions in central Inner Mongolia, North China: *Geological Society of America Bulletin*, v. 134, p. 629–657, <https://doi.org/10.1130/B35907.1>.
- Zhu, Z., Ding, Y., Li, Z., Dong, Y., Wang, H., Liu, J., Zhu, J., Li, X., Chu, F., and Jin, X., 2021, Hafnium isotopic constraints on crustal assimilation in response to the tectono–magmatic evolution of the Okinawa Trough: *Lithos*, v. 398–399, 106352, <https://doi.org/10.1016/j.lithos.2021.106352>.

Received June 2022

Accepted as revised September 2022

