

# Geochemistry of Carboniferous peralkaline felsic volcanic rocks, central New Brunswick, Canada: examination of uranium potential

Taryn R. Gray, Jaroslav Dostal, Malcolm McLeod, Duncan Keppie and Yuanyuan Zhang

Volume 46, 2010

URI: <https://id.erudit.org/iderudit/ageo46art10>

[See table of contents](#)

Publisher(s)

Atlantic Geoscience Society

ISSN

0843-5561 (print)

1718-7885 (digital)

[Explore this journal](#)

## Article abstract

Felsic volcanic rocks from the ca. 335 Ma Cumberland Hill Formation of southern New Brunswick consist of peralkaline rhyolite and trachyte. These rocks are interpreted to have undergone extensive fractional crystallization from an alkali basaltic magma, although their different geochemical signatures indicate that the rhyolite was not derived from the trachyte via continuous fractional crystallization. The rhyolite is highly enriched in incompatible trace elements including uranium (up to ca. 20 ppm). Cumberland Hill peralkaline rhyolite suggests that a potentially economic source of uranium exists in the basin, based on the elevated uranium concentrations and high potential for remobilization into surrounding sedimentary units.

## Cite this article

Gray, T. R., Dostal, J., McLeod, M., Keppie, D. & Zhang, Y. (2010). Geochemistry of Carboniferous peralkaline felsic volcanic rocks, central New Brunswick, Canada: examination of uranium potential. *Atlantic Geology*, 46, 173–184.

# Geochemistry of Carboniferous peralkaline felsic volcanic rocks, central New Brunswick, Canada: examination of uranium potential

TARYN R. GRAY<sup>1\*</sup>, JAROSLAV DOSTAL<sup>1</sup>, MALCOLM McLEOD<sup>2</sup>, DUNCAN KEPPIE<sup>3</sup>, YUANYUAN ZHANG<sup>4</sup>

1. Department of Geology, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada

2. New Brunswick Department of Natural Resources and Energy, Geological Surveys Branch,  
P.O. Box 5040, Sussex, New Brunswick E4E 5L2, Canada

3. Instituto de Geología, Universidad Nacional Autónoma de México, 04510, México

4. Key Laboratory of Orogenic Belts and Crustal Evolution, Ministry of Education,  
School of Earth and Space Sciences, Peking University, Beijing, China 100871

\*Corresponding author <Taryn.Gray@smu.ca>

*Date received: 30 April 2010* ¶ *Date accepted: 16 September 2010*

## ABSTRACT

Felsic volcanic rocks from the ca. 335 Ma Cumberland Hill Formation of southern New Brunswick consist of peralkaline rhyolite and trachyte. These rocks are interpreted to have undergone extensive fractional crystallization from an alkali basaltic magma, although their different geochemical signatures indicate that the rhyolite was not derived from the trachyte via continuous fractional crystallization. The rhyolite is highly enriched in incompatible trace elements including uranium (up to ca. 20 ppm). Cumberland Hill peralkaline rhyolite suggests that a potentially economic source of uranium exists in the basin, based on the elevated uranium concentrations and high potential for remobilization into surrounding sedimentary units.

## RÉSUMÉ

Les roches volcaniques felsiques de la Formation de Cumberland Hill, au sud du Nouveau-Brunswick, dont l'origine remonte à quelque 335 Ma se composent de rhyolite et de trachyte hyperalkalines. On estime que ces roches ont subi une cristallisation fractionnée intense à partir d'un magma de basalte alcalin, même si leurs diverses signatures géochimiques indiquent que la rhyolite n'est pas issue de la trachyte par un processus permanent de cristallisation fractionnée. La rhyolite est fortement enrichie d'éléments traces incompatibles, au nombre duquel se trouve de l'uranium (concentration qui peut atteindre environ 20 ppm). Si l'on se fie aux fortes concentrations d'uranium et à la grande possibilité de remobilisation dans les unités sédimentaires environnantes, la présence de la rhyolite hyperalkaline dans la Formation de Cumberland Hill porte à croire que le bassin pourrait contenir une source d'uranium vraisemblablement rentable.

*[Traduit par la rédaction]*

## INTRODUCTION

Igneous rocks are sparsely distributed throughout the Middle Devonian–Permian Maritimes Basin (Fyffe and Barr 1986; Dunning *et al.* 2002). They appear to decrease in volume through time from: (i) Middle to Late Devonian tholeiite, minor alkali basalt, bimodal gabbro and A-type to evolved I-type granite; (ii) Lower Carboniferous bimodal igneous rocks, and most sparsely, (iii) Upper Carboniferous bimodal igneous rocks (New Brunswick Department of Natural Resources 2010a). The felsic igneous rocks, as well as the interbedded lithic sandstone, conglomerate, and feldspathic sandstone

of the Cumberland Hill Formation, the topic of this paper, were considered part of the Late Wesphalian Pictou Group, and consequently associated with the latter episode of magmatism based on stratigraphic relationships (Fyffe and Barr 1986). However, recent mapping indicated that these volcanic rocks lie within the Visean to early Namurian Mabou Group, and therefore are associated with the middle episode of volcanic activity (St. Peter 1997; New Brunswick Department of Natural Resources 2010a). A mid-Visean U–Pb zircon date of  $335 \pm 2$  Ma substantiates this interpretation (New Brunswick Department of Natural Resources 2010a). Hence the felsic volcanic rocks in the Cumberland Hill Formation appear to

have approximately the same age as several alkali mafic units in New Brunswick (e.g., the Hardwood Ridge, Royal Road, and Queenstown basalts), and the tholeiitic-alkali mafic rocks of the Cap aux Diables Formation in the Magdalen Islands (Barr *et al.* 1985; Fyffe and Barr 1986; LaFleche *et al.* 1998; New Brunswick Department of Natural Resources 2010a).

The mafic rocks have continental within-plate characteristics (Barr *et al.* 1985), and HIMU-OIB Pb-isotopic signatures that have been related either to a mantle plume (Pe-Piper and Piper 1998) or to decompression melting below a pull-apart rift (LaFleche *et al.* 1998). However, the plume model (Murphy *et al.* 1999; Keppie and Krogh 1999) appears inconsistent with U-Pb chronology of Devonian to early Carboniferous volcanic rocks within the southern Magdalen Basin (Dunning *et al.* 2002).

In this paper, we selected key samples from the Cumberland Hill Formation in order to petrologically and geochemically characterize the rocks and compare the rhyolite and trachyte to similar rocks elsewhere in the Maritimes Basin. This work is critical when assessing uranium potential in the Cumberland Hill Formation, as similar rocks in the Maritimes Basin have been identified as hosts of economic uranium mineralization (e.g., Dahlkamp 1993; Plant *et al.* 1999; Cuney 2009; Nash 2010).

## GEOLOGICAL SETTING

Late Devonian to Early Permian rocks of New Brunswick comprise the western part of the Maritimes Basin of Atlantic Canada. The Maritimes Basin is a successor basin within which there are a number of sub-basins separated by fault-bounded ridges of basement (Fig. 1a). In New Brunswick, these rocks occur within deep depositional centres or subbasins, or on shallowly buried or partially exposed basement uplifts and platforms (e.g., St. Peter and Johnson 2009).

The volcanic rocks of the Cumberland Hill Formation crop out as several inliers and are unconformably overlain by the Upper Carboniferous Pictou Group on the New Brunswick Platform (Fig. 1b). The outcrops are likely related to a single volcanic centre partially hidden beneath Pictou Group rocks, because the area incorporating the inliers clearly defines a circular area of high magnetic response on aeromagnetic maps (Thomas and Kiss 2005).

Fyffe and Barr (1986) examined some of the felsic volcanic rocks in the vicinity of Cumberland Hill, including lava flows and tuffs in a rhyolitic unit and underlying trachytic unit. Subsequently, St. Peter (1997) delineated the distribution of these units in more detail, depicting the trachyte as mostly underlying, but in part interdigitating with, the rhyolite (Fig. 2). There is also one massive intrusive porphyry plug with several smaller dykes/pipes, interpreted by Thomas and Kiss (2005) as volcanic feeder dykes, based on their bulls-eye positive magnetic anomalies (50–375 nT) within the magnetically anomalous area. The exact thickness of these units is unknown as they

are poorly exposed; however, both trachyte and rhyolite units are believed to be tens of metres thick (St. Peter 1997).


## PETROGRAPHY

The rhyolite occurs as aphanitic to porphyritic flows and ignimbritic tuffs with flow banding and microporphyritic to porphyritic textures (Fig. 3a). Flow-aligned phenocrysts of feldspar (mainly anorthoclase; Fyffe and Barr 1986; 0.1–0.5 mm in length) and quartz (0.3–1 mm) are enclosed in a quartzofeldspathic groundmass with rare green clinopyroxene (hedenbergite; Fyffe and Barr 1986). The groundmass is composed of quartz, feldspar, devitrified glass, minor magnetite, ilmenite, pyrite, rare galena, fluorite and accessory minerals similar to those in trachyte samples (i.e., microlites of zircon, apatite, monazite, and allanite). The rhyolite is also extensively altered, although to a lesser degree than the trachyte, with most primary magmatic phases replaced by a secondary assemblage (i.e., carbonate, sericite, albite, and chlorite).

The trachyte occurs in two varieties, porphyritic and aphanitic, the latter of which is dominant. The porphyritic trachyte contains phenocrysts of feldspar (sanidine or anorthoclase) up to 2 mm in size, rare clinopyroxene, and plagioclase (Fig. 3b). The groundmass is largely composed of devitrified glass with phenocrysts of feldspar, clinopyroxene, and quartz, and rare ilmenite, hematite, sphalerite, and galena. Accessory minerals (fine-grained microlitic zircon, apatite, monazite, and allanite) were identified and confirmed using a scanning electron microscope including backscattered electron imaging, energy-dispersive X-ray spectroscopy and X-ray element mapping. The aphanitic variety displays a typical trachytic texture with flow-aligned feldspars (0.1–0.5 mm in length) enclosed in a fine-grained matrix similar to that of the porphyritic type. Both varieties are pervasively altered, with only relicts of primary minerals present, and secondary minerals dominated by calcite, sericite, chlorite, albite, and hematite alteration.

## GEOCHEMISTRY

Fifteen samples were collected from the Cumberland Hill Formation, crushed and pulped in a soft iron swing mill, then analyzed for major elements using glass discs by X-ray fluorescence (XRF) spectrometry in the Nova Scotia Regional Geochemical Centre at Saint Mary's University, Halifax. Samples were subsequently analyzed for trace element compositions using a Perkin Elmer Optima 3000 inductively coupled plasma-mass spectrometer in the Activation Laboratories in Ancaster, Ontario. Prior to analysis, samples were digested in nitric acid after lithium metaborate fusion to ensure the complete dissolution of refractory accessory minerals. Precision and accuracy of the XRF data are reported by Dostal *et al.* (1994). The analytical error of the trace element determinations is 2–10%.




**Fig. 1.** (a) Map of Atlantic Canada depicting the Maritimes Basin, adapted from Roliff (1962), Williams (1974), Bradley (1982), and Fyffe and Barr (1986). Major faults are indicated as dashed lines; study area is indicated with a box. Red circles, blue stars and green squares represent different types of uranium occurrences within the Maritimes Basin (McLeod and Smith 2010; Dostal *et al.* 1983); (b) Geological map of the Cumberland Hill area, southern New Brunswick (adapted from Smith 2007) indicating the areal extent of the Cumberland Hill trachyte and rhyolite, both of which are interbedded with lithic and feldspathic sandstone and conglomerate, and contain interdigitating feldspar porphyry, rhyolite ash tuff, breccia, and lapilli tuff (New Brunswick Department of Natural Resources 2010a). The Hurley Creek Formation is composed of mudstone, fine-grained sandstone, coarse-grained feldspathic and lithic sandstone and polymictic pebble conglomerate at the top (Muller 1951). The Minto Formation contains feldspathic, pebbly and fine-grained sandstone, polymictic conglomerate and minor laminated mudstone, shale and thin coal seams (Dyer 1926; Muller 1951; St. Peter 2000).




According to the  $\text{Na}_2\text{O} + \text{K}_2\text{O}$  vs  $\text{SiO}_2$  classification of LeMaitre *et al.* (1989; Fig. 4a) and the  $\text{Zr}/\text{TiO}_2$  vs  $\text{SiO}_2$  diagram (Fig. 4b) of Winchester and Floyd (1977), the rocks can be subdivided into two groups, trachyte and peralkaline rhyolite, analogous to comendite/pantellerite fields according to Winchester and Floyd (1977). Most rocks are peralkaline with a

high agpaite index ( $\text{AI} = \text{mole}(\text{Na}+\text{K})/\text{Al}$ ; Shand 1951) and acmite in their CIPW normative mineralogy. Considering loss on ignition-free analyses (Table 1), the trachyte on average ( $n = 7$ ,  $1\sigma$ ) has moderate  $\text{SiO}_2$  ( $65 \pm 2.48$  wt. %) accompanied by high total alkalis ( $9.5 \pm 1.81$  wt. %), low  $\text{CaO}$  ( $2.5 \pm 1.21$  wt. %), high  $\text{FeO}^*/\text{MgO}$  (ca. 25) and elevated concentrations of high field



**Fig. 2.** Carboniferous stratigraphy in the central New Brunswick Basin (adapted from St. Peter 1997). The Cumberland Hill and Hardwood Ridge formations are part of the Mabou Group, whereas the Minto Formation, Hurley Creek Formation and Sunbury Creek Formation belong to the Pictou Group.



**Fig. 3.** Slab photographs indicating (a) flow-banded rhyolite with a quartzofeldspathic groundmass, and (b) vuggy trachyte with chlorite infillings.

**Table 1.** Major and trace element concentrations in trachyte and rhyolite from the Cumberland Hill Formation.


	Trachyte							Rhyolite							
	46°01'56"N, 65°51'57"W to 46°01'50"N, 65°52'04"W							46°01'11"N, 65°51'15"W to 46°01'13"N, 65°51'14"W							
	1-202	1-203	1-206	1-207	1-01	1-12	1-14	2-01	2-8A	2-8B	2-09	2-10	2-11	2-12	2-13
SiO <sub>2</sub> (wt. %)	61.71	59.93	66.95	63.63	64.15	62.39	62.30	74.22	72.81	73.40	72.72	71.69	71.85	72.42	74.29
TiO <sub>2</sub>	0.48	0.47	0.63	0.37	0.60	0.63	0.38	0.17	0.17	0.17	0.17	0.17	0.17	0.18	0.18
Al <sub>2</sub> O <sub>3</sub>	14.29	14.14	13.87	14.94	13.75	13.51	14.43	11.64	11.23	11.44	11.25	11.60	11.71	11.65	11.93
Fe <sub>2</sub> O <sub>3</sub> <sup>T</sup>	9.30	8.80	4.07	7.83	6.11	5.10	7.84	3.39	4.16	4.13	4.13	4.37	4.22	4.36	3.38
MnO	0.18	0.25	0.11	0.22	0.19	0.17	0.22	0.04	0.10	0.10	0.12	0.10	0.10	0.09	0.03
MgO	0.13	0.37	0.20	0.08	0.37	0.29	0.07	0.02	0.02	-	0.02	0.02	0.05	0.02	0.03
CaO	1.52	2.36	2.30	1.69	3.13	4.50	1.38	0.11	0.15	0.15	0.22	0.31	0.26	0.23	0.34
Na <sub>2</sub> O	4.78	4.43	3.73	5.77	3.03	2.21	6.17	3.65	3.93	4.21	3.67	3.81	4.26	4.50	5.33
K <sub>2</sub> O	4.93	5.02	4.97	5.16	4.85	3.87	5.68	4.59	4.96	4.84	5.56	4.97	4.53	4.18	3.46
P <sub>2</sub> O <sub>5</sub>	0.14	0.13	0.23	0.09	0.17	0.19	0.08	-	0.01	-	0.01	-	-	-	-
LOI	1.84	2.48	2.84	0.66	4.02	6.31	0.62	0.96	1.31	1.17	1.99	1.80	2.06	1.03	1.40
Total	99.30	98.37	99.89	100.44	100.37	99.17	99.17	98.79	98.85	99.61	99.85	98.84	99.22	98.66	100.37
Pb (ppm)	26.0	13.0	18.0	21.0	9.00	16.0	90.0	15.0	62.0	21.0	37.0	28.0	82.0	24.0	17.0
Zn	250	150	110	210	90.0	60.0	210	390	280	300	360	300	300	310	290
W	1.60	1.40	1.80	1.40	1.10	3.50	0.70	3.20	6.60	8.20	2.90	3.50	3.40	22.9	9.40
Rb	81.0	79.0	92.0	104	98.0	91.0	124	308	290	299	347	306	262	244	192
Cs	0.60	0.20	1.30	0.70	1.70	1.70	0.70	1.90	1.70	1.90	1.90	2.10	1.50	1.70	1.10
Ba	384	403	552	259	664	398	280	15.00	4.00	-	12.0	6.00	9.00	-	9.00
Sr	107	281	100	42.0	93.0	96.0	39.0	25.0	16.0	17.0	37.0	39.0	38.0	21.0	28.0
Ga	36.0	34.0	26.0	40.0	30.0	28.0	39.0	56.0	54.0	58.0	57.0	57.0	56.0	58.0	55.0
Ta	6.72	6.68	3.67	8.09	3.70	4.33	8.29	15.1	18.4	18.9	18.4	18.2	18.4	19.4	18.7
Nb	95.5	92.6	51.5	118	51.1	65.6	118	281	235	287	222	249	223	306	208
Hf	22.3	21.9	11.7	24.9	14.4	16.0	31.0	57.3	51.1	55.5	53.0	50.4	51.7	54.5	55.0
Zr	991	944	557	1150	679	755	1550	2640	2190	2420	2290	2300	2280	2380	2450
Y	86.90	83.3	51.7	101	54.4	62.3	88.8	177	179	175	179	184	162	187	191
Th	11.5	11.1	11.3	12.2	12.6	13.6	14.8	29.6	31.2	30.8	38.7	32.9	29.2	33.1	35.0
U	2.36	2.35	4.81	3.71	4.65	5.79	3.99	20.80	9.27	11.70	18.50	13.80	16.10	10.20	13.80
La	114	116	93.0	139	86.0	88.3	120	164	150	151	144	142	146	148	146
Ce	239	243	195	286	183	189	259	337	331	334	330	327	315	319	322
Pr	25.9	26.3	20.7	30.5	21.7	22.1	30.6	39.3	39.1	39.5	37.7	37.5	38.3	38.7	38.7
Nd	92.5	94.8	70.3	110	86.6	88.8	122	152	155	155	147	147	151	152	152
Sm	20.8	20.7	14.9	23.7	17.1	17.5	24.8	35.6	36.6	37.3	35.3	35.5	36.1	36.3	36.4
Eu	5.33	5.40	3.62	5.40	3.40	3.55	4.85	3.06	3.17	3.17	3.11	3.08	3.18	3.12	3.20
Gd	19.3	19.6	13.3	22.2	14.0	14.7	21.5	33.2	34.9	35.9	34.6	34.7	34.8	35.5	36.0
Tb	3.12	3.04	2.02	3.49	2.10	2.26	3.38	6.08	6.04	6.11	6.12	6.12	6.00	6.27	6.35
Dy	16.9	16.6	10.8	19.7	11.3	12.6	18.4	33.3	34.5	34.3	35.1	35.1	33.0	35.9	36.1
Ho	3.15	3.12	2.00	3.66	2.17	2.46	3.56	6.62	6.67	6.67	7.16	6.97	6.24	7.13	7.20
Er	8.85	8.61	5.61	10.0	5.91	6.78	9.72	18.2	18.1	17.8	19.5	19.0	16.6	19.4	19.6
Tm	1.27	1.24	0.81	1.42	0.88	1.02	1.45	2.85	2.64	2.63	2.95	2.93	2.55	3.01	2.97
Yb	8.35	7.98	5.29	9.07	5.61	6.50	9.37	17.6	17.7	15.5	17.0	17.9	15.7	18.1	17.7
Lu	1.30	1.30	0.80	1.42	0.89	1.03	1.48	2.54	2.68	2.23	2.36	2.63	2.30	2.60	2.56

Note: - symbol indicates no data available, or value below detection limits


strength elements (HFSE) and other incompatible elements (e.g.,  $Zr + Nb + Ce + Y = 1334 \pm 409.70$ ; Whalen *et al.* 1987). The average of the rhyolitic rocks ( $n = 8$ ,  $1\sigma$ ) has higher  $SiO_2$  ( $75.0 \pm 0.66$  wt. %), equivalent total alkali concentrations ( $9.0 \pm 0.29$  wt. %), very low  $CaO$  ( $0.23 \pm 0.08$  wt. %), high  $FeO^I/MgO$  (ca. 140), and high HFSE and other incompatible elements (e.g.,  $Zr + Nb + Ce + Y = 3126 \pm 161.75$ ; Whalen *et al.* 1987).

The major and trace element signatures of the volcanic rocks of the Cumberland Hill Formation are interpreted to vary predominantly due to the degree of fractionation. Conversely, a heterogeneous magma source could also contribute to the variations exhibited. In order to determine the fractionation trends, Nb was utilized as a differentiation index following White *et al.* (2006; Fig. 5). With increasing concentration of Nb, the trachyte samples display positive correlation with  $Al_2O_3$ , and negative correlation with  $TiO_2$ . The reduction in  $TiO_2$  concentration can most likely be attributed to crystallization of Fe-Ti oxides. Rhyolite samples, in contrast, have relatively constant  $Al_2O_3$  (Fig. 5a) and  $TiO_2$  (Fig. 5b). Trace elements also display different fractionation trends in trachyte and rhyolite samples. Trachyte samples show marked decreases in Ba with increasing Nb (Fig. 5c), reflecting the crystallization of feldspars, as well as an increase in incompatible trace elements including Zr, La, and Ga (Fig. 5d, e, f). However, the abundances of these trace elements in the rhyolite remain virtually unchanged with increasing Nb. The contrast in fractional crystallization trends between trachyte and rhyolite is highlighted by the  $CaO$ - $K_2O$ - $Na_2O$  ternary diagram (Fig. 6), which illustrates that trachyte samples plot in a trend away from the  $CaO$  apex, whereas rhyolite sample variation is parallel to the  $K_2O$ - $Na_2O$  join. This difference is most readily explained by the crystallization of Ca-bearing plagioclase and clinopyroxene in the trachyte magma, and alkali feldspar crystallization in the rhyolite magma.

The chondrite-normalized rare earth element (REE) patterns for each suite (Fig. 7a) are strongly enriched in light REE (LREE). The rhyolite has higher LREE contents with  $La_n$  values of ca. 440–510 compared to the trachyte with  $La_n$  ca. 270–430. The trachyte shows steeper LREE to heavy rare earth element (HREE) profiles, with  $La_n/Yb_n$  ratios ca. 8–11 compared to ca. 4–6 in rhyolite. A similar relationship is observed for  $La_n/Sm_n$ , which in the trachyte is ca. 2.6–3.4 and in the rhyolite is ca. 2.2–2.6. The shape of the REE patterns in the trachyte does not change significantly, although the absolute abundances are escalating with the increasing degree of differentiation. The REE patterns of the rhyolite are characterized by a pronounced negative Eu anomaly (average  $Eu/Eu^* = 0.08$ ), consistent with low-pressure fractional crystallization of alkali and/or plagioclase feldspar. Primitive mantle-normalized trace element patterns in both rock types (Fig. 7b) are fairly similar. Both are highly fractionated and peak at Th-Nb; however, the rhyolite displays strong, negative Ba, Sr, and moderately negative Eu anomalies, whereas the trachyte contains only slightly negative anomalies in these elements. The Nb/Ta ratios of the trachyte ( $14.27 \pm 0.47$ ) and rhyolite ( $13.91 \pm 2.48$ ) are the same,



**Fig. 4.** Plots of (a)  $Na_2O + K_2O$  versus  $SiO_2$  (wt. %; Le Maitre *et al.* 1989); (b)  $SiO_2$  versus  $Zr/TiO_2$  (wt. %; Winchester and Floyd 1977); and (c)  $Zr/TiO_2$  versus  $Nb/Y$  (Winchester and Floyd 1977) for volcanic samples from the Cumberland Hill Formation.




**Fig. 5.** Plots of Nb (ppm) versus (a)  $\text{Al}_2\text{O}_3$  (wt. %), (b)  $\text{TiO}_2$  (wt. %), (c) Ba (ppm), (d) Zr (ppm), (e) La (ppm), and (f) Ga (ppm) for volcanic samples from the Cumberland Hill Formation.

but slightly lower than that of the mantle (typically  $\sim 17.5$ ; Sun and McDonough 1989).

The abundances of U and Th in the felsic rocks of the Cumberland Hill Formation are high and variable. Rhyolite display significantly higher contents of U ( $14.0 \pm 4.0$  ppm,  $n = 8$ ), and Th ( $33.0 \pm 3.1$  ppm,  $n = 8$ ) compared to trachyte (U =

$2.4 \pm 2.4$  ppm,  $n = 7$ ; Th =  $12 \pm 1.4$  ppm,  $n = 7$ ). The large range in U may be related to devitrification and alteration of the rhyolite groundmass. The rhyolite samples also have a higher U/Th ratio ( $0.44 \pm 0.13$ ,  $n = 8$ ) than trachyte ( $0.32 \pm 0.09$ ,  $n = 7$ ), which is comparable to the whole rock or melt inclusion data from non-mineralized peralkaline rhyolite from various






**Fig. 6.** CaO-K<sub>2</sub>O-Na<sub>2</sub>O ternary diagram for volcanic samples from the Cumberland Hill Formation. Trends indicate that fractionation of Ca-plagioclase and clinopyroxene could have occurred in trachyte, whereas alkali feldspar fractionation is more likely in rhyolite.

U districts in the world (Cuney and Kyser 2009). The increase of U/Th ratio in the rhyolite compared to trachyte may reflect monazite fractionation. Based on the variability of the U and Th concentrations in the rhyolite and trachyte samples, we interpret that remobilization of these elements has occurred. This remobilization was likely caused by devitrification of the groundmass because zircon is modally insignificant, and not the principal host of these elements based on the lack of correlation of radioactive trace elements with Zr (see Dupuy and Dostal 1983).


## PETROGENESIS

Overall the geochemical and mineralogical similarities of the spatially and temporally associated trachyte and rhyolite imply that they are genetically related. However, discontinuous and/or contrasting variation trends and significant differences in incompatible trace element ratios indicate that the rhyolite was not derived from the trachyte via continuous fractional crystallization, although fractional crystallization may be responsible for the variations within the individual suites. The evolution of the trachyte in the melt phase involved crystallization of feldspars, clinopyroxene, and Fe-Ti oxides, whereas the evolution of the rhyolite was dominated by crystallization and fractionation of alkali feldspar.




**Fig. 7.** (a) Chondrite-normalized REE patterns of the averages of trachyte ( $n = 7$ ) and rhyolite ( $n = 8$ ) samples from the Cumberland Hill Formation and rhyolite from Streltsovka ( $n = 1$ ; Chabiron *et al.* 2003); (b) Primitive mantle-normalized multi-element plots of the average of trachyte ( $n = 7$ ) and rhyolite ( $n = 8$ ) samples from the Cumberland Hill Formation and rhyolite from Streltsovka ( $n = 1$ ; Chabiron *et al.* 2003). Normalizing values are after Sun and McDonough (1989).

An alternative model to continuous fractional crystallization for the origin of the rhyolite is assimilation-fractional crystallization. However, the low and overlapping Th/Ta ratio of the trachyte and rhyolite (Fig. 8a), a sensitive indicator of crustal contamination (Gorton and Schandl 2000), demonstrates that assimilation-fractional crystallization and crustal contamination processes cannot solely account for derivation of the rhyolite from the trachyte. In addition, Pe-Piper and Piper (1998) reported an  $\epsilon_{\text{Nd}}$  value of +3, with an assumed age of ca. 310 Ma from the Cumberland Hill rhyolite. This relatively high value implies that significant assimilation of crustal material is not likely to have taken place.



**Fig. 8.** Discrimination diagrams for felsic volcanic rocks from the Cumberland Hill Formation. (a) Yb (ppm) versus Th/Ta (Gorton and Schandl 2000). Abbreviations: WPVZ = within plate volcanic zones; MORB = mid-ocean ridge basalt; WPB = within-plate basalt. (b) Rb vs Nb + Y and (c) Ta vs Yb (Pearce *et al.* 1984). Abbreviations: syn-COLG = syn-collisional granite; WPG = within-plate granite; VAG = volcanic-arc granite; ORG = oceanic-ridge granite.



**Fig. 9.** Plot of K/Rb versus (a) Rb (ppm) and (b) U (ppm) in felsic volcanic samples from the Cumberland Hill Formation.

K/Rb ratios in trachyte samples are much higher than in rhyolite samples (Fig. 9a, b), which suggest that the evolution of the rhyolite may have been affected by fluids (Dostal and Chatterjee 1995). Again, this mechanism cannot explain the origin of the rhyolite, but rather indicates that fluids played an important role during the evolution of these rocks and perhaps even led to the mobilization of U (Cuney and Kyser 2009; Nash 2010) in the rhyolite (Fig. 9b).

A comparison with similar rocks elsewhere (e.g., Pecerrillo *et al.* 2003) suggests that the Cumberland Hill trachyte and rhyolite could represent various stages of fractional crystallization in an evolving alkali basaltic magma. The absence of a negative Nb anomaly suggests that the parent magmas were not derived from subcontinental lithospheric mantle, but more plausibly originated from asthenospheric mantle. This inter-

pretation is supported by the rhyolite  $\epsilon_{\text{Nd}}$  value of +3 (Pe-Piper and Piper 1998), which implies upward migration of enriched asthenospheric magma into a juvenile lithosphere.

LaFleche *et al.* (1998) and Pe-Piper and Piper (1998) described alkali basaltic rocks of comparable age from other parts of the Maritimes Basin. The presence of alkali gabbroic rocks of a similar age (Johnson 2008) in the area suggests that the felsic rocks of the Cumberland Hill Formation could be derived from such parent magma by fractional crystallization. In fact, peralkaline felsic volcanic complexes are typically associated with a shallow-seated alkali intrusions ranging in composition from gabbro to highly fractionated granitic rocks. Although geochemical data are insufficient from the mafic alkali rocks in the vicinity of the Cumberland Hill Formation, chemical analyses of alkali basaltic rocks reported by LaFleche *et al.* (1998) from the Magdalen Islands indicate that similar basalts could have been a parent to trachyte and rhyolite. Trachyte of similar compositions have been documented to evolve from alkali basalt by extensive fractional crystallization (Peccherillo *et al.* 2003) and peralkaline rhyolite could have evolved from such magma by further fractional crystallization in a subvolcanic magma chamber. A likely scenario for the Cumberland Hill felsic rocks is that they are the products of fractional crystallization of alkali basaltic magma, and that the trachyte and rhyolite represent separate magma pulses with the latter being younger having undergone more extensive fractionation prior to emplacement.

### ECONOMIC POTENTIAL

Silica-oversaturated peralkaline felsic volcanic rocks, particularly rhyolite, are commonly enriched in U and represent a potential source of uranium for volcanic-related hydrothermal U deposits, such as the Streltsovka caldera (Transbaikalia, Russia: Ischukova 1997; Chabiron *et al.* 2001, 2003; Cuney and Kyser 2009; Nash 2010), sediment-hosted U deposits in siliciclastic sedimentary basins, such as the U deposits in the Tim Merso basin in Niger (Forbes *et al.* 1984; Forbes 1989; Plant *et al.* 1999; Pagel *et al.* 2005; International Atomic Energy Agency 2009), and the New Horton mineralization (U-Cu-Ag-Au) within the Carboniferous basin (Boyd 1978; McLeod and Smith 2010; New Brunswick Department of Natural Resources 2010b).

The crystallization of U-bearing accessory minerals in highly depolymerized peralkaline melts is typically suppressed (Cuney 2009; Cuney and Kyser 2009) leading to progressive enrichment of U in residual melts during fractional crystallization and entrapment of U in the glassy groundmass. Subsequently U can be leached during the alteration or devitrification of the glass (Nash 2010). Cuney (2009) and Cuney and Kyser (2009) inferred that ignimbritic tuffs, in which U is hosted in the glass, are the most favourable type of volcanic rock; such rhyolitic pyroclastic rocks are abundant in the Cumberland Hill Formation.

Rhyolite of the Cumberland Hill Formation has relatively

high abundances of U compared to associated trachyte and could be a source of U mineralization. The U concentrations in these rhyolite samples are elevated compared to similar felsic and mafic Late Tournaisian–Early Viséan deposits of the Maritimes Basin (Magdalen Islands, 0.4–1.9 ppm; Central New Brunswick, 0.8–1.9 ppm and the Cobequid Highland dykes, 0.2 ppm; Pe-Piper and Piper 1998). The abundances of U in the rhyolite are comparable to the non-mineralized U concentrations of the peralkaline rhyolite of the Streltsovka caldera (Fig. 7b), the largest, volcanic-related, U ore field in the world (Cuney 2009). Despite similar U concentration, it is important to consider where U is hosted in rhyolite. When U is hosted in zircon or other resistant accessory minerals it cannot be readily remobilized. Based on current geochemical work, it appears U is present within the glassy groundmass of the rhyolite in the Cumberland Hill Formation and can therefore be released and re-deposited and thus could be a source of U within U- (Cu-V-Ag) deposits within local parts of the Carboniferous basin.

### CONCLUSIONS

Felsic volcanic rocks from the Cumberland Hill Formation consist of peralkaline rhyolite and trachyte. Both contain elevated concentrations of incompatible trace elements, with the rhyolite more enriched and fractionated than the trachyte. Although both rock types appear to be genetically related, differences in their geochemical signatures imply the rhyolite was not derived from the trachyte by continuous fractional crystallization. Examination of the Th/Ta ratio also indicated crustal contamination cannot account for the observed compositional contrast. A preferred model is that the trachyte and rhyolite originated from the same or highly similar parent magma, and represent separate magma pulses with the latter being younger, having undergone more extensive fractionation prior to emplacement. The absence of a negative Nb anomaly suggests the parent magma was not derived from subcontinental lithospheric mantle, but more plausibly originated from an asthenospheric mantle source. Lastly, Cumberland Hill rhyolite contains elevated U concentrations, with U hosted within the glassy groundmass. Since U can be leached and re-deposited, the rhyolite could be a source of sediment-hosted U mineralization within the basin.

### ACKNOWLEDGEMENTS

A grant from the Geological Survey Branch of the New Brunswick Department of Natural Resources and Energy (to JD) covered fieldwork expenses. Analytical work was predominantly funded by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant (to JD), and in part by a Society of Economic Geologists Student Research Grant (to TRG). The manuscript benefited from constructive reviews by David Lentz, an anonymous reviewer, and journal editor Sandra Barr.

## REFERENCES

- Barr, S.M., Brisebois, D., and MacDonald, A.S. 1985. Carboniferous volcanic rocks of the Magdalen Islands, Gulf of St. Lawrence. *Canadian Journal of Earth Sciences*, 22, pp. 1679–1688.
- Boyd, R.T. 1978. Gulf Minerals Canada Limited report on the New Horton claims Albert County, New Brunswick. Report Number 472212, 36 p.
- Bradley, D.C. 1982. Subsidence in the late Paleozoic Basins in the eastern Appalachians. *Tectonics*, 1, pp. 107–123. [doi:10.1029/TC001i001p00107](https://doi.org/10.1029/TC001i001p00107)
- Chabiron, A., Alyoshin, A.P., Cuney, M., Deloule, E., Golubev, V.N., Velitchkin, V.I., and Poty, B. 2001. Geochemistry of the rhyolitic magmas from the Streltsovka caldera (Transbaikalia, Russia): a melt inclusion study. *Chemical Geology*, 175, pp. 273–290. [doi:10.1016/S0009-2541\(00\)00300-4](https://doi.org/10.1016/S0009-2541(00)00300-4)
- Chabiron, A., Cuney, M., and Poty, B. 2003. Possible uranium sources for the largest uranium district associated with volcanism: the Streltsovka caldera (Transbaikalia, Russia). *Mineralium Deposita*, 38, pp. 127–140.
- Cuney, M. 2009. The extreme diversity of uranium deposits. *Mineralium Deposita*, 44, pp. 3–9. [doi:10.1007/s00126-008-0223-1](https://doi.org/10.1007/s00126-008-0223-1)
- Cuney, M., and Kyser, K. (Editors). 2009. Recent and not-so-recent developments in uranium deposits and implications for exploration. *Mineralogical Association of Canada, Short Course Series 39*, 257 p.
- Dahlkamp, F.J. 1993. *Uranium ore deposits*. Springer-Verlag, Heidelberg. 460 p.
- Dostal, J., and Chatterjee, A.K. 1995. Origin of topaz-bearing and related peraluminous granites of late Devonian Davis Lake pluton, Nova Scotia, Canada. *Chemical Geology*, 123, pp. 67–88. [doi:10.1016/0009-2541\(95\)00047-P](https://doi.org/10.1016/0009-2541(95)00047-P)
- Dostal, J., Dupuy, C., and Keppie, J.D. 1983. uranium and thorium in Paleozoic rhyolites of Nova Scotia. *Canadian Journal of Earth Science*, 20 pp. 266–274.
- Dostal, J., Dupey, C., and Caby, R. 1994. Geochemistry of the Neoproterozoic Tilemsi belt of Iforas (Mali, Sahara): a crustal section of an oceanic island arc. *Precambrian Research*, 65, pp. 55–69. [doi:10.1016/0301-9268\(94\)90099-X](https://doi.org/10.1016/0301-9268(94)90099-X)
- Dunning, G.R., Barr, S.M., Giles, P.S., McGregor, D.C., Piper, G., and Piper, D.J.W. 2002. Chronology of Devonian to early Carboniferous rifting and igneous activity in southern Magdalen Basin based on U-Pb (zircon) dating. *Canadian Journal of Earth Sciences*, 39, pp. 1219–1237. [doi:10.1139/e02-037](https://doi.org/10.1139/e02-037)
- Dupuy, C., and Dostal, J. 1983. Uranium and thorium in Paleozoic basalts of Nova Scotia. *Bulletin of Volcanology*, 46, pp. 125–133. [doi:10.1007/BF02597581](https://doi.org/10.1007/BF02597581)
- Dyer, W.S. 1926. Minto coal basin, New Brunswick. *Geological Survey of Canada, Memoir 151*, 42 p.
- Forbes, P. 1989. Roles des structures sédimentaires et tectoniques, du volcanisme alcalin régional et des fluides diagenétiques hydrothermaux pour la Formation des minéralisations a U-Zr-Zn-V-Mo d'Akouta (Niger). *Géologie Géochimie Uranium Mémoires Nancy*, 17, 376 p.
- Forbes, P., Pagel, M., Pacquet, A., Chantret, F., and Oumarou, J. 1984. Volcanic components in the Akouta uranium deposit (Niger). *Comptes Rendus de l'Académie des Sciences Serie II*. 298, pp. 647–650.
- Fyffe, L.R., and Barr, S.M. 1986. Petrochemistry and tectonic significance of Carboniferous volcanic rocks in New Brunswick. *Canadian Journal of Earth Sciences*, 23, pp. 1243–1256.
- Gorton, M.P., and Schandl, E.S. 2000. From continents to island arcs: a geochemical index of tectonic setting for arc-related and within-plate felsic to intermediate volcanic rocks. *Canadian Mineralogist*, 38, pp. 1065–1073. [doi:10.2113/gscanmin.38.5.1065](https://doi.org/10.2113/gscanmin.38.5.1065)
- International Atomic Energy Agency 2009. *World Distribution of Uranium Deposits (UDEPO) with Uranium Deposit Classification*. Vienna, Austria, 126 p.
- Ischukova, L.P. 1997. The Streltsovskoye uranium district: Vienna, International Atomic Energy Agency. *Tecdoc-961*, pp. 237–250.
- Johnson, S.C. 2008. Bedrock geology of pre-Carboniferous basement inliers in the Coal Creek, Canaan River, and Thorne Brook areas, southeastern New Brunswick Platform. In *Abstracts 2008: Exploration Mining and Petroleum New Brunswick*, Edited by S.A.A. Merlini. New Brunswick Department of Natural Resources, Minerals, Policy and Planning Division, Information Circular 2008-1, 39 p.
- Keppie, J.D., and Krogh, T.E. 1999. U-Pb geochronology of Devonian granites in the Meguma Terrane of Nova Scotia, Canada: evidence for hotspot melting of a Neoproterozoic source. *Journal of Geology*, 107, pp. 555–568. [doi:10.1086/314369](https://doi.org/10.1086/314369)
- LaFleche, M.R., Camire, G., and Jenner, G.A. 1998. Geochemistry of post-Acadian, Carboniferous continental intra-plate basalts from the Maritimes Basin, Magdalen Islands, Quebec, Canada. *Chemical Geology*, 148, pp. 115–136. [doi:10.1016/S0009-2541\(98\)00002-3](https://doi.org/10.1016/S0009-2541(98)00002-3)
- Le Maitre, R.W., Bateman P., Dudek A., Keller J., Lameyre J., Le Bas M.J., Sabine P.A., Schmid R., Sorensen H., Streck-eisen A., Wooley A.R., and Zanettin B. 1989. *A Classification of Igneous Rocks and Glossary of Terms*, Blackwell, Oxford. 193 p.
- McLeod, M.J., and Smith, E.A. 2010. Uranium. New Brunswick Department of Natural Resources; Lands, Minerals and Petroleum Division, Mineral Commodity Profile, 6, 7 p.
- Muller, J.E. 1951. Geology and coal deposits of Minto and Chipman map-areas, New Brunswick. *Geological Survey of Canada, Memoir 260*, 40 p.
- Murphy, J.B., van Staal, C.R., and Keppie, J.D. 1999. Middle to Late Paleozoic Acadian Orogeny in the northern Appalachians: a Laramide-style plume-modified orogeny? *Geology*, 27, pp. 653–656. [doi:10.1130/0091-7613\(1999\)027<0653:MTLPAO>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0653:MTLPAO>2.3.CO;2)
- Nash, J.T. 2010. *Volcanogenic Uranium Deposits: Geology, Geochemical Processes, and Criteria for Resource Assess-*

- ment. US Geological Survey, Open-File Report 2010–1001, 99 p.
- New Brunswick Department of Natural Resources. 2010a. Bedrock lexicon of New Brunswick Minerals and Energy Division. URL <<http://www1.gnb.ca/0078/GeoscienceDatabase/Lexicon/qryFormationSummary-e.asp?UnitId=251>>, April 2010.
- New Brunswick Department of Natural Resources. 2010b. Mineral Occurrence Data Base. URL <<http://dnre-mrne.gnb.ca/mineraloccurrence/>>, April 2010.
- Pagel, M., Cavellec, S., Forbes, P., Gerbaud, O., Vergely, P., and Wagani, I. 2005. Uranium deposits in the Arlit area (Niger). In *Mineral deposit research, Meeting the global challenge. Edited by J. Mao and M.P. Bierlein. Proceedings of the 8th SGA Meeting in Beijing, China*, pp. 303–305.
- Pearce, J.A., Harris, N.B.W., and Tindle, A.G. 1984. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *Journal of Petrology*, 25, pp. 956–983.
- Peccerillo, A., Barberio, M.R., Yirgu, G., Ayalew, D., Barbieri, M., and Wu, T.W. 2003. Relationships between mafic and peralkaline silicic magmatism in continental rift settings: a petrological, geochemical, and isotopic study of the Gedomsa Volcano, central Ethiopian Rift. *Journal of Petrology*, 44, pp. 2003–2032. doi:10.1093/petrology/egg068
- Pe-Piper, G., and Piper, D.J.W. 1998. Geochemical evolution of Devonian-Carboniferous igneous rocks of the Magdalen basin, Eastern Canada: Pb- and Nd-isotope evidence for mantle and lower crustal sources. *Canadian Journal of Earth Science*, 35, pp. 201–221. doi:10.1139/cjes-35-3-201
- Plant, J., Simpson, P.R., Smith, B., and Windley, B.F. 1999. Uranium ore deposits: products of the radioactive earth. In *Uranium—Mineralogy, geochemistry, and the environment. Edited by P.C. Burns and Robert Finch. Mineralogical Society of America, Reviews in Mineralogy* 38, pp. 255–320.
- Roliff, W.A., 1962. The Maritime Carboniferous basin of eastern Canada. *Geological Association of Canada, Proceedings* 14, pp. 21–41.
- Shand, S.J. 1951. *Eruptive rocks; their genesis, composition, classification, and their relation to ore-deposits*. 4<sup>th</sup> edition. John Wiley and Sons, N.Y. p. 488.
- Smith, E.A. (Compiler) 2007. Bedrock geology of the Chipman Area (NTS 21 I/04). Sunbury and Queens Counties, New Brunswick. Department of Natural Resources: Mineral, Policy and Planning Division. Plate 2007-50, scale 1:50 000.
- St. Peter, C. J. 1997. Bedrock geology of Chipman – Canaan River map area (parts of NTS 21 I/04 and 21 H/13), Sunbury, Queens and Kings counties, New Brunswick. New Brunswick Department of Natural Resources and Energy, Mineral Resources, Plate 97-34, scale 1:20 000.
- St. Peter, C.J. 2000. Carboniferous geology of the southwestern New Brunswick platform (Maugerville Subbasin). New Brunswick Department of Natural Resources and Energy, Mineral Resources, Plate 2000-16, scale 1:150 000.
- St. Peter C.J., and Johnson S.C. 2009. Stratigraphy and structural history of the late Paleozoic Maritimes Basin in southeastern New Brunswick, Canada. New Brunswick Department of Natural Resources; Minerals, Policy and Planning Division, Memoir 3, 348 p.
- Sun, S.S., and McDonough, W.F. 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. In *Magmatism in the Ocean Basins. Edited by A.D. Saunders and M.J. Norry. Geological Society London, Special Publication* 42, pp. 313–345.
- Thomas, M.D., and Kiss, F. 2005. Geological interpretation of the 2004 Marrtown aeromagnetic survey, Southeastern New Brunswick. Geological Survey of Canada, Open File 4953, New Brunswick Department of Natural Resources, Minerals, Policy and Planning Division. Plate 2005-21D, scale 1:125 000.
- Whalen, J.B., Currie, K.L., and Chappell, B.W. 1987. A-type granites: geochemical characteristics, discrimination and petrogenesis, *Contributions to Mineralogy and Petrology*, 95, pp. 407–419. doi:10.1007/BF00402202
- White, J.C., Benker, S.C., Ren, M., Urbanczyk, K.M., and Corrick, D.W. 2006. Petrogenesis and tectonic setting of the peralkaline Pine Canyon caldera, Trans-Pecos Texas, USA. *Lithos*, 91, pp. 74–94. doi:10.1016/j.lithos.2006.03.015
- Williams, E.P. 1974. Geology and petroleum possibilities in and around Gulf of St. Lawrence. *American Association of Petroleum Geologists, Bulletin* 58, 6, pp. 1137–1155.
- Winchester, J.A., and Floyd, P.A. 1977. Geochemical discrimination of different magma series and their differentiation products using immobile elements. *Chemical Geology*, 20, pp. 325–343. doi:10.1016/0009-2541(77)90057-2

*Editorial responsibility: Sandra M. Barr*