

## **Igneous Rock Associations 13** **Focusing on the Central American Subduction Zone**

James A. Walker and Esteban Gazel

Volume 41, Number 1, 2014

URI: <https://id.erudit.org/iderudit/1023630ar>

[See table of contents](#)

Publisher(s)

The Geological Association of Canada

ISSN

0315-0941 (print)

1911-4850 (digital)

[Explore this journal](#)


Cite this article

Walker, J. A. & Gazel, E. (2014). Igneous Rock Associations 13: Focusing on the Central American Subduction Zone. *Geoscience Canada*, 41(1), 57–74.

### Article abstract

Central America has recently been an important focus area for investigations into the complex processes occurring in subduction zones. Here we review some of the new findings concerning subduction input, magma production and evolution, and resultant volcanic output. In the Nicaraguan portion of the subduction zone, subduction input is unusually wet, likely caused by extensive serpentinization of the mantle portion of the incoming plate associated with bending-related faulting seaward of the Middle America trench. The atypical influx of water into the Nicaraguan section of the subduction zone ultimately leads to a regional maximum in the degree of mantle melting. In central Costa Rica, subduction input is also unusual in that it includes oceanic crust flavored by the Galapagos plume. Both of these exotic subduction inputs are recognizable in the compositions of magmas erupted along the volcanic front. In addition, Nicaraguan magmas bear a strong chemical imprint from subducting hemipelagic sediments. The high-fieldstrength- element depletions of magmas from El Salvador through Costa Rica are related to local variations in the depth to the subducting Cocos plate and, therefore, to segmentation of the volcanic front. Minor phases, probably amphibole or rutile, control these variable depletions. Silicic magmas erupted along the volcanic front exhibit the same along-arc geochemical variations as their mafic brethren. This and their mantle-like radiogenic isotopic compositions suggest the production of juvenile continental crust all along the Central American subduction zone. Punctuated times of enhanced magmatic input from the mantle may aid in crustal development.

# SERIES



## Igneous Rock Associations 13. Focusing on the Central American Subduction Zone

James A. Walker<sup>1</sup> and Esteban  
Gazel<sup>2</sup>

<sup>1</sup>*Department of Geology and Environmental  
Geosciences  
Northern Illinois University  
DeKalb, IL, 60115, USA  
E-mail: jwalker@niu.edu*

<sup>2</sup>*Department of Geosciences  
Virginia Polytechnic Institute and  
State University  
Blacksburg, VA, 24061, USA*

### SUMMARY

Central America has recently been an important focus area for investigations into the complex processes occurring in subduction zones. Here we review some of the new findings concerning subduction input, magma production and evolution, and resultant volcanic output. In the Nicaraguan portion of the subduction zone, subduction input is unusually wet, likely caused by exten-

sive serpentinization of the mantle portion of the incoming plate associated with bending-related faulting seaward of the Middle America trench. The atypical influx of water into the Nicaraguan section of the subduction zone ultimately leads to a regional maximum in the degree of mantle melting. In central Costa Rica, subduction input is also unusual in that it includes oceanic crust flavored by the Galapagos plume. Both of these exotic subduction inputs are recognizable in the compositions of magmas erupted along the volcanic front. In addition, Nicaraguan magmas bear a strong chemical imprint from subducting hemipelagic sediments. The high-field-strength-element depletions of magmas from El Salvador through Costa Rica are related to local variations in the depth to the subducting Cocos plate and, therefore, to segmentation of the volcanic front. Minor phases, probably amphibole or rutile, control these variable depletions. Silicic magmas erupted along the volcanic front exhibit the same along-arc geochemical variations as their mafic brethren. This and their mantle-like radiogenic isotopic compositions suggest the production of juvenile continental crust all along the Central American subduction zone. Punctuated times of enhanced magmatic input from the mantle may aid in crustal development.

### SOMMAIRE

L'Amérique centrale a récemment été le lieu de recherches sur les processus complexes se produisant dans les zones de subduction. Ici nous passons en revue certaines découvertes sur nature

des intrants de subduction, la production et l'évolution des magmas, ainsi que les extrants volcaniques résultants. Dans le segment nicaraguayen de la zone de subduction, les intrants de subduction sont exceptionnellement humides, probablement à cause de la serpentinisation généralisée de la portion mantélique de la plaque en subduction, fissurée par flexure dans partie marine de la fosse océanique de l'Amérique centrale. L'afflux atypique en eau dans le segment nicaraguayen de la zone de subduction induit ultimement un maximum régional de la proportion de fusion du manteau. Dans la portion centrale du Costa Rica l'intrant de subduction est lui aussi atypique en ce qu'il comprend une croûte océanique teintée par le panache des Galápagos. Ces deux intrants de subduction atypiques sont répercutés dans la composition des magmas éjectés le long du front volcanique. En outre, les magmas nicaraguayens affichent une forte empreinte chimique héritée des sédiments hémipélagiques en subduction. Les appauvrissements en éléments à fortes liaisons atomiques des magmas, du El Salvador jusqu'au Costa Rica, sont liés à des variations localisées de la profondeur de la plaque en subduction de Cocos, et donc, à la segmentation du front volcanique. Des phases mineures, probablement amphibole et rutile, déterminent ces appauvrissements variables. Les magmas siliceux éjectés le long du même front volcanique montrent les mêmes variations géochimiques le long de l'arc que leur contrepartie mafique. De plus, les compositions radiogéniques de leurs contreparties mantéliques évoquent la


production d'une croûte continentale juvénile le long de la zone de subduction de l'Amérique centrale. Des épisodes d'accroissements ponctuels des intrants magmatiques du manteau peuvent contribuer au développement d'une croûte.

## INTRODUCTION

The Central American subduction zone is awash with volcanologic and petrologic diversity over its 1100 km-long length (McBirney 1969; Carr et al. 1982, 2003, 2007a, b; Carr 1984; van Wyk de Vries et al. 2007). Over the past decade and a half, encouraged in part by relative political stability, this diversity has been examined in finer detail, yielding considerable new and important insights into magmatic and volcanic processes occurring in the Central American subduction zone. Recognizing and understanding the intricacies of this subduction zone is crucial, as all of Central America's population is at risk from earthquakes and volcanic eruptions (Ewert and Harpel 2004; Witham 2005; Dilley et al. 2005; Auker et al. 2013). In this brief review, we will summarize some of the newer insights concerning Central American magmatism and volcanism with the aim of stimulating future research along this fascinating convergent margin. We acknowledge that this review is in no way comprehensive, but it does provide an overall view of the efforts to understand Central American magma genesis. There have also been many recent, significant studies concerning other aspects of the Central American subduction zone and its volcanoes, including neotectonics, eruption dynamics, tephra fallout, volatile emissions, volatile cycling, volcanic stratigraphy, groundwater – volcano interactions, volcanic surveillance, and hazard assessments.

## TECTONIC, GEOLOGIC, AND VOLCANIC FRAMEWORK

In this paper we will restrict the boundaries of the Central American subduction zone (CASZ) to the region that is experiencing subduction of the Cocos Plate beneath the Caribbean Plate, i.e. from the Guatemala – Mexico border to central Costa Rica (Fig.



**Figure 1.** Simplified tectonic framework for Central America. Red triangles are Quaternary volcanic complexes of the volcanic front. Arrows point to discontinuities (segment boundaries) in the volcanic front after Carr et al. (2007b). Arrows also indicate estimated convergence directions (from Syracuse and Abers 2006). Numbers in parentheses are calculated convergence rates also from Syracuse and Abers (2006). CNA is the Caribbean – North American plate boundary.

1). The rate of Cocos – Caribbean convergence slightly increases south-eastward along the trench from about 6 to 9 cm/yr (DeMets 2001; Syracuse and Abers 2006) and slab dips below the active volcanoes are variable, with the steepest dips beneath Nicaragua (Carr 1984; Protti et al. 1995; Syracuse and Abers 2006; Syracuse et al. 2008; MacKenzie et al. 2010). Slab depths are also greatest beneath Nicaragua (Syracuse and Abers 2006). Oceanic crust formed at the East Pacific Rise (ca. 25 Ma), with a normal mid-ocean ridge basalt composition, subducts from Guatemala to northern Costa Rica, whereas 15–20 Ma oceanic crust that formed at the Cocos – Nazca Spreading Center and was overprinted by Galapagos hotspot tracks, subducts in central Costa Rica (Werner et al. 1999; O'Connor et al. 2007). There is no evidence of sediment accretion along the CASZ, so all sediments are assumed to be subducted into the mantle (Aubouin et al. 1984; Ranero and von Huene 2000; Moritz et al. 2000). The total thickness of subduct-

ing sediments is similar all along the CASZ, as is their lithologic architecture, which consists of an overlying sequence of hemipelagic muds underlain by carbonate oozes (von Huene et al. 1980; Plank and Langmuir 1993, 1998; Kimura et al. 1997; Patino et al. 2000). The presence of the North American – Caribbean plate boundary transecting Guatemala (Fig. 1) adds an appreciable level of complexity to the tectonics of the northern CASZ (Burkart and Self 1985; Guzmán-Speziale 2001; Lyon-Caen et al. 2006; Rogers and Mann 2007; Álvarez-Gómez et al. 2008; Rodríguez et al. 2009; Walker et al. 2011).


Although Cocos – Caribbean convergence has a much longer history (Mann et al. 2007; Gazel et al. 2009, 2011; Alvarado and Gans 2012), in this review we will restrict discussion to Quaternary volcanism, focusing on the volcanic front where volcanism has been overwhelmingly concentrated during the Quaternary (Carr et al. 1982, 2003, 2007a). Crustal thickness below the volcanic front is shallowest

in Nicaragua (ca. 25–30 km) and thickens to ca. 35–45 km toward the extremities of the CASZ (Carr 1984; Carr et al. 1990, 2003; Sallarès et al. 2001; MacKenzie et al. 2010; Lücke et al. 2010). According to Rogers et al. (2007), much of the volcanic front, from southeastern Guatemala through Nicaragua, has been built on the southern Chortis terrane, which is floored by a post-Paleozoic arc-type or ocean-crust basement (Geldmacher et al. 2008). The northwestern portion of the volcanic front in central and northwestern Guatemala, by contrast, is likely underlain, at least in part, by rocks of the central Chortis terrane, which has a continental Paleozoic to Precambrian substrate (Dengo 1985; Rogers et al. 2007). The crustal character below the other end of the volcanic front, in Costa Rica, is also distinctive, being thickly anchored by rocks of the Caribbean Large Igneous Province, interpreted as a product of the Galapagos plume head (Hauff et al. 1997, 2000; Sallarès et al. 2001).

Magmas erupted along the volcanic front range from basalts to rhyolites that largely exhibit the characteristic elemental signatures of a subduction zone origin (Carr et al. 1982, 2003, 2007b; Carr 1984; Walker 1989; Leeman et al. 1994; Patino et al. 2000; Sadofsky et al. 2008). Along most of the CASZ there is a bimodal distribution of compositions with peaks in basalt – basaltic andesite and rhyolite (Vogel et al. 2006).

### THE VOLCANIC FRONT – ARE ALL SEGMENTS EQUALLY CREATED?

The Central American volcanic front is thought to be among the world's most active volcanic belts (Bluth and Rose 2002). The application of high precision  $^{40}\text{Ar}/^{39}\text{Ar}$  dating to a larger number of volcanic rocks associated with Quaternary volcanism in Central America (Rose et al. 1999; Vogel et al. 2004; Carr et al. 2007a; Escobar-Wolf et al. 2010; Singer et al. 2011; Alvarado and Gans 2012) has allowed renewed appraisals of extrusive and magmatic fluxes along the volcanic front. Carr et al. (2007a) estimate that extrusion rates along the Nicaraguan and Costa Rican segments of the front are roughly



**Figure 2.** a. Estimates of magma flux in four portions of the Central American volcanic front from Kutterolf et al. (2008b). b. Estimates of the volumes of volcanic centers for same four portions of the Central American volcanic front from Bolge et al. (2009).

equivalent within error. In contrast, Bolge et al. (2009) calculate raw eruptive volumes along the volcanic front in Costa Rica that are substantially larger than those in Nicaragua, consistent with earlier inferences that Costa Rica has had a greater volcanic flux than Nicaragua (Carr 1984; Carr et al. 1990). For their volume estimates, Bolge et al. (2009) incorporate important new data from Kutterolf et al. (2008b) on tephra erupted along the CASZ over the past 322,000 years. According to Kutterolf et al. (2008b), these Plinian

products account for approximately 65% of the total magmatic output in the CASZ. Moreover, their analysis indicates that the overall magma fluxes have been greatest in the northern half of the CASZ (Kutterolf et al. 2008b; Fig. 2a). If, however, the volume estimates of Bolge et al. (2009) are taken as a simple reflection of overall magmatic flux, then Costa Rica emerges as the most magmatically productive portion of the CASZ (Fig. 2b).

All of the investigations summarized in this section so far have



been regional in scope. Singer et al. (2011) instead focus on a single volcano – Santa María in Guatemala – and employ an extensive collection of  $^{40}\text{Ar}/^{39}\text{Ar}$  age determinations, mostly from Escobar-Wolf et al. (2010). Their results indicate that this prototypical composite cone was constructed in four stages spanning approximately 75 kyr. The magma flux calculated by Singer et al. (2011) for building of the Santa María cone, based in part by energy-constrained recharge, assimilation, and fractional crystallization (EC–RAXFC) modeling (Bohrson and Spera 2007), is impressive – ca. 46 660 g/s, about 4 times the rate estimated by Kutterolf et al. (2008b). Moreover, Singer et al. (2011) point out that this magmatic input is also several times greater than those assumed in thermal models of the interaction between mantle-derived melts and the crust (Dufek and Bergantz 2005; Annen et al. 2006). Singer et al. (2011) also estimate a regional extrusion flux for Guatemala during cone construction at Santa María of  $5\text{--}9\text{ km}^3\text{km}^{-1}\text{Myr}^{-1}$ , which is comparable to those reported by Carr et al. (2007a) for Nicaragua and Costa Rica.

## NICARAGUA – WET INPUT AND OUTPUT?

Ranero et al. (2001) first report the occurrence of extensive bending-related faulting in the incoming Cocos Plate at the outer rise offshore Nicaragua. This structural control would lead to deep and widespread serpentinization of the mantle portion of the approaching plate, and hence an anomalously wet subduction input (Rüpke et al. 2002; Ranero et al. 2003). Subsequent geophysical investigations off Nicaragua lend support to this hypothesis (Grevemeyer et al. 2005, 2007; Ivandic et al. 2008, 2010; Lefeldt et al. 2009; Key et al. 2012). Grevemeyer et al. (2005) document lower than expected heat flow in the subducting Cocos Plate, which they attribute to enhanced hydrothermal circulation associated with bending-related faulting. A series of studies document slow seismic velocities in the outer rise offshore Nicaragua, suggesting

15–30% serpentinization of Cocos Plate mantle to at least a few kilometres below the Moho (Grevemeyer et al. 2007; Ivandic et al. 2008, 2010; Van Avendonk et al. 2011). Van Avendonk et al. (2011) estimate that the incoming Cocos slab is about 2.5 times wetter offshore Nicaragua versus offshore Costa Rica, and like all of the previous geophysical investigations, attribute this to widespread serpentinization of the mantle portion of the incoming Cocos Plate.


Onshore, Abers et al. (2003) find unusually slow seismic velocities at the top of subducting Cocos Plate beneath Nicaragua, suggesting an anomalously wet slab. In a more detailed study, Syracuse et al. (2008) also report anomalously low seismic velocities in the upper part of the subducting Cocos plate, extending 20–30 km below the slab surface, i.e. well into the mantle lithosphere. The observed velocity anomalies are consistent with 10–20% serpentinization of the subducting Cocos mantle. Therefore, compelling evidence exists on both sides of the trench for an uncommonly wet input into the Nicaraguan portion of the CASZ.

The consequences of this unusual hydrous influx have also been seismically imaged. In the Nicaraguan mantle wedge, Syracuse et al. (2008) distinguish a vertically extensive region having high ratios of primary to secondary wave velocities ( $V_P/V_S$ ), thought to outline melt generated by the large water additions from the slab. Rychert et al. (2008), in a related investigation, show that the Nicaraguan mantle wedge exhibits a relatively wide zone of high shear wave attenuation that would indicate considerable wedge melting, hydration, excess temperatures, or some combination of all three. Hence, both studies provide convincing evidence for unusually large amounts of melting in the Nicaraguan mantle wedge that can be linked to an anomalously wet, serpentinized subducting plate.

There are a number of geochemical indicators that support the geophysical picture of an arc segment with enhanced melt production linked to an amplified delivery of water from

the subducting Cocos plate. The first is that Nicaraguan mafic magmas and olivine-hosted melt inclusions define a regional minimum in La/Yb along the CASZ (Fig. 3a; Carr et al. 1990, 2003, 2007b; Sadofsky et al. 2008; Bolge et al. 2009). Carr et al. (1990, 2003, 2007b) attribute this to higher degrees of wedge melting which, in turn, implies a greater hydrous flux from the subducted slab, since flux-melting is generally thought to be the predominant means of magma production in subduction zones (Ringwood 1974; Ulmer 2001; Wallace 2005; Grove et al. 2012). Higher degrees of wedge melting in Nicaragua are also consistent with the lower  $\text{Na}_2\text{O}$  contents of mafic magmas and olivine-hosted melt inclusions erupted in Nicaragua compared with those emitted elsewhere along the CASZ (Carr 1984; Plank and Langmuir 1988; Eiler et al. 2005; Syracuse and Abers 2006; Carr et al. 2007b; Sadofsky et al. 2008). Sadofsky et al. (2008) conclude that (western) Nicaragua has a wetter mantle wedge and erupts somewhat wetter magmas than Guatemala and Costa Rica, based on water analyses of olivine-hosted melt inclusions (they had no data for El Salvador). Finally, Eiler et al. (2005) find that olivine phenocrysts from Nicaraguan lavas have unusually low  $\delta^{18}\text{O}$ , both for the CASZ and for subduction zones worldwide (Fig. 3b). The Nicaraguan olivines also have anomalously low  $\delta^{18}\text{O}$  relative to normal mid-ocean ridge basalts and mantle peridotites (Fig. 3b). These distinctively low values are ascribed to an aqueous fluid component from hydrothermally altered rocks deep within the subducting Cocos plate, possibly subducted serpentinites (Eiler et al. 2005). According to the modeling results of Eiler et al. (2005), this hydrous component has a much reduced to non-existent influence elsewhere along the CASZ.

Other geochemical proxies of water involvement in Nicaraguan magma genesis are more problematic or fail to single out Nicaraguan magmas. The first is the somewhat iconic regional peak in Ba/La exhibited by (western) Nicaraguan mafic lavas (Fig. 4a). Because a number of experimen-




**Figure 3.** Variations of La/Yb (a) and  $\delta^{18}\text{O}$  (b) in mafic volcanic rocks (45–55 wt%  $\text{SiO}_2$ ) erupted along the Central American volcanic front. Data from RU\_CAGeochem2013 database (<http://rci.rutgers.edu/~carr/>) supplemented with data from La Femina et al. (2004) and Singer et al. (2011). Shaded band represents  $\delta^{18}\text{O}$  for most mantle peridotites and normal mid-ocean ridge basalts (Eiler et al. 2005).

tal studies have demonstrated that Ba is mobile during slab dehydration, whereas La is not (Tatsumi et al. 1986; Keppler 1996; Martin et al. 2011), Ba/La, like Ba/Nb and Ba/Th, is com-

monly employed as a proxy for a slab fluid component (Woodhead and Johnson 1993; Woodhead et al. 1998; Walker et al. 2000; Sadofsky et al. 2008; Bolge et al. 2009). However, the

regional variation of Ba/La in mafic output in the CASZ actually reflects regional variability in La, not Ba (Carr et al. 1990, 2007a), as may be the case on a global scale (Morris and Hart 1983), and thus is more difficult to directly relate to a variable fluid flux. In addition, as first pointed out by Carr et al. (1990), and subsequently highlighted by others (Leeman et al. 1994; Jicha et al. 2010), the regional peak in Ba/La corresponds with a regional peak in a less ambiguous geochemical proxy,  $^{10}\text{Be}$  (Fig. 4b).  $^{10}\text{Be}$  is a clear tracer of the recycling of young subducting sediment (Tera et al. 1986; Morris et al. 1990), which in Central America would pinpoint source contributions from the upper sequence of hemipelagic sediments (Leeman et al. 1994; Patino et al. 2000). Since subducting sediments in the CASZ are distinctly enriched in Ba (Plank and Langmuir 1993; Leeman et al. 1994; Patino et al. 2000) and Ba/La (Patino et al. 2000), the Nicaraguan peak in Ba/La most likely reflects an increased slab signal from subducted (hemipelagic) sediments (Jicha et al. 2010). An enhanced flux from subducted (hemipelagic) sediments can also explain the along-arc peaks in B/La, U/Th, Ba/Th, and ( $^{230}\text{Th}/^{232}\text{Th}$ ) seen in Nicaragua (Leeman et al. 1994; Patino et al. 2000; Carr et al. 2003; Walker et al. 2007; Sadofsky et al. 2008; Jicha et al. 2010). The He –  $\text{CO}_2$  relationships and the  $\delta^{15}\text{N}$  ratios of volcanic and geothermal fluids in Nicaragua also suggest elevated contributions from subducted sediments (Shaw et al. 2003; Elkins et al. 2006), although  $\delta^{15}\text{N}$  ratios in Nicaragua are similar to those in Guatemala (Fischer et al. 2002; Elkins et al. 2006). Jicha et al. (2010) point out that  $^{238}\text{U}$  excesses do not peak in Nicaragua, as might be expected if fluid input is maximized in this segment of the CASZ. Although  $^{238}\text{U}$  excess is considered to be a robust indicator of the addition of  $\text{U}^{+6}$  as part of a fluid component from subducting lithosphere (Gill 1981; Allègre and Condomines 1982; Turner et al. 2003), it is a function of the fractionation of U from Th in aqueous fluids and not necessarily a measure of the overall water flux. The fact that  $^{238}\text{U}$  excesses



**Figure 4.** Variations of Ba/La (a) and  $^{10}\text{Be}/^9\text{Be}$  (b) in mafic volcanic rocks (45–55 wt %  $\text{SiO}_2$ ) erupted along the Central American volcanic front. Sources of data as in Figure 3.

extend from Guatemala to Costa Rica is, however, an important observation as it indicates that a fluid component of some sort is involved in magma genesis all along the CASZ.

An additional problem with the idea that wedge melting is maxi-

mized beneath the Nicaraguan portion of the CASZ is that, no matter the means of estimation, extrusive fluxes in Nicaragua do not stand out, as summarized above. In other words, the logical expectation is that with more melting there would be greater surface

volcanism, particularly given that Nicaragua has the thinnest crust in the CASZ (Carr 1984; Carr et al. 1990, 2003). This discrepancy was first addressed by Carr et al. (1990), who attribute it to the steeper slab dip beneath Nicaragua. With a steep dip, water influx from the subducted Cocos plate is focused into a tighter volume, producing smaller quantities of high percentage partial melts that possess a more concentrated slab signal (Carr et al. 1990, 2003; Feigenson and Carr 1993).

We think a coupling of the models by Carr et al. (1990) and Eiler et al. (2005) best links the compelling evidence for an unusually wet input into the Nicaraguan segment of the CASZ and its equally distinctive volcanic output. The model of Carr et al. (1990) reconciles a regionally large degree of melting in the mantle wedge with a non-distinctive volcanic output. The enhanced degree of melting in Nicaragua is directly related to an anomalously wet input from the subducting Cocos plate, likely from serpentinites formed in the outer rise (Ranero et al. 2001, 2003; Rüpke et al. 2002; Eiler et al. 2005). In the model of Eiler et al. (2005), this wet addition is provided by their water-rich, low  $\delta^{18}\text{O}$  component. As demonstrated by Eiler et al. (2005), a second slab component, required in Nicaragua to provide the necessary enhancements of  $^{10}\text{Be}$ , Ba, B, U and other trace elements, must be subducted hemipelagic sediments (Morris et al. 1990; Plank and Langmuir 1993; Leeman et al. 1994; Patino et al. 2000; Rüpke et al. 2002; Shaw et al. 2003; Eiler et al. 2005; Elkins et al. 2006; Jicha et al. 2010). Eiler et al. (2005), following the current consensus (e.g. Elliott 2003), suggests that this second component is a sediment melt, although they provide it with very un-sedimentary Sr and Nd isotopic compositions. Leeman et al. (1994), on the other hand, favor addition of the sediment signal via fluid transport. The agent transporting the sedimentary component in Nicaragua remains an open question and may hinge on the relative mobilities of trace elements, particularly Be, Sr, Nd, and Hf, in various slab fluids (Tatsumi and




Isoyama 1988; You et al. 1994, 1996; Johnson and Plank 1999; Woodhead et al. 2001; Eiler et al. 2005; Marschall et al. 2007).


### CENTRAL COSTA RICA – INPUT FROM THE GALAPAGOS PLUME

As shown in Figure 3a, volcanic rocks erupted in central Costa Rica have notably elevated La/Yb (Carr et al. 1990, 2003, 2007b; Herrstrom et al. 1995). Higher La/Yb implies a lower degree of partial melting or derivation from an enriched magma source. As shown in Figure 5, central Costa Rican volcanic rocks also have unusually enriched Pb isotopic compositions indicative of an enriched mantle source (Feigenson et al. 2004; Hoernle et al. 2008; Gazel et al. 2009, 2011). The overall correlation between La/Yb and Pb isotope ratios indicates that enriched magma sources are present where La/Yb is >10 (Gazel et al. 2009, 2011). The low Zr/Nb in central Costa Rican lavas is also supportive of an enriched mantle source (Bolge et al. 2009).

The observed source enrichment in central Costa Rica has been the subject of much debate. However, most studies have linked source enrichment to the influence of the Galapagos plume (Johnston and Thorkelson 1997; Abratis and Wörner 2001; Feigenson et al. 2004; Goss and Kay 2006; Hoernle et al. 2008; Gazel et al. 2009, 2011), which has had a fundamental role in the history of both the Caribbean and Cocos plates (Sinton et al. 1998; Werner et al. 1999, 2003; Hauff et al. 2000; Barckhausen et al. 2001; Hoernle et al. 2002; Denyer and Gazel 2009). Recall that the oceanic crust that subducts beneath central Costa Rica was formed at the Cocos – Nazca spreading center, and then overprinted by Galapagos hotspot tracks (Werner et al. 1999; O'Connor et al. 2007). In detail, the subducting Galapagos Seamount Province outboard of central Costa Rica has an alkaline composition and an isotopic signature of the Northern Galapagos Domain (Wolf – Darwin Lineament in the Galapagos Archipelago; Hoernle et al. 2000; Werner et al. 2003; Fig. 6). The subducting Cocos and Coiba ridges



**Figure 5.** Variation of  $^{206}\text{Pb}/^{204}\text{Pb}$  in mafic volcanic rocks (45–55 wt%  $\text{SiO}_2$ ) erupted along the Central American volcanic front. Sources of data as in Figure 3.




**Figure 6.** Tectonic setting of southern Central America (modified from Gazel et al. 2011). Black circles are volcanic complexes of the volcanic front. Orange hexagons are locations of adakitic volcanism. EPR: East Pacific Rise; CNS: Cocos-Nazca Spreading Center; C.I.: Cocos Island; PFZ: Panama Fracture Zone. The depth contours of the subducting Cocos slab are from Protti et al. (1994).



have a tholeiitic composition with a dominant isotopic composition belonging to the central Galapagos Domain (Fernandina Island; Hoernle et al. 2000; Werner et al. 2003; Fig. 6).

The Pb isotopic compositions of samples from central Costa Rica can be explained by three isotopic end-members, a depleted component (depleted mantle) and two enriched subducting Galapagos components, the Seamount Province and the Cocos – Coiba Ridge (Fig. 7; Hoernle et al. 2008; Gazel et al. 2009). The Seamount Province is a recently arrived (<7 Ma) component; before that, the Galapagos interaction was dominated by a component similar to the Coiba and Cocos ridges that arrived at the subduction system ca. 15–10 Ma (Gazel et al. 2011). Based on radiogenic isotope systematics, geochemical variations with time, and geochemical modeling, Gazel et al. (2009) propose that the process to produce magmas with a Galapagos signature requires partial melting of subducting Galapagos tracks and reaction of those melts with the mantle wedge. These conclusions are in agreement with other recent studies (Benjamin et al. 2007; Hoernle et al. 2008) that provide convincing evidence that the anomalous enriched signature in the central Costa Rican portion of the volcanic front is derived from the interaction of the mantle wedge with Galapagos hotspot tracks. Hoernle et al. (2008) suggest that trench-parallel mantle flow, perhaps coupled with oblique subduction, causes diffusion of this Galapagos signature from central Costa Rica northwestward into Nicaragua. Trench-parallel mantle flow is consistent with recent seismic anisotropy data for the southern portion of the CASZ (Hoernle et al. 2008; Abt et al. 2009, 2010).

Gazel et al. (2009, 2011) show that the appearance of the Galapagos signature in the central Costa Rican volcanic front correlates with the production of magmas having primitive andesitic/adakitic compositions. Adakites are intermediate magmas produced by high-pressure melting of a mafic protolith, such as a subducted slab, and are recognized in part by their high La/Yb and Sr compositions (Kay



**Figure 7.**  $^{208}\text{Pb}/^{204}\text{Pb}$  versus  $^{206}\text{Pb}/^{204}\text{Pb}$  for Costa Rican lavas showing mixing lines from the three proposed source components: depleted mantle (DM), the subducting seamount province (SP), and the subducting Cocos – Coiba ridge (CCR). Sources of data as in Figure 3.

1978; Defant and Drummond 1990; Gazel et al. 2011; Whattam et al. 2012). Their presence in central Costa Rica strongly supports the occurrence of slab melting and its importance in explaining the enriched signature seen in southernmost CASZ magmas. Moreover, Gazel et al. (2011) report a migration of adakitic lavas of 35 mm/y towards the southeast, tracking the eastward movement of the triple junction where the Panama Fracture Zone intersects the Middle America Trench (Fig. 6). Seismic evidence (Protti et al. 1994) suggests that there is a slab ‘window’ beneath southern Costa Rica and Panama, in the area where adakites are common (orange hexagons in Fig. 6). The numerous hotspot tracks and fracture zones on the subducting Cocos and Nazca plates (Werner et al. 1999) could make the subducting slab below this part of the CASZ relatively easy to tear. Therefore, Gazel et al. (2011) hypothesize that a collision between the Galapagos hotspot tracks (Coiba Ridge?) and the CASZ that started ca. 15–10 Ma (Denyer and Arias 1991; Silver et al. 2004; MacMillan et al. 2004; Gazel et al. 2009) clogged the subduction zone and triggered slab detachment below


southern Costa Rica and Panama. The detached slab segments were then replaced by hot asthenosphere, which is consistent with the elevated mantle potential temperatures (1400–1450 °C) in the mantle wedge below southern Central America (Gazel et al. 2011). The slab-free area correlates with the highest elevations (i.e. the Talamanca Cordillera at ca. 4 km) in southern Central America. These surface elevations are possibly related to the isostatic effect of the influx of hot mantle together with shortening related to the collision of the Cocos Ridge (Gazel et al. 2011).

## VOLCANIC SEGMENTATION AND HIGH-FIELD-STRENGTH-ELEMENTS


Stoiber and Carr (1973), Carr et al. (1982, 2007b), and Carr (1984), building on observations made by early explorers of Central America (Dollfus and Montserrat 1868; Sapper 1917), divided the Central American volcanic front into seven or eight segments, each from 100 to 300 km long, separated by changes of strike, ‘volcanic gaps’, or right-hand step-outs (Fig. 1). These along-front discontinuities are generally associated with transverse structures, such as faults and align-

ments of volcanic vents (Stoiber and Carr 1973). Although originally thought to reflect severance of both the overriding and subducting plates (Stoiber and Carr 1973; Carr et al. 1982), segmentation is now thought to be solely an upper plate phenomenon (Burkart and Self 1985; Bolge et al. 2009). Bolge et al. (2009) show that Zr/Nb correlates with the well-recognized segmentation of the Central American volcanic front; specifically, Zr/Nb in erupted, high-field-strength-element (HFSE)-depleted mafic magmas declines abruptly on the north-western side of four proposed segment boundaries in El Salvador through Costa Rica (Fig. 8; Bolge et al. 2009). Bolge et al. (2009) demonstrate that these variations are controlled by changes in Nb, not Zr. All of the observed discontinuities in Zr/Nb correspond with right-hand steps in the volcanic front (Bolge et al. 2009). As a result, the variations in Zr/Nb are closely mimicked by sharp changes in slab depth, which abruptly decreases on the southeastern side of the segment boundaries, i.e. with each right-hand (trenchward) step (Fig. 1; Syracuse and Abers 2006; Bolge et al. 2009). Thus, at each right-hand step, slab depth decreases and Zr/Nb increases, caused by increasing Nb depletion in the erupted magmas (Bolge et al. 2009). It is important to point out that the discontinuous variations in Zr/Nb are superimposed on an overall along-arc trend in which western Nicaraguan lavas define a weak regional peak in Zr/Nb (Fig. 8; Bolge et al. 2009), grossly analogous to the along-arc variations in Ba/La and  $^{10}\text{Be}$  (Fig. 4). Although not examined by Bolge et al. (2009), Hf/Ta variations in the CASZ are identical to Zr/Nb as is evident from the amazingly good correlation between the two incompatible element ratios (Fig. 9).

Bolge et al. (2009) speculate that the segmented Zr/Nb changes along the CASZ are controlled by variable amphibole stability during melting of the subducted Cocos plate. At shallower slab depths, residual amphibole is present during slab melting, resulting in high Zr/Nb if  $^{Amph/L}D_{Nb/Zr} > 1$  (e.g. Tiepolo et al. 2001). At greater slab



**Figure 8.** Variation of Zr/Nb in low-Ti mafic volcanic rocks (45–55 wt%  $\text{SiO}_2$ ) erupted along the Central American volcanic front from El Salvador through central Costa Rica. Vertical lines show positions of proposed segment boundaries (see Fig. 1). Sources of data as in Figure 3.



**Figure 9.** Zr/Nb versus Hf/Ta in low-Ti mafic volcanic rocks (45–55 wt%  $\text{SiO}_2$ ) erupted along the Central American volcanic front from El Salvador through central Costa Rica. Sources of data as in Figure 3.


depths, amphibole would not be stable during melting (e.g. Niida and Green 1999) resulting in low Zr/Nb (Bolge et al. 2009). Although amphibole control cannot presently be ruled out, we think

rutile is a more likely option for directing relative HFSE fractionation along the CASZ. Rutile, unlike other possible accessory phases such as titanite, is well known for its ability to decouple

Nb and Ta from Zr and Hf (Jenner et al. 1993; Foley et al. 2000; Klemme et al. 2005; Xiong et al. 2005; Bromiley and Redfern 2008), which is the critical necessity along the CASZ. Rutile would also be a likely accessory phase in subducting siliceous sediments (Hermann and Spandler 2008; Skora and Blundy 2010) which, as discussed above, clearly exert a large influence on the trace element budget in Nicaragua and may do so further north along the CASZ as well. Bolge et al. (2009) dismissed rutile control for two reasons: first, because of its inability to explain a negative correlation between Zr/Nb and Nb/Ta; however, this negative correlation is seen only in El Salvador and there only moderately ( $r^2 = 0.694$ ). Second, the stability of rutile is not pressure dependent; but pressure dependence is not essential, as the stability of the guiding mineral could be a stronger function of temperature, or in the case of partial melting, the degree of melting. For instance, at shallow slab depths and lower slab surface temperatures, rutile would likely be stable during melting or dehydration, resulting in high Zr/Nb and Hf/Ta. At higher temperatures, deeper along the slab surface, rutile might not be stable (Skora and Blundy 2010) or might become exhausted during melting, producing lower Zr/Nb and Hf/Ta in generated magmas.

#### SILICIC VOLCANISM – GENERATION OF JUVENILE CONTINENTAL CRUST?

Kutterolf et al. (2008b) have shown that silicic volcanism is volumetrically significant along the CASZ and may in fact dominate the overall magmatic output over the past few hundred thousand years. However, geochemical investigations of Central American silicic rocks are still too few and far between. One of the most important recent studies is that by Vogel et al. (2006), which expands on thoughts first presented in Vogel et al. (2004). In both papers, Vogel et al. (2004, 2006) stress that erupted felsic magmas along the Central American volcanic front display many of the same regional geochemical variations as their mafic brethren. For instance, the Ba/La ratios of silicic magmas, with two




**Figure 10.** Variations in Ba/La (a) and La/Yb (b) in silicic volcanic rocks (65–77 wt% SiO<sub>2</sub>) erupted along the Central American volcanic front. Data from Vogel et al. (2006) supplemented by data from Kutterolf et al. (2008a), Garrison et al. (2012), and the RU\_CAGeochem2013 database.

exceptions, peak in Nicaragua and fall off to the northwest and southeast (Fig. 10a). The regional pattern for La/Yb ratios is, on the other hand, the almost mirror image of Ba/La, exhibiting a regional minimum in Nicaragua (Fig. 10b) – in fact, a more pronounced regional minimum than

that shown by mafic magmas (Fig. 3a). In addition, the few available radiogenic isotopic compositions for silicic rocks overlap those of contiguous mafic rocks (Fig. 11). Thus, the silicic magmas of Central America seem to have had little, if any, crustal inheritance, no matter the thickness or com-





**Figure 11.** Nd isotopic compositions of mafic (a) and silicic (b) volcanic rocks (as delineated in previous figures) erupted along the Central American volcanic front. Colour distinctions as in previous diagrams, i.e. red – Guatemala, blue – El Salvador, green – Nicaragua, and purple – Costa Rica. Data from RU\_CAGeochem2013 database, Vogel et al. (2006), Singer et al. (2011), and Garrison et al. (2012).

position of crust they inhabited and traversed (Vogel et al. 2006). This would suggest that juvenile continental crust is being created all along the CASZ (Vogel et al. 2004; Deering et al. 2012). The preferred model of Vogel et al. (2006) for the origin of Central American silicic magmas is by penecontemporaneous partial melting of recently crystallized mafic intrusions (Tamura and Tatsumi 2002), or by melt extraction from more mafic crystal mushes (Deering et al. 2012).

Nevertheless, partial melting of subducting lithosphere may also play a significant role in the generation of continental crust (Drummond and Defant 1990; Rapp and Watson 1995; Hacker et al. 2011). This process would be facilitated if the slab were enriched in incompatible-elements by plume interaction as in the case of Costa Rica (Gazel et al. 2009, 2011). Partial melting of subducting hotspot tracks can ‘re-fertilize’ the arc mantle wedge. Once the enriched starting material is produced, intra-crustal processes such as fractional crystallization, assimilation and anatexis (e.g. Hildreth and Moorbath 1988; Annen et al. 2006) will complete the development of juvenile continental crust. This process, which may be occurring on a large scale in Costa Rica, is consistent with a recent geophysical study (Hayes et al. 2013) showing that Costa Rica has average *P*-wave velocities that are closest to continental crust of any non-continental subduction zone, worldwide.

#### DIRECTIONS AND QUESTIONS FOR THE FUTURE

A number of important avenues for future research in the CASZ fall out of this review. The first is further applications of high-precision  $^{40}\text{Ar}/^{39}\text{Ar}$  dating to individual volcanoes in order to better constrain extrusive and magmatic fluxes along the CASZ (e.g. Singer et al. 2011). A second is to examine whether the sediment signature seen in Nicaraguan volcanic rocks is delivered by a fluid or a melt from the subducting Cocos plate. Acquisition of Hf isotopic data could shed light on this question, depending on whether the Hf isotopic composition of the CASZ

mantle wedge is distinct from that of the subducting Cocos crust (Tollstrup et al. 2010). Definitive geochemical evidence of a serpentinite-derived component in Nicaragua is also lacking: chlorine isotope values for volcanic rocks, like those of oxygen isotopes, are only suggestive of a serpentinite signature (Barnes et al. 2009). Tonarini et al. (2007) utilize B isotopes to argue for fluid inputs in El Salvador; a similar B isotopic study of Nicaraguan volcanic rocks is needed to validate the approach of Tonarini et al. (2007) in the segment of the CASZ with, at present, the strongest geophysical evidence for serpentinite subduction (e.g. Van Avendonk et al. 2011) and the most robust geochemical evidence for a wet slab input. Another important research question is the character of fluid addition from the subducting Cocos plate in portions of the CASZ away from Nicaragua, as U-series data suggest fluid involvement all along the CASZ, even in Costa Rica (Benjamin et al. 2007; Jicha et al. 2010). Future investigations of HFSE variability along the Central American volcanic front are also vital to identify which minor or accessory phases (or phase) control(s) the observed variation and how it relates to CASZ segmentation and slab depth (Bolge et al. 2009). One aspect of HFSE variability not discussed in this review is the occurrence of unusual HFSE-enriched volcanic rocks along the volcanic front, particularly in Nicaragua and Costa Rica (Ui 1972; Walker 1984; Reagan and Gill 1989; Walker et al. 1990, 2001; Alvarado and Carr 1993; Feigenson and Carr 1993; Reagan et al. 1994; Patino et al. 2000; Carr et al. 2003, 2007b; Sadofsky et al. 2008; Freundt et al. 2010; Rausch and Schmincke 2010). The origin of these atypical volcanic rocks is still controversial and wants further study, bearing in mind that the Nicaraguan and Costa Rican examples are geochemically quite distinct (Walker et al. 1990, 2001). Another research focus for the CASZ in the future should be the origin and evolution of silicic volcanism, particularly within large caldera systems. The Vogel et al. (2006) hypothesis of important juvenile crustal production

all along the volcanic front requires geochemical testing, especially with radiogenic isotope data. Last, and perhaps most important, both onshore and offshore geophysical studies of northern Central America are imperative. These would provide an important foundation for vital assessments of the geochemical variations along the northern half of the Central American volcanic front, which are sometimes as robust as those in the southern half (Figs. 3b and 4), but have attracted scant scientific attention. Readers interested in examining the geochemistry of CASZ volcanic rocks further are encouraged to download the RU\_CAGeochem database from <http://rci.rutgers.edu/~carr/>.

### ACKNOWLEDGEMENTS

Many thanks to Jarda Dostal for the invitation to review aspects of magmatic/volcanic activity in the Central American subduction zone. We also thank Mike Carr and an anonymous reviewer for their beneficial reviews of the manuscript. Mark Howland, cartographer extraordinaire, drafted all of the figures.

### REFERENCES

- Abers, G.A., Plank, T., and Hacker, B.R., 2003, The wet Nicaraguan slab: *Geophysical Research Letters*, v. 30, 1098, <http://dx.doi.org/10.1029/2002GL015649>.
- Abratis, M., and Wörner, G., 2001, Ridge collision, slab-window formation, and the flux of Pacific asthenosphere into the Caribbean realm: *Geology*, v. 29, p. 127–130, [http://dx.doi.org/10.1130/0091-7613\(2001\)029<0127:RCSWFA>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2001)029<0127:RCSWFA>2.0.CO;2).
- Abt, D.L., Fischer, K.M., Abers, G.A., Strauch, W., Protti, J.M., and González, V., 2009, Shear wave anisotropy beneath Nicaragua and Costa Rica: Implications for flow in the mantle wedge: *Geochemistry, Geophysics, Geosystems*, v. 10, Q05S15, <http://dx.doi.org/10.1029/2009GC002375>.
- Abt, D.L., Fischer, K.M., Abers, G.A., Protti, M., González, V., and Strauch, W., 2010, Constraints on upper mantle anisotropy surrounding the Cocos slab from SK(K)S splitting: *Journal of Geophysical Research*, v. 115, B06316, <http://dx.doi.org/10.1029/2009JB006710>.
- Allègre, C.J., and Condomines, M., 1982, Basalt genesis and mantle structure studied through Th-isotopic geochemistry: *Nature*, v. 299, p. 21–24, <http://dx.doi.org/10.1038/299021a0>.
- Alvarado, G.E., and Carr, M.J., 1993, The Platanar–Agua Zarcas volcanic centers, Costa Rica: spatial-temporal association of Quaternary calc-alkaline and alkaline volcanism: *Bulletin of Volcanology*, v. 55, p. 443–453, <http://dx.doi.org/10.1007/BF00302004>.
- Alvarado, G.E., and Gans, P.B., 2012, Síntesis geocronológica del magmatismo, metamorfismo y metalogenia de Costa Rica, América Central: *Revista Geológica de América Central*, v. 46, p. 7–122.
- Álvarez-Gómez, J.A., Meijer, P.T., Martínez-Díaz, J.J., and Capote, R., 2008, Constraints from finite element modeling on the active tectonics of northern Central America and the Middle America trench: *Tectonics*, v. 27, TC1008, <http://dx.doi.org/10.1029/2007TC002162>.
- Annen, C., Blundy, J.D., and Sparks, R.S.J., 2006, The genesis of intermediate and silicic magmas in deep crustal hot zones: *Journal of Petrology*, v. 47, p. 505–539, <http://dx.doi.org/10.1093/petrology/egi084>.
- Aubouin, J., Bourgois, J., and Azéma, J., 1984, A new type of active margin: the convergent-extensional margin, as exemplified by the Middle America Trench off Guatemala: *Earth and Planetary Science Letters*, v. 67, p. 211–218, [http://dx.doi.org/10.1016/0012-821X\(84\)90116-X](http://dx.doi.org/10.1016/0012-821X(84)90116-X).
- Auker, M.R., Sparks, R.S.J., Siebert, L., Crossweller, H.S., and Ewert, J., 2013, A statistical analysis of the global historical volcanic fatalities record: *Journal of Applied Volcanology*, v. 2, 24 p., <http://dx.doi.org/10.1186/2191-5040-2-2>.
- Barckhausen, U., Ranero, C.R., von Huene, R., Cande, S.C., and Roeser, H.A., 2001, Revised tectonic boundaries in the Cocos Plate off Costa Rica: Implications for the segmentation of the convergent margin and for plate tectonic models: *Journal of Geophysical Research*, v. 106, p. 19207–19220, <http://dx.doi.org/10.1029/2001JB000238>.
- Barnes, J.D., Sharp, Z.D., Fischer, T.P., Hilton, D.R., and Carr, M.J., 2009, Chlorine isotope variations along the Central American volcanic front and back arc: *Geochemistry, Geophysics, Geosystems*, v. 10, Q11S17,

- <http://dx.doi.org/10.1029/2009GC002587>.
- Benjamin, E.R., Plank, T., Wade, J.A., Kelley, K.A., Hauri, E.H., and Alvarado, G.E., 2007, High water contents in basaltic magmas from Irazú Volcano, Costa Rica: *Journal of Volcanology and Geothermal Research*, v. 168, p. 68–92, <http://dx.doi.org/10.1016/j.jvolgeores.2007.08.008>.
- Bluth, G.J., and Rose, W.I., 2002, Collaborative studies target volcanic hazards in Central America: *EOS Transactions, American Geophysical Union*, v. 83, p. 429–435, <http://dx.doi.org/10.1029/2002EO000309>.
- Bohrson, W.A., and Spera, F.J., 2007, Energy-constrained recharge, assimilation, and fractional crystallization (EC-RAXFC): A Visual Basic computer code for calculating trace element and isotope variations of open-system magmatic systems: *Geochemistry, Geophysics, Geosystems*, v. 8, Q11003, <http://dx.doi.org/10.1029/2007GC001781>.
- Bolge, L.L., Carr, M.J., Milidakis, K.I., Lindsay, F.N., and Feigenson, M.D., 2009, Correlating geochemistry, tectonics, and volcanic volume along the Central American volcanic front: *Geochemistry, Geophysics, Geosystems*, v. 10, Q12S18, <http://dx.doi.org/10.1029/2009GC002704>.
- Bromiley, G.D., and Redfern, S.A.T., 2008, The role of TiO<sub>2</sub> phases during melting of subduction-modified crust: Implications for deep mantle melting: *Earth and Planetary Science Letters*, v. 267, p. 301–308, <http://dx.doi.org/10.1016/j.epsl.2007.11.033>.
- Burkart, B., and Self, S., 1985, Extension and rotation of crustal blocks in northern Central America and effect on the volcanic arc: *Geology*, v. 13, p. 22–26, [http://dx.doi.org/10.1130/0091-7613\(1985\)13<22:EAROCB>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(1985)13<22:EAROCB>2.0.CO;2).
- Carr, M.J., 1984, Symmetrical and segmented variation of physical and geochemical characteristics of the Central American volcanic front: *Journal of Volcanology and Geothermal Research*, v. 20, p. 231–252, [http://dx.doi.org/10.1016/0377-0273\(84\)90041-6](http://dx.doi.org/10.1016/0377-0273(84)90041-6).
- Carr, M.J., Rose, W.I., and Stoiber, R.E., 1982, Central America, in Thorpe, R.S., ed., *Andesites*: John Wiley, p. 149–166.
- Carr, M.J., Feigenson, M.D., and Bennett, E.A., 1990, Incompatible element and isotopic evidence for tectonic control of source mixing and melt extraction along the Central American arc: *Contributions to Mineralogy and Petrology*, v. 105, p. 369–380, <http://dx.doi.org/10.1007/BF00286825>.
- Carr, M.J., Feigenson, M.D., Patino, L.C., and Walker, J.A., 2003, Volcanism and geochemistry in Central America: Progress and problems, in Eiler, J., ed., *Inside the Subduction Factory*: American Geophysical Union, Geophysical Monograph Series, v. 138, p. 153–174, <http://dx.doi.org/10.1029/138GM09>.
- Carr, M.J., Saginor, I., Alvarado, G.E., Bolge, L.L., Lindsay, F.N., Milidakis, K., Turrin, B.D., Feigenson, M.D., and Swisher, C.C., III, 2007a, Element fluxes from the volcanic front of Nicaragua and Costa Rica: *Geochemistry, Geophysics, Geosystems*, v. 8, Q06001, <http://dx.doi.org/10.1029/2006GC001396>.
- Carr, M.J., Patino, L.C., and Feigenson, M.D., 2007b, Petrology and geochemistry of lavas, in Bundschuh, J., and Alvarado, G.E., eds., *Central America: Geology, Resources and Hazards*, v. 1, Taylor and Francis, p. 565–590.
- Deering, C.D., Vogel, T.A., Patino, L.C., Szymanski, D.W., and Alvarado, G.E., 2012, Magmatic processes that generate chemically distinct silicic magmas in NW Costa Rica and the evolution of juvenile continental crust in oceanic arcs: *Contributions to Mineralogy and Petrology*, v. 163, p. 259–275, <http://dx.doi.org/10.1007/s00410-011-0670-z>.
- Defant, M.J., and Drummond, M.S., 1990, Derivation of some modern arc magmas by melting of young subducted lithosphere: *Nature*, v. 347, p. 662–665, <http://dx.doi.org/10.1038/347662a0>.
- DeMets, C., 2001, A new estimate for present-day Cocos-Caribbean plate motion: Implications for slip along the Central American volcanic arc: *Geophysical Research Letters*, v. 28, p. 4043–4046, <http://dx.doi.org/10.1029/2001GL013518>.
- Dengo, G., 1985, Mid America: Tectonic setting for the Pacific margin from southern Mexico to northwestern Colombia, in Nairn, A.E.M., Stehli, F.G., and Uyeda, S., eds., *The Ocean Basins and Margins: Volume 7A The Pacific Ocean*: Plenum Press, New York, p. 123–180, [http://dx.doi.org/10.1007/978-1-4613-2351-8\\_4](http://dx.doi.org/10.1007/978-1-4613-2351-8_4).
- Denyer, P., and Arias, O., 1991, Estratigrafía de la región central de Costa Rica: *Revista Geológica de América Central*, v. 12, p. 1–59.
- Denyer, P., and Gazel, E., 2009, The Costa Rican Jurassic to Miocene oceanic complexes: Origin, tectonics, and relations: *Journal of South American Earth Sciences*, v. 28, p. 429–442, <http://dx.doi.org/10.1016/j.jsames.2009.04.010>.
- Dille, M., Chen, R.S., Deichmann, U., Lerner-Lam, A.L., and Arnold, M., 2005, Natural disaster hotspots: A global risk analysis: *World Bank Publications*, Washington D.C., 145 p., <http://dx.doi.org/10.1596/0-8213-5930-4>.
- Dollfus, A., and Montserrat, E., 1868, *Voyage Géologique Dans Les Républiques De Guatemala Et De El Salvador*: Imprimerie Imp, Paris, 539 p.
- Drummond, M.S., and Defant, M.J., 1990, A model for trondhjemite-tonalite-dacite genesis and crustal growth via slab melting: Archean to modern comparisons: *Journal of Geophysical Research*, v. 95, p. 21503–21521, <http://dx.doi.org/10.1029/JB095iB13p21503>.
- Dufek, J., and Bergantz, G.W., 2005, Lower crustal magma genesis and preservation: a stochastic framework for the evaluation of basalt–crust interaction: *Journal of Petrology*, v. 46, p. 2167–2195, <http://dx.doi.org/10.1093/petrology/egi049>.
- Eiler, J.M., Carr, M.J., Reagan, M., and Stolper, E., 2005, Oxygen isotope constraints on the sources of Central American arc lavas: *Geochemistry, Geophysics, Geosystems*, v. 6, Q07007, <http://dx.doi.org/10.1029/2004GC000804>.
- Elkins, L.J., Fischer, T.P., Hilton, D.R., Sharp, Z.D., McKnight, S., and Walker, J., 2006, Tracing nitrogen in volcanic and geothermal volatiles from the Nicaraguan volcanic front: *Geochimica et Cosmochimica Acta*, v. 70, p. 5215–5235, <http://dx.doi.org/10.1016/j.gca.2006.07.024>.
- Elliott, T., 2003, Tracers of the slab, in Eiler, J., ed., *Inside the Subduction Factory*: American Geophysical Union, Geophysical Monograph Series, v. 138, p. 23–45, <http://dx.doi.org/10.1029/138GM03>.
- Escobar-Wolf, R.P., Diehl, J.F., Singer, B.S., and Rose, W.I., 2010, <sup>40</sup>Ar/<sup>39</sup>Ar and paleomagnetic constraints on the evolution of Volcán de Santa María, Guatemala: *Geological Society of America Bulletin*, v. 122, p. 757–771, <http://dx.doi.org/10.1130/B26569.1>.
- Ewert, J.W., and Harpel, C.J., 2004, In harm's way: Population and volcanic risk: *Geotimes*, v. 49, p. 14–17.



- Feigenson, M.D., and Carr, M.J., 1993, The source of Central American lavas: inferences from geochemical inverse modeling: *Contributions to Mineralogy and Petrology*, v. 113, p. 226–235, <http://dx.doi.org/10.1007/BF00283230>.
- Feigenson, M.D., Carr, M.J., Maharaj, S.V., Juliano, S., and Bolge, L.L., 2004, Lead isotope composition of Central American volcanoes: Influence of the Galapagos plume: *Geochemistry, Geophysics, Geosystems*, v. 5, Q06001, <http://dx.doi.org/10.1029/2003GC000621>.
- Fischer, T.P., Hilton, D.R., Zimmer, M.M., Shaw, A.M., Sharp, Z.D., and Walker, J.A., 2002, Subduction and recycling of nitrogen along the Central American margin: *Science*, v. 297, p. 1154–1157, <http://dx.doi.org/10.1126/science.1073995>.
- Foley, S.F., Barth, M.G., and Jenner, G.A., 2000, Rutile/melt partition coefficients for trace elements and an assessment of the influence of rutile on the trace element characteristics of subduction zone magmas: *Geochimica et Cosmochimica Acta*, v. 64, p. 933–938, [http://dx.doi.org/10.1016/S0016-7037\(99\)00355-5](http://dx.doi.org/10.1016/S0016-7037(99)00355-5).
- Freundt, A., Hartmann, A., Kutterolf, S., and Strauch, W., 2010, Volcaniclastic stratigraphy of the Tiscapa maar crater walls (Managua, Nicaragua): implications for volcanic and seismic hazards and Holocene climate changes: *International Journal of Earth Sciences*, v. 99, p. 1453–1470, <http://dx.doi.org/10.1007/s00531-009-0469-6>.
- Garrison, J.M., Reagan, M.K., and Sims, K.W.W., 2012, Dacite formation at Ilopango Caldera, El Salvador: U-series disequilibrium and implications for petrogenetic processes and magma storage time: *Geochemistry, Geophysics, Geosystems*, v. 13, Q06018, <http://dx.doi.org/10.1029/2012GC004107>.
- Gazel, E., Carr, M.J., Hoernle, K., Feigenson, M.D., Szymanski, D., Hauff, F., and van den Bogaard, P., 2009, Galapagos-OIB signature in southern Central America: Mantle refertilization by arc-hot spot interaction: *Geochemistry, Geophysics, Geosystems*, v. 10, Q02S11, <http://dx.doi.org/10.1029/2008GC002246>.
- Gazel, E., Hoernle, K., Carr, M.J., Herzberg, C., Saginor, I., van den Bogaard, P., Hauff, F., Feigenson, M., and Swisher, C., III, 2011, Plume-subduction interaction in southern Central America: Mantle upwelling and slab melting: *Lithos*, v. 121, p. 117–134, <http://dx.doi.org/10.1016/j.lithos.2010.10.008>.
- Geldmacher, J., Hoernle, K., Van Den Bogaard, P., Hauff, F., and Klügel, A., 2008, Age and geochemistry of the Central American forearc basement (DSDP Leg 67 and 84): Insights into Mesozoic arc volcanism and seamount accretion on the fringe of the Caribbean LIP: *Journal of Petrology*, v. 49, p. 1781–1815, <http://dx.doi.org/10.1093/petrology/egn046>.
- Gill, J.B., 1981, *Orogenic Andesites and Plate Tectonics*: Springer-Verlag, Berlin, 390 p., <http://dx.doi.org/10.1007/978-3-642-68012-0>.
- Goss, A.R., and Kay, S.M., 2006, Steep REE patterns and enriched Pb isotopes in southern Central American arc magmas: Evidence for forearc subduction erosion?: *Geochemistry, Geophysics, Geosystems*, v. 7, Q05016, <http://dx.doi.org/10.1029/2005GC001163>.
- Grevenmeyer, I., Kaul, N., Diaz-Naveas, J.L., Villinger, H.W., Ranero, C.R., and Reichert, C., 2005, Heat flow and bending-related faulting at subduction trenches: Case studies offshore of Nicaragua and Central Chile: *Earth and Planetary Science Letters*, v. 236, p. 238–248, <http://dx.doi.org/10.1016/j.epsl.2005.04.048>.
- Grevenmeyer, I., Ranero, C.R., Flueh, E.R., Kläschen, D., and Bialas, J., 2007, Passive and active seismological study of bending-related faulting and mantle serpentinization at the Middle America trench: *Earth and Planetary Science Letters*, v. 258, p. 528–542, <http://dx.doi.org/10.1016/j.epsl.2007.04.013>.
- Grove, T.L., Till, C.B., and Krawczynski, M.J., 2012, The role of H<sub>2</sub>O in subduction zone magmatism: *Annual Review of Earth and Planetary Sciences*, v. 40, p. 413–439, <http://dx.doi.org/10.1146/annurev-earth-042711-105310>.
- Guzmán-Speziale, M., 2001, Active seismic deformation in the grabens of northern Central America and its relationship to the relative motion of the North America–Caribbean plate boundary: *Tectonophysics*, v. 337, p. 39–51, [http://dx.doi.org/10.1016/S0040-1951\(01\)00110-X](http://dx.doi.org/10.1016/S0040-1951(01)00110-X).
- Hacker, B.R., Kelemen, P.B., and Behn, M.D., 2011, Differentiation of the continental crust by relamination: *Earth and Planetary Science Letters*, v. 307, p. 501–516, <http://dx.doi.org/10.1016/j.epsl.2011.05.024>.
- Hauff, F., Hoernle, K., Schmincke, H.-U., and Werner, R., 1997, A Mid Cretaceous origin for the Galapagos hotspot: volcanological, petrological and geochemical evidence from Costa Rican oceanic crustal segments: *Geologische Rundschau*, v. 86, p. 141–155, <http://dx.doi.org/10.1007/PL00009938>.
- Hauff, F., Hoernle, K., van den Bogaard, P., Alvarado, G., and Garbe-Schönberg, D., 2000, Age and geochemistry of basaltic complexes in western Costa Rica: Contributions to the tectonic evolution of Central America: *Geochemistry, Geophysics, Geosystems*, v. 1, 1009, <http://dx.doi.org/10.1029/1999GC000020>.
- Hayes, J.L., Holbrook, W.S., Lizarralde, D., Van Avendonk, H.J.A., Bullock, A.D., Mora, M., Harder, S., Alvarado, G.E., and Ramirez, C., 2013, Crustal structure across the Costa Rican volcanic arc: *Geochemistry, Geophysics, Geosystems*, v. 14, p. 1087–1103, <http://dx.doi.org/10.1002/ggge.20079>.
- Hermann, J., and Spandler, C.J., 2008, Sediment melts at sub-arc depths: an experimental study: *Journal of Petrology*, v. 49, p. 717–740, <http://dx.doi.org/10.1093/petrology/egm073>.
- Herrstrom, E.A., Reagan, M.K., and Morris, J.D., 1995, Variations in lava composition associated with flow of asthenosphere beneath southern Central America: *Geology*, v. 23, p. 617–620, [http://dx.doi.org/10.1130/0091-7613\(1995\)023<0617:VILCAW>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1995)023<0617:VILCAW>2.3.CO;2).
- Hildreth, W., and Moorbath, S., 1988, Crustal contributions to arc magmatism in the Andes of Central Chile: *Contributions to Mineralogy and Petrology*, v. 98, p. 455–489, <http://dx.doi.org/10.1007/BF00372365>.
- Hoernle, K., Werner, R., Morgan, J.P., Garbe-Schönberg, D., Bryce, J., and Mrazek, J., 2000, Existence of complex spatial zonation in the Galapagos plume: *Geology*, v. 28, p. 435–438, [http://dx.doi.org/10.1130/0091-7613\(2000\)28<435:EOCSZI>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2000)28<435:EOCSZI>2.0.CO;2).
- Hoernle, K., van den Bogaard, P., Werner, R., Lissinna, B., Hauff, F., Alvarado, G., and Garbe-Schönberg, D., 2002, Missing history (16–71 Ma) of the Galapagos hotspot: Implications for the tectonic and biological evolution

- of the Americas: *Geology*, v. 30, p. 795–798, [http://dx.doi.org/10.1130/0091-7613\(2002\)030<0795:MHMOTG>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2002)030<0795:MHMOTG>2.0.CO;2).
- Hoernle, K., Abt, D.L., Fischer, K.M., Nichols, H., Hauff, F., Abers, G.A., van den Bogaard, P., Heydolph, K., Alvarado, G., Protti, M., and Strauch, W., 2008, Arc-parallel flow in the mantle wedge beneath Costa Rica and Nicaragua: *Nature*, v. 451, p. 1094–1097, <http://dx.doi.org/10.1038/nature06550>.
- Ivancic, M., Grevemeyer, I., Berhorst, A., Flueh, E.R., and McIntosh, K., 2008, Impact of bending related faulting on the seismic properties of the incoming oceanic plate offshore of Nicaragua: *Journal of Geophysical Research*, v. 113, B05410, <http://dx.doi.org/10.1029/2007JB005291>.
- Ivancic, M., Grevemeyer, I., Bialas, J., and Petersen, C.J., 2010, Serpentinization in the trench-outer rise region offshore of Nicaragua: constraints from seismic refraction and wide-angle data: *Geophysical Journal International*, v. 180, p. 1253–1264, <http://dx.doi.org/10.1111/j.1365-246X.2009.04474.x>.
- Jenner, G.A., Foley, S.F., Jackson, S.E., Green, T.H., Fryer, B.J., and Longgerich, H.P., 1993, Determination of partition coefficients for trace elements in high pressure-temperature experimental run products by laser ablation microprobe-inductively coupled plasma-mass spectrometry (LAM-ICP-MS): *Geochimica et Cosmochimica Acta*, v. 57, p. 5099–5103, [http://dx.doi.org/10.1016/0016-7037\(93\)90611-Y](http://dx.doi.org/10.1016/0016-7037(93)90611-Y).
- Jicha, B.R., Smith, K.E., Singer, B.S., Beard, B.L., Johnson, C.M., and Rogers, N.W., 2010, Crustal assimilation no match for slab fluids beneath Volcán de Santa María, Guatemala: *Geology*, v. 38, p. 859–862, <http://dx.doi.org/10.1130/G31062.1>.
- Johnson, M.C., and Plank, T., 1999, Dehydration and melting experiments constrain the fate of subducted sediments: *Geochimica Geophysica Geosystems*, v. 1, 1007, <http://dx.doi.org/10.1029/1999GC00014>.
- Johnston, S.T., and Thorkelson, D.J., 1997, Cocos-Nazca slab window beneath Central America: *Earth and Planetary Science Letters*, v. 146, p. 465–474, [http://dx.doi.org/10.1016/S0012-821X\(96\)00242-7](http://dx.doi.org/10.1016/S0012-821X(96)00242-7).
- Kay, R.W., 1978, Aleutian magnesian andesites: Melts from subducted Pacific ocean crust: *Journal of Volcanology and Geothermal Research*, v. 4, p. 117–132, [http://dx.doi.org/10.1016/0377-0273\(78\)90032-X](http://dx.doi.org/10.1016/0377-0273(78)90032-X).
- Keppeler, H., 1996, Constraints from partitioning experiments on the composition of subduction-zone fluids: *Nature*, v. 380, p. 237–240, <http://dx.doi.org/10.1038/380237a0>.
- Key, K., Constable, S., Matsuno, T., Evans, R.L., and Myer, D., 2012, Electromagnetic detection of plate hydration due to bending faults at the Middle America trench: *Earth and Planetary Science Letters*, v. 351–352, p. 45–53, <http://dx.doi.org/10.1016/j.epsl.2012.07.020>.
- Kimura, G., Silver, E., and Blum, P., 1997, *Proceedings of the Ocean Drilling Program, Initial Reports, Volume 170: Ocean Drilling Program, College Station, Texas*, 458 p.
- Klemme, S., Prowatke, S., Hametner, K., and Günther, D., 2005, Partitioning of trace elements between rutile and silicate melts: Implications for subduction zones: *Geochimica et Cosmochimica Acta*, v. 69, p. 2361–2371, <http://dx.doi.org/10.1016/j.gca.2004.11.015>.
- Kutterolf, S., Freundt, A., Pérez, W., Mörz, T., Schacht, U., Wehrmann, H., and Schmincke, H.-U., 2008a, Pacific offshore record of plinian arc volcanism in Central America: 1. Along-arc correlations: *Geochimica, Geophysics, Geosystems*, v. 9, Q02S01, <http://dx.doi.org/10.1029/2007GC001631>.
- Kutterolf, S., Freundt, A., and Pérez, W., 2008b, Pacific offshore record of plinian arc volcanism in Central America: 2. Tephra volumes and erupted masses: *Geochimica, Geophysics, Geosystems*, v. 9, Q02S02, <http://dx.doi.org/10.1029/2007GC001791>.
- La Femina, P.C., Connor, C.B., Hill, B.E., Strauch, W., and Saballos, J.A., 2004, Magma-tectonic interactions in Nicaragua: the 1999 seismic swarm and eruption of Cerro Negro volcano: *Journal of Volcanology and Geothermal Research*, v. 137, p. 187–199, <http://dx.doi.org/10.1016/j.jvolgeores.2004.05.006>.
- Leeman, W.P., Carr, M.J., and Morris, J.D., 1994, Boron geochemistry of the Central American volcanic arc: Constraints on the genesis of subduction-related magmas: *Geochimica et Cosmochimica Acta*, v. 58, p. 149–168, [http://dx.doi.org/10.1016/0016-7037\(94\)90453-7](http://dx.doi.org/10.1016/0016-7037(94)90453-7).
- Lefeldt, M., Grevemeyer, I., Goßler, J., and Bialas, J., 2009, Intraplate seismicity and related mantle hydration at the Nicaraguan trench outer rise: *Geophysical Journal International*, v. 178, p. 742–752, <http://dx.doi.org/10.1111/j.1365-246X.2009.04167.x>.
- Lücke, O.H., Götze, H.-J., and Alvarado, G.E., 2010, A constrained 3D density model of the upper crust from gravity data interpretation for central Costa Rica: *International Journal of Geophysics*, v. 2010, 860902, <http://dx.doi.org/10.1155/2010/860902>.
- Lyon-Caen, H., Barrier, E., Lasserre, C., Franco, A., Arzu, I., Chiquin, L., Chiquin, M., Duquesnoy, T., Flores, O., Galicia, O., Luna, J., Molina, E., Porras, O., Requena, J., Robles, V., Romero, J., and Wolf, R., 2006, Kinematics of the North American–Caribbean–Cocos plates in Central America from new GPS measurements across the Polochic–Motagua fault system: *Geophysical Research Letters*, v. 33, L19309, <http://dx.doi.org/10.1029/2006GL027694>.
- MacKenzie, L.S., Abers, G.A., Rondenay, S., and Fischer, K.M., 2010, Imaging a steeply dipping subducting slab in southern Central America: *Earth and Planetary Science Letters*, v. 296, p. 459–468, <http://dx.doi.org/10.1016/j.epsl.2010.05.033>.
- MacMillan, I., Gans, P.B., and Alvarado, G., 2004, Middle Miocene to present plate tectonic history of the southern Central American Volcanic Arc: *Tectonophysics*, v. 392, p. 325–348, <http://dx.doi.org/10.1016/j.tecto.2004.04.014>.
- Mann, P., Rogers, R.D., and Gahagan, L., 2007, Overview of plate tectonic history and its unresolved problems, *in* Bundschuh, J., and Alvarado, G.E., eds., *Central America: Geology, Resources and Hazards*, v. 1: Taylor & Francis, p. 201–237.
- Marschall, H.R., Altherr, R., and Rüpke, L., 2007, Squeezing out the slab – modelling the release of Li, Be and B during progressive high-pressure metamorphism: *Chemical Geology*, v. 239, p. 323–335, <http://dx.doi.org/10.1016/j.chemgeo.2006.08.008>.
- Martin, L.A.J., Wood, B.J., Turner, S., and Rushmer, T., 2011, Experimental measurements of trace element partitioning between lawsonite, zoisite and fluid and their implication for the composition of arc magmas: *Journal of Petrology*, v. 52, p. 1049–1075, <http://dx.doi.org/10.1093/petrology>

- gy/egr018.
- McBirney, A.R., 1969, Compositional variations in Cenozoic calc-alkaline suites of Central America, *in* McBirney, A.R., *ed.*, Proceedings of the Andesite Conference: Oregon Department of Geology and Mineral Industries, Bulletin, 65, p. 185–189.
- Moritz, E., Bornholdt, S., Westphal, H., and Meschede, M., 2000, Neural network interpretation of LWD data (ODP Leg 170) confirms complete sediment subduction at the Costa Rica convergent margin: *Earth and Planetary Science Letters*, v. 174, p. 301–312, [http://dx.doi.org/10.1016/S0012-821X\(99\)00270-8](http://dx.doi.org/10.1016/S0012-821X(99)00270-8).
- Morris, J.D., and Hart, S.R., 1983, Isotopic and incompatible element constraints on the genesis of island arc volcanics from Cold Bay and Amak Island, Aleutians, and implications for mantle structure: *Geochimica et Cosmochimica Acta*, v. 47, p. 2015–2030, [http://dx.doi.org/10.1016/0016-7037\(83\)90217-X](http://dx.doi.org/10.1016/0016-7037(83)90217-X).
- Morris, J.D., Leeman, W.P., and Tera, F., 1990, The subducted component in island arc lavas: constraints from Be isotopes and B–Be systematics: *Nature*, v. 344, p. 31–36, <http://dx.doi.org/10.1038/344031a0>.
- Niida, K., and Green, D.H., 1999, Stability and chemical composition of pargasitic amphibole in MORB pyroxene under upper mantle conditions: *Contributions to Mineralogy and Petrology*, v. 135, p. 18–40, <http://dx.doi.org/10.1007/s004100050495>.
- O'Connor, J.M., Stoffers, P., Wijbrans, J.R., and Worthington, T.J., 2007, Migration of widespread long-lived volcanism across the Galápagos Volcanic Province: Evidence for a broad hotspot melting anomaly?: *Earth and Planetary Science Letters*, v. 263, p. 339–354, <http://dx.doi.org/10.1016/j.epsl.2007.09.007>.
- Patino, L.C., Carr, M.J., and Feigenson, M.D., 2000, Local and regional variations in Central American arc lavas controlled by variations in subducted sediment input: *Contributions to Mineralogy and Petrology*, v. 138, p. 265–283, <http://dx.doi.org/10.1007/s004100050562>.
- Plank, T., and Langmuir, C.H., 1988, An evaluation of the global variations in the major element chemistry of arc basalts: *Earth and Planetary Science Letters*, v. 90, p. 349–370, [http://dx.doi.org/10.1016/0012-821X\(88\)90135-5](http://dx.doi.org/10.1016/0012-821X(88)90135-5).
- Plank, T., and Langmuir, C.H., 1993, Tracing trace elements from sediment input to volcanic output at subduction zones: *Nature*, v. 362, p. 739–742, <http://dx.doi.org/10.1038/362739a0>.
- Plank, T., and Langmuir, C.H., 1998, The chemical composition of subducting sediment and its consequences for the crust and mantle: *Chemical Geology*, v. 145, p. 325–394, [http://dx.doi.org/10.1016/S0009-2541\(97\)00150-2](http://dx.doi.org/10.1016/S0009-2541(97)00150-2).
- Protti, M., Güendel, F., and McNally, K., 1994, The geometry of the Wadati–Benioff zone under southern Central America and its tectonic significance: results from a high-resolution local seismographic network: *Physics of the Earth and Planetary Interiors*, v. 84, p. 271–287, [http://dx.doi.org/10.1016/0031-9201\(94\)90046-9](http://dx.doi.org/10.1016/0031-9201(94)90046-9).
- Protti, M., Güendel, F., and McNally, K., 1995, Correlation between the age of the subducting Cocos plate and the geometry of the Wadati–Benioff zone under Nicaragua and Costa Rica, *in* Mann, P., *ed.*, *Geologic and Tectonic Development of the Caribbean Plate Boundary in Southern Central America*: Geological Society of America, Special Papers, v. 295, p. 309–326, <http://dx.doi.org/10.1130/SPE295-p309>.
- Ranero, C.R., and von Huene, R., 2000, Subduction erosion along the Middle America convergent margin: *Nature*, v. 404, p. 748–752, <http://dx.doi.org/10.1038/35008046>.
- Ranero, C.R., Morgan, J.P., McIntosh, K.D., and Reichert, C., 2001, Flexural faulting and mantle serpentinization at the Middle America trench (abstract): *American Geophysical Union, Fall Meeting, 2001, Abstracts*, v. 82, #T22D-04.
- Ranero, C.R., Morgan, J.P., McIntosh, K., and Reichert, C., 2003, Bending-related faulting and mantle serpentinization at the Middle America trench: *Nature*, v. 425, p. 367–373, <http://dx.doi.org/10.1038/nature01961>.
- Rapp, R.P., and Watson, E.B., 1995, Dehydration melting of metabasalt at 8–32 kbar: Implications for continental growth and crust–mantle recycling: *Journal of Petrology*, v. 36, p. 891–931, <http://dx.doi.org/10.1093/petrology/36.4.891>.
- Rausch, J., and Schmincke, H.-U., 2010, Nejapa Tephra: The youngest (c. 1 ka BP) highly explosive hydroclastic eruption in western Managua (Nicaragua): *Journal of Volcanology and Geothermal Research*, v. 192, p. 159–177, <http://dx.doi.org/10.1016/j.jvolgeores.2010.02.010>.
- Reagan, M.K., and Gill, J.B., 1989, Coexisting calcalkaline and high-niobium basalts from Turrialba Volcano, Costa Rica: Implications for residual titanates in arc magma sources: *Journal of Geophysical Research*, v. 94, p. 4619–4633, <http://dx.doi.org/10.1029/JB094iB04p04619>.
- Reagan, M.K., Morris, J.D., Herrstrom, E.A., and Murrell, M.T., 1994, Uranium series and beryllium isotope evidence for an extended history of subduction modification of the mantle below Nicaragua: *Geochimica et Cosmochimica Acta*, v. 58, p. 4199–4212, [http://dx.doi.org/10.1016/0016-7037\(94\)90273-9](http://dx.doi.org/10.1016/0016-7037(94)90273-9).
- Ringwood, A.E., 1974, The petrological evolution of island arc systems: Twenty-seventh William Smith Lecture: *Journal of the Geological Society*, v. 130, p. 183–204, <http://dx.doi.org/10.1144/gsjgs.130.3.0183>.
- Rodriguez, M., DeMets, C., Rogers, R., Tenorio, C., and Hernandez, D., 2009, A GPS and modelling study of deformation in northern Central America: *Geophysical Journal International*, v. 178, p. 1733–1754, <http://dx.doi.org/10.1111/j.1365-246X.2009.04251.x>.
- Rogers, R.D., and Mann, P., 2007, Transtensional deformation of the western Caribbean–North America plate boundary zone, *in* Mann, P., *ed.*, *Geologic and Tectonic Development of the Caribbean Plate Boundary in Northern Central America*: Geological Society of America, Special Papers, v. 428, p. 37–64, [http://dx.doi.org/10.1130/2007.2428\(03\)](http://dx.doi.org/10.1130/2007.2428(03)).
- Rogers, R.D., Mann, P., and Emmet, P.A., 2007, Tectonic terranes of the Chortis block based on integration of regional aeromagnetic and geologic data, *in* Mann, P., *ed.*, *Geologic and Tectonic Development of the Caribbean Plate Boundary in Northern Central America*: Geological Society of America, Special Papers, v. 428, p. 65–88, [http://dx.doi.org/10.1130/2007.2428\(04\)](http://dx.doi.org/10.1130/2007.2428(04)).
- Rose, W.I., Conway, F.M., Pullinger, C.R., Deino, A., and McIntosh, W.C., 1999, An improved age framework for late Quaternary silicic eruptions in northern Central America: *Bulletin of Volcanology*, v. 61, p. 106–120, <http://dx.doi.org/10.1007/s004450050266>.
- Rüpke, L.H., Morgan, J.P., Hort, M., and Connolly, J.A.D., 2002, Are the region-



- al variations in Central American arc lavas due to differing basaltic versus peridotitic slab sources of fluids?: *Geology*, v. 30, p. 1035–1038, [http://dx.doi.org/10.1130/0091-7613\(2002\)030<1035:ATRVIC>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(2002)030<1035:ATRVIC>2.0.CO;2).
- Rychert, C.A., Fischer, K.M., Abers, G.A., Plank, T., Syracuse, E., Protti, J.M., Gonzalez, V., and Strauch, W., 2008, Strong along-arc variations in attenuation in the mantle wedge beneath Costa Rica and Nicaragua: *Geochemistry, Geophysics, Geosystems*, v. 9, Q10S10, <http://dx.doi.org/10.1029/2008GC002040>.
- Sadofsky, S.J., Portnyagin, M., Hoernle, K., and van den Bogaard, P., 2008, Subduction cycling of volatiles and trace elements through the Central American volcanic arc: evidence from melt inclusions: *Contributions to Mineralogy and Petrology*, v. 155, p. 433–456, <http://dx.doi.org/10.1007/s00410-007-0251-3>.
- Sallarès, V., Dañobeitia, J.J., and Flueh, E.R., 2001, Lithospheric structure of the Costa Rican Isthmus: Effects of subduction zone magmatism on an oceanic plateau: *Journal of Geophysical Research*, v. 106, p. 621–643, <http://dx.doi.org/10.1029/2000JB900245>.
- Sapper, K., 1917, *Katalog Der Geschichtlichen Vulkanausbrüche*: Karl J. Trubner, Strasbourg, 358 p.
- Shaw, A.M., Hilton, D.R., Fischer, T.P., Walker, J.A., and Alvarado, G.E., 2003, Contrasting He–C relationships in Nicaragua and Costa Rica: insights into C cycling through subduction zones: *Earth and Planetary Science Letters*, v. 214, p. 499–513, [http://dx.doi.org/10.1016/S0012-821X\(03\)00401-1](http://dx.doi.org/10.1016/S0012-821X(03)00401-1).
- Silver, E., Costa Pisani, P., Hutnak, M., Fisher, A., DeShon, H., and Taylor, B., 2004, An 8–10 Ma tectonic event on the Cocos Plate offshore Costa Rica: Result of Cocos Ridge collision?: *Geophysical Research Letters*, v. 31, L18601, <http://dx.doi.org/10.1029/2004GL020272>.
- Singer, B.S., Smith, K.E., Jicha, B.R., Beard, B.L., Johnson, C.M., and Rogers, N.W., 2011, Tracking open-system differentiation during growth of Santa María Volcano, Guatemala: *Journal of Petrology*, v. 52, p. 2335–2363, <http://dx.doi.org/10.1093/petrology/egr047>.
- Sinton, C.W., Duncan, R.A., Storey, M., Lewis, J., and Estrada, J.J., 1998, An oceanic flood basalt province within the Caribbean plate: *Earth and Planetary Science Letters*, v. 155, p. 221–235, [http://dx.doi.org/10.1016/S0012-821X\(97\)00214-8](http://dx.doi.org/10.1016/S0012-821X(97)00214-8).
- Skora, S., and Blundy, J., 2010, High-pressure hydrous phase relations of radiolarian clay and implications for the involvement of subducted sediment in arc magmatism: *Journal of Petrology*, v. 51, p. 2211–2243, <http://dx.doi.org/10.1093/petrology/egq054>.
- Stoiber, R.E., and Carr, M.J., 1973, Quaternary volcanic and tectonic segmentation of Central America: *Bulletin Volcanologique*, v. 37, p. 304–325, <http://dx.doi.org/10.1007/BF02597631>.
- Syracuse, E.M., and Abers, G.A., 2006, Global compilation of variations in slab depth beneath arc volcanoes and implications: *Geochemistry, Geophysics, Geosystems*, v. 7, Q05017, <http://dx.doi.org/10.1029/2005GC001045>.
- Syracuse, E.M., Abers, G.A., Fischer, K., MacKenzie, L., Rychert, C., Protti, M., González, V., and Strauch, W., 2008, Seismic tomography and earthquake locations in the Nicaraguan and Costa Rican upper mantle: *Geochemistry, Geophysics, Geosystems*, v. 9, Q07S08, <http://dx.doi.org/10.1029/2008GC001963>.
- Tamura, Y., and Tatsumi, Y., 2002, Remelting of an andesitic crust as a possible origin for rhyolitic magma in oceanic arcs: an example from the Izu-Bonin arc: *Journal of Petrology*, v. 43, p. 1029–1047, <http://dx.doi.org/10.1093/petrology/43.6.1029>.
- Tatsumi, Y., and Isoyama, H., 1988, Transportation of beryllium with H<sub>2</sub>O at high pressures: Implication for magma genesis in subduction zones: *Geophysical Research Letters*, v. 15, p. 180–183, <http://dx.doi.org/10.1029/GL015i002p00180>.
- Tatsumi, Y., Hamilton, D.L., and Nesbitt, R.W., 1986, Chemical characteristics of fluid phase released from a subducted lithosphere and origin of arc magmas: Evidence from high-pressure experiments and natural rocks: *Journal of Volcanology and Geothermal Research*, v. 29, p. 293–309, [http://dx.doi.org/10.1016/0377-0273\(86\)90049-1](http://dx.doi.org/10.1016/0377-0273(86)90049-1).
- Tera, F., Brown, L., Morris, J., Sacks, I.S., Klein, J., and Middleton, R., 1986, Sediment incorporation in island-arc magmas: Inferences from <sup>10</sup>Be: *Geochimica et Cosmochimica Acta*, v. 50, p. 535–550, [http://dx.doi.org/10.1016/0016-7037\(86\)90103-1](http://dx.doi.org/10.1016/0016-7037(86)90103-1).
- Tiepolo, M., Bottazzi, P., Foley, S.F., Oberli, R., Vannucci, R., and Zanetti, A., 2001, Fractionation of Nb and Ta from Zr and Hf at mantle depths: the role of titanite and kaersutite: *Journal of Petrology*, v. 42, p. 221–232, <http://dx.doi.org/10.1093/petrology/42.1.221>.
- Tollstrup, D., Gill, J., Kent, A., Prinkey, D., Williams, R., Tamura, Y., and Ishizuka, O., 2010, Across-arc geochemical trends in the Izu-Bonin arc: Contributions from the subducting slab, revisited: *Geochemistry, Geophysics, Geosystems*, v. 11, Q01X10, <http://dx.doi.org/10.1029/2009GC002847>.
- Tonardini, S., Agostini, S., Doglioni, C., Innocenti, F., and Manetti, P., 2007, Evidence for serpentine fluid in convergent margin systems: The example of El Salvador (Central America) arc lavas: *Geochemistry, Geophysics, Geosystems*, v. 8, Q09014, <http://dx.doi.org/10.1029/2006GC001508>.
- Turner, S., Bourdon, B., and Gill, J., 2003, Insights into magma genesis at convergent margins from U–series isotopes, *in* Bourdon, B., Henderson, G.M., Lundstrom, C.C., and Turner, S.P., eds., *Uranium-Series Geochemistry: Mineralogical Society of America, Reviews in Mineralogy and Geochemistry*, v. 52, p. 255–310.
- Ui, T., 1972, Recent volcanism in the Masaya–Granada area, Nicaragua: *Bulletin Volcanologique*, v. 36, p. 174–190, <http://dx.doi.org/10.1007/BF02596989>.
- Ulmer, P., 2001, Partial melting in the mantle wedge – the role of H<sub>2</sub>O in the genesis of mantle-derived ‘arc-related’ magmas: *Physics of the Earth and Planetary Interiors*, v. 127, p. 215–232, [http://dx.doi.org/10.1016/S0031-9201\(01\)00229-1](http://dx.doi.org/10.1016/S0031-9201(01)00229-1).
- Van Avendonk, H.J.A., Holbrook, W.S., Lizarralde, D., and Denyer, P., 2011, Structure and serpentinization of the subducting Cocos plate offshore Nicaragua and Costa Rica: *Geochemistry, Geophysics, Geosystems*, v. 12, Q06009, <http://dx.doi.org/10.1029/2011GC003592>.
- van Wyk de Vries, B., Grosse, P., and Alvarado, G.E., 2007, Volcanism and volcanic landforms, *in* Bundschuh, J., and Alvarado, G.E., eds., *Central America: Geology, Resources and Hazards*, v. 1: Taylor and Francis, p. 123–154, <http://dx.doi.org/10.1201/9780203947043.ch4>.

- Vogel, T.A., Patino, L.C., Alvarado, G.E., and Gans, P.B., 2004, Silicic ignimbrites within the Costa Rican volcanic front: evidence for the formation of continental crust: *Earth and Planetary Science Letters*, v. 226, p. 149–159, <http://dx.doi.org/10.1016/j.epsl.2004.07.013>.
- Vogel, T.A., Patino, L.C., Eaton, J.K., Valley, J.W., Rose, W.I., Alvarado, G.E., and Viray, E.L., 2006, Origin of silicic magmas along the Central American volcanic front: Genetic relationship to mafic melts: *Journal of Volcanology and Geothermal Research*, v. 156, p. 217–228, <http://dx.doi.org/10.1016/j.jvolgeores.2006.03.002>.
- von Huene, R., Aubouin, J., Azema, J., Blackington, G., Carter, J.A., Coulbourn, W.T., Cowan, D.S., Curiale, J.A., Dengo, C.A., Faas, R.W., Harrison, W., Hesse, R., Hussong, D.M., Laad, J.W., Muzylov, N., Shiki, T., Thompson, P.R., and Westberg, J., 1980, Leg 67: The Deep Sea Drilling Project Mid-America Trench transect off Guatemala: *Geological Society of America Bulletin*, v. 91, p. 421–432, [http://dx.doi.org/10.1130/0016-7606\(1980\)91<421:LTSDP>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1980)91<421:LTSDP>2.0.CO;2).
- Walker, J.A., 1984, Volcanic rocks from the Nejapa and Granada cinder cone alignments, Nicaragua, Central America: *Journal of Petrology*, v. 25, p. 299–342, <http://dx.doi.org/10.1093/petrology/25.2.299>.
- Walker, J.A., 1989, Caribbean arc tholeiites: *Journal of Geophysical Research*, v. 94, p. 10539–10548, <http://dx.doi.org/10.1029/JB094iB08p10539>.
- Walker, J.A., Carr, M.J., Feigenson, M.D., and Kalamarides, R.I., 1990, The petrogenetic significance of interstratified high- and low-Ti basalts in central Nicaragua: *Journal of Petrology*, v. 31, p. 1141–1164, <http://dx.doi.org/10.1093/petrology/31.5.1141>.
- Walker, J.A., Patino, L.C., Cameron, B.I., and Carr, M.J., 2000, Petrogenetic insights provided by compositional transects across the Central America arc: Southeastern Guatemala and Honduras: *Journal of Geophysical Research*, v. 105, p. 18949–18963, <http://dx.doi.org/10.1029/2000JB900173>.
- Walker, J.A., Patino, L.C., Carr, M.J., and Feigenson, M.D., 2001, Slab control over HFSE depletions in central Nicaragua: *Earth and Planetary Science Letters*, v. 192, p. 533–543, [http://dx.doi.org/10.1016/S0012-821X\(01\)00476-9](http://dx.doi.org/10.1016/S0012-821X(01)00476-9).
- Walker, J.A., Mickelson, J.E., Thomas, R.B., Patino, L.C., Cameron, B., Carr, M.J., Feigenson, M.D., and Edwards, R.L., 2007, U-series disequilibria in Guatemalan lavas, crustal contamination, and implications for magma genesis along the Central American subduction zone: *Journal of Geophysical Research*, v. 112, B06205, <http://dx.doi.org/10.1029/2006JB004589>.
- Walker, J.A., Singer, B.S., Jicha, B.R., Cameron, B.I., Carr, M.J., and Olney, J.L., 2011, Monogenetic, behind-the-front volcanism in southeastern Guatemala and western El Salvador:  $^{40}\text{Ar}/^{39}\text{Ar}$  ages and tectonic implications: *Lithos*, v. 123, p. 243–253, <http://dx.doi.org/10.1016/j.lithos.2010.09.016>.
- Wallace, P.J., 2005, Volatiles in subduction zone magmas: concentrations and fluxes based on melt inclusions and volcanic gas data: *Journal of Volcanology and Geothermal Research*, v. 140, p. 217–240, <http://dx.doi.org/10.1016/j.jvolgeores.2004.07.023>.
- Werner, R., Hoernle, K., van den Bogaard, P., Ranero, C., von Huene, R., and Korich, D., 1999, Drowned 14-m.y.-old Galápagos archipelago off the coast of Costa Rica: Implications for tectonic and evolutionary models: *Geology*, v. 27, p. 499–502, [http://dx.doi.org/10.1130/0091-7613\(1999\)027<0499:DMYO-GP>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1999)027<0499:DMYO-GP>2.3.CO;2).
- Werner, R., Hoernle, K., Barckhausen, U., and Hauff, F., 2003, Geodynamic evolution of the Galápagos hot spot system (Central East Pacific) over the past 20 m.y.: Constraints from morphology, geochemistry, and magnetic anomalies: *Geochemistry, Geophysics, Geosystems*, v. 4, 1108, <http://dx.doi.org/10.1029/2003GC000576>.
- Whattam, S.A., Montes, C., McFadden, R.R., Cardona, A., Ramirez, D., and Valencia, V., 2012, Age and origin of earliest adakitic-like magmatism in Panama: Implications for the tectonic evolution of the Panamanian magmatic arc system: *Lithos*, v. 142–143, p. 226–244, <http://dx.doi.org/10.1016/j.lithos.2012.02.017>.
- Witham, C.S., 2005, Volcanic disasters and incidents: A new database: *Journal of Volcanology and Geothermal Research*, v. 148, p. 191–233, <http://dx.doi.org/10.1016/j.jvolgeores.2005.04.017>.
- Woodhead, J.D., and Johnson, R.W., 1993, Isotopic and trace-element profiles across the New Britain island arc, Papua New Guinea: *Contributions to Mineralogy and Petrology*, v. 113, p. 479–491, <http://dx.doi.org/10.1007/BF00698317>.
- Woodhead, J.D., Eggins, S.M., and Johnson, R.W., 1998, Magma genesis in the New Britain Island Arc: Further insights into melting and mass transfer processes: *Journal of Petrology*, v. 39, p. 1641–1668, <http://dx.doi.org/10.1093/petrology/39.9.1641>.
- Woodhead, J.D., Hergt, J.M., Davidson, J.P., and Eggins, S.M., 2001, Hafnium isotope evidence for ‘conservative’ element mobility during subduction zone processes: *Earth and Planetary Science Letters*, v. 192, p. 331–346, [http://dx.doi.org/10.1016/S0012-821X\(01\)00453-8](http://dx.doi.org/10.1016/S0012-821X(01)00453-8).
- Xiong, X.L., Adam, J., and Green, T.H., 2005, Rutile stability and rutile/melt HFSE partitioning during partial melting of hydrous basalt: Implications for TTG genesis: *Chemical Geology*, v. 218, p. 339–359, <http://dx.doi.org/10.1016/j.chemgeo.2005.01.014>.
- You, C.-F., Morris, J.D., Gieskes, J.M., Rosenbauer, R., Zheng, S.H., Xu, X., Ku, T.L., and Bischoff, J.L., 1994, Mobilization of beryllium in the sedimentary column at convergent margins: *Geochimica et Cosmochimica Acta*, v. 58, p. 4887–4897, [http://dx.doi.org/10.1016/0016-7037\(94\)90219-4](http://dx.doi.org/10.1016/0016-7037(94)90219-4).
- You, C.-F., Castillo, P.R., Gieskes, J.M., Chan, L.H., and Spivack, A.J., 1996, Trace element behavior in hydrothermal experiments: Implications for fluid processes at shallow depths in subduction zones: *Earth and Planetary Science Letters*, v. 140, p. 41–52, [http://dx.doi.org/10.1016/0012-821X\(96\)00049-0](http://dx.doi.org/10.1016/0012-821X(96)00049-0).

Received June 2013

Accepted as revised August 2013

First published on the web

February 2014